

[Available Issues](#) | [Japanese](#)Author: [ADVANCED](#)Volume Page Keyword: [TOP](#) > [Available Issues](#) > [Table of Contents](#) > **Abstract**

Tropical Medicine and Health

Vol. 38 (2010), No. 1 p.39

Incidence of Chikungunya Fever in Singapore: Implementation of Health Measures and Transnational Movements of I

[Minako Jen Yoshikawa](#)¹⁾, [Choon Siang Tang](#)²⁾ and [Mitsu](#)

1) Graduate School of Asian and African Area Studies, Kyoto University

2) National Environment Agency

3) Center for Southeast Asia Studies, Kyoto University

(Accepted January 20, 2010)

Abstract: In addition to Dengue virus infection, another mosquito-borne disease has become a significant health hazard. After the first domestic case of Chikungunya fever was reported in the Republic of Singapore, the outbreak was brought under control. Despite the quick and thorough public health measures including international travel restrictions, this country known as a developed island state could not stop the spread of the disease. Both domestic and imported cases surged. Our in-depth investigation and field activities recapitulates the governmental interventions to control the disease. We discuss potential factors contributing to the subsequent

Singapore and then review strategies that other governments may co
vector-borne infectious diseases in urban areas with transnational m

Key words: [Chikungunya](#), [vector control](#), [public health measures](#),
[movements](#), [Singapore](#)

[\[PDF \(55K\)\]](#) [\[References\]](#)

Downlo

To cite this article:

Minako Jen Yoshikawa, Choon Siang Tang and Mitsuaki Nishibuc
Chikungunya Fever in Singapore: Implications of Public Health Me
Movements of People”. *Tropical Medicine and Health*, Vol. **38**, pp

doi:10.2149/tmh.2009-18

JOI JST.JSTAGE/tmh/2009-18

Copyright (c) 2010 by The Japanese Society of Tropic