

Received: July 12, 2003

Accepted: November 21, 2003

Ref: ● Kaur K and Garg RK. Results Of Detection On The Persistence Of Insecticides (Dimethoate And Phosphamidon) From Soil And Paper Substrates By Thin Layer Chromatography *Anil Aggrawal's Internet Journal of Forensic Medicine and Toxicology*, 2003; Vol. 4, No. 2, (July - December 2003); ; Published: November 24, 2003, ([Accessed:](#)

● Email Dr. Kuldeep Kaur by [clicking here](#)



: **EMBASE Accession Number: 2004204909**



Kuldeep Kaur



R.K.Garg

Results Of Detection On The Persistence Of Insecticides (Dimethoate And Phosphamidon) From Soil And Paper Substrates By Thin Layer Chromatography

● **Kuldeep Kaur**
*Former, M.Sc. Student, Forensic Science
Department Punjabi University, Patiala -
147002, India*

● **R.K.Garg**
*Reader and Former, HOD Forensic Science,
Department Punjabi University, Patiala
147002, India. (Corresponding author)*

Abstract

In the present investigation dimethoate and phosphamidon pesticides presents on soil and paper material have been examined using Thin Layer Chromatography. Among the fifteen different solvent systems examined, it has been observed that dimethoate can be separated by using Benzene: Acetone (9 : 1) and phosphamidon by using cyclohexane . Acetone' chloroform (70 : 25 : 5) followed by iodine fuming as visualizing aid. The analysis of these two compounds could be successfully performed even up to 8 weeks times from soil and paper containing these pesticides in small amounts and it remains therein. The study will provide a useful information to the forensic toxicologists..

Keywords

insecticides, dimethoate, phosphamidon, thin layer chromatography

Introduction

Poisoning due to insecticides and pesticides have an important role in crimes all over the world. Forensic toxicologists are dealing with maximum number of cases' due to insecticides and pesticide poisoning involved in various types of crimes. Pesticides also cause much hazard to the surrounding environments and other organisms. Since the analysis of insecticide residue poses an entirely different type of problem for the toxicologists because these residues are present in extremely small quantity in heterogeneous materials including the biological materials. The importance of insecticide residue problem led to intensive search for analytical methods for accurate and rapid analysis (Zweig, 1964), The determination of the pesticide in various biological materials often faced with the problem of determining the minute quantity of insecticides mixed with large amount of extraneous material or intermixing. Qualitative and quantitative methods are required to be applied keeping in view the sensitivity and specificity of the methods on one hand and nature of type of pesticides on the other hand. The technique of thin layer chromatography has made a strong impact on analytical toxicology. Extensive literature on both the qualitative and quantitative analysis of insecticides/pesticides is available (Beynon and Elgar, 1966 ; Fisher and Vanwazar, 1961 ; Cautselinis and Dinvopoulos, 1971 ; Stanley, 1964 ; Abbott et al. 1964 ; Tewari, 1976 ; Ganguly and Bhattacharya, 1975 ; Das, 1977 ; Kaur, 1998 ; Grant et al. 1969 ; Guru et al. 1981 ; Dewey, 1958 ; Stewart and Stolman, 1960, 1961 ; Gunther and Blinn, 1963 ; Zweig, 1964 ; KawaJe and Joglekar, 1968, 1976 ; Fargo, 1967 ; Prasad et al. 1982 ; Clarke, 1971 ; Bami, 1972, 1981 ; Tewari, 1976 ; Kirchner, 1978 ; 1973b. Tewari et al., 1981 ; Gouldon et al. 1965) Tewari and Ram, 1969 ; Wong and Chou, 1969 ; Narayanswami et al., 1974; Sandroni and Schlitt, 1971 ; Stahl, 1969; Ludwick et. al., 1977, Bunyan, 1964 ; Chakravarti and Roy, 1976 ; Joglekar and MahaI, 1968 ; Kulkarni and Ramakrishna, 1972).

In the present investigation an attempt has been made to investigate the persistence of two different organophosphorus compounds present on two different substrates (soil and paper) after a varied length of time (upto 8 weeks) using thin layer chromatography technique applying different solvent systems.

Materials And Methods

The present investigation has been conducted on two commercial grade organophosphorus compounds namely Dimethoate and Phosphamidon using TLC. One ml of each of these two samples were' kept in 20 gms dried sol kept in two different containers and in another case on two different filter papers. All these samples were kept in cupboards and examined periodically upto two months time. They were extracted and identified with a varied length of time using different solvent systems. Smooth glass plates of 5x20 cms size coated with silica gel G.(E Merck Ltd. Bombay, India) in the ratio of 1 : 2 in water were used (thickness 0.25mm). The silica gel layer was allowed to settle and dried in the air for 10-15 minutes and later kept in an oven at 1100C for an hour for proper activation. Fifteen different solvents were tried on the sample for their separation and identification based on their polarity. The soil and paper samples having pesticides were extracted and analyzed with a varied period of time (from one week to eight weeks) using 800mg of soil and making small cutting of filter paper portion having insecticides/pesticides with acetone (1ml and 0.5ml) in a test tube respectively. All the samples were centrifuged before use and supernatant used for TLC. The samples were spotted with the help of five capillaries separately for each sample. These spots were allowed to dry before being placed to run in respective solvent systems.

The solvent run was restricted to the distance of 10 cms from the site of application of sample spot by marking a horizontal line on the plate with the help of a scale. All the TLC experimentation work was accomplished by ascending development. The development of colourless and separated constituents was carried but using. (i) iodine fuming (one minute exposure) and (ii) silver nitrate solution (1.0 gm AgNO₃ dissolved in 95ml of redistilled ethanol and 5ml of conc. Ammonia). After development of the colourless constituents, colour, size, shape and hRf value was noted in each case.

Results And Discussion

The results of the present investigation are given in Table -1. It is observed that amongst the used solvents, the Benzene: Acetone (9 : 1), cyclohexane : Acetone : Chloroform (70 : 25 : 5) and Hexane: Acetone (8: 2) followed by iodine fuming gave more clearer and easily interpretable results upto the period of analysis. The Dimethoate and Phosphamidon constituents did not disintegrated upto the period of analysis i.e. upto 8 weeks at the laboratory conditions. Iodine fuming gave much better results as compared to the silver nitrate spray on both the substrates for the development of spots. Benzene: Acetone (9 : 1) followed by iodine fuming as a visualizing aids has been found to be most suitable combination for Dimethoate detection while cyclohexane : Acetone: Chloroform (70 : 25 : 5) for the phosphamidon in commercial grade insecticides and pesticides in this study. The results show that these compounds can still be examined for the separation and detection upto 8 weeks period from soil and paper substrates. It persists in and on soil and filter paper upto 8 weeks times. The hRf value of dimethoate and phosphamidon are different in different solvent systems and spraying reagents. However, the hRf value may differ slightly depending on the variation of batch of the- compound, silica gel G, thickness of TLC plates, temperature etc. It needs to be examined further for various other factors such as on different types of soils, papers, prolonged time, etc. The study will provide a useful information in toxicological analysis.

● Click [here](#) to download table 1.

Acknowledgement

We are thankful to all the laboratory staff of the Forensic Science Department for helping in carrying out the present study at various stages of this work.

REFERENCES

1. Abott, D.C, Egan, H. and Thomson, J. (1964) : J chromat. 16: 481 (cf. Beynon and Elgar 1966).
2. Bami, H.L. (1972) : Hazards of Modern Insecticides, J. Ind. Acad. of Foren. Sci., 11 : 157- 170.
3. Bami., H.L. (1981) Pesticides in criminal poisoning, Journal of Forensic Science Society, 21 : 1527.
4. Beynon, K.I and Elgar, K.E. (1966) : The analysis for residues of chlorinated insecticides and acaricides, A review, Analyst 91; 143-175.
5. Bunyan, P.J. (1964) : The detection of organo-phosphorus pesticides on TLC, Analyst, 89 : 615 : 618.
6. Chakravarti, J. and Roy, B.R. (1976) : A generalized procedure for separation and identification of pesticides belonging to different groups by TLC ; J. Ind. Acad. For. Sci. 15(2) : 20.
7. Coutselinis, H. and Dinvopoulos, G. (1971) : organo phosphorus pesticides in liver and their identification J. Forensic medicine, 18 (1) : 11.

8. Dass, B. (1977) : Thin layer chromatography with reference to forensic application, J. Police Research and Development, 5 : 22-25.
9. Dewey, J.E. (1958) : qualitative and quantitative analyse of Pesticides, J. Agri and Food Chemistry, 6: 274.
10. Fargo, A. (1967) : Thin layer chromatographic detection and quantitative determination of malathion in biological materials, J. Foren. Sci., 12 : 547.
11. Fisher, E.B. and Vanwazar, J.R, (1961) : Uses of organophosphorus compounds in phosphorus, Editor: Vanwazares Vol-II - Interscience, New York.
12. Ganguly, S.K. and Battacharya, J. (1973) : Detection of small amount of pesticides in human biological material by TLC, J. of Indian Academy of Forensic Sciences, 2 : 333-338.
13. Grant, D.L. and Sherwood, C.R. and McCully, K.A. (1969) : Detection of organophosphorus compounds by using different spraying reagents, J. Chromat, 44 : 491.
14. Gunther, F. A. and Blinn, RC. (1963) : Analysis of insecticides: A treatise on sampling isolation and determination including residual methods, interscience publisher; New York.
15. Joglekar, V.D. and Mahal, H.S. (1968) : A new reagent for the identification of organophosphorus of pesticides in biological materials by TLC., J. Foren. Sciences, 142-170.
16. Kaur, Harmeet (1998) : Detection and estimation of insecticide residues in human blood and milk and their forensic significance, Ph.D thesis, submitted to Punjabi University, Patiala (Unpublished).
17. Kirchner, J. G. Ed. (1978) Thin layer chromatography, John, Willey Interscience, New York (2nd ed.)
18. Kulkarni, J.R. and Ramakrishna Reddy K. (1972): Qualitative and Quantitative analysis of pesticides, J. Ind. Acad. For. Sciences, 11: 15.
19. Ludwick Adriane, G; Lau Aldrich, N. K. and Ludwick larry, M. (1977) : Simple and sensitive technique for detecting organo chlorine pesticides on thin layer chromatogram, J. Assoc. of Anal. Chem. 60 (5) : 1077-1080.
20. Narayamswam, K; Moitra, B; Kotangle, R. S. and Bami., H. L. (1974) : Thin layer chromatography in the detection of poisoning by pesticides, J. chromatography, 95: 181- 188.
21. Prasad, B.N., Kawale, G.B. ,and Joglekar, V.D. (1982) : Detection of Phosphamidon insecticide in presence of other insecticides on thin layer chromatogram. The Indian Journal of Criminology and Criminalistics, New Delhi, 2(3-4) : 234-235.
22. Sandroni, S, And Schlitt, H. (1971) : Screening method for organo-chlorine and phosphorus pesticides residues in vegetables using TLC, J. chromatography, 55(2) : 385- 395.
23. Stahl, E. (1969) : Thin layer chromatography, Springer Berlin, N Y. P. 643.
24. Stanley, C.W. (1964) : TLC analysis on microchromato plates of some organo-phosphorus compounds, J. Chromatography, 16 : 467-475.
25. Stewart, C. P and Stolman, A (1960) : Toxicology mechanical and analytical methods, Vo1.1, Academic Press, New York.
26. Tewari S. N. (1976) : analytical Toxicology, Govt. of India Press Nasik., New Delhi.
27. Tewari, S. N. and Harpalani S.P. (1977) : Isolation and thin layer chromatographic detection and determination of phosphamidon (Dimecron) in a case of human poisoning, J. Ind. Acad. For. Science, 16 : 1-4.
28. Tewari, S. N. and Harpalani, S.P. (1973) : Detection of organophosphorus pesticide residue in autopsy tissues by thin layer chromatography, Microchim Acta, 2 : 321-24.
29. Villeneuve, D. C., Butterfield, A. G, Grant, D. L. and McCully, K.A, (1970) : Detection, separation and quantitative recovery of thirteen organo-phosphorus pesticides on silica Gel GF 254 thin layer, Choromatograms, J. Chromatography, 48; .567.
30. Zweig, O. (1964) : Analytical methods for pesticides, Plant growth regulators and Food additives, Academic Press, New York (Vol. II).

*Corresponding author and requests for clarifications and further details:

[Dr. R.K.Garg](#)

Reader and Former HOD Forensic Science,

Department Punjabi University,

Patiala 147002

India

E-mail: panky8@yahoo.com

You've been on Dr. Kuldeep Kaur's Paper for seconds.

● *N.B. It is essential to read this journal - and especially this paper as it contains several tables and high resolution graphics - under a screen resolution of 1600 x 1200 dpi or more, and preferably on a 17" or bigger monitor. If the resolution is less than this, you may see broken or overlapping tables/graphics, graphics overlying text or other anomalies. It is strongly advised to switch over to this resolution to read this journal - and especially this paper. These pages are viewed best in Netscape Navigator 4.7 and above.*

-Anil Aggrawal



[Sign My Guestbook](#)  [View My Guestbook](#)

● Click [here](#) to contact us.



You are Visitor No:



**since January 1, 2002
when this page was created.**

● This page has been constructed and maintained by Dr. Anil Aggrawal, Professor of Forensic Medicine, at the Maulana Azad Medical College, New Delhi-110002. You may want to give me the feedback to make this pages better. Please be kind enough to write your comments in the guestbook maintained above. These comments would help me make these pages better.

● IMPORTANT NOTE: ALL PAPERS APPEARING IN THIS ONLINE JOURNAL ARE COPYRIGHTED BY "ANIL AGRAWAL'S INTERNET JOURNAL OF FORENSIC MEDICINE AND TOXICOLOGY" AND MAY NOT BE REPOSTED, REPRINTED OR OTHERWISE USED IN ANY MANNER WITHOUT THE WRITTEN PERMISSION OF THE WEBMASTER

Questions or suggestions ? Please use ICQ [19727771](#) or email to dr_anil@hotmail.com

[Page Professor Anil Aggrawal via ICQ](#)



