

西妥昔单抗联合放化疗的不良反应及其防治

立即下载

Adverse reactions of cetuximab along with chemo-radiotherapy regimens and its prevention and treatments

韩佳容;邹梅娟;程刚;

摘要:

西妥昔单抗等靶向药物已在肿瘤疾病治疗中发挥越来越重要的作用,但是随着西妥昔单抗的广泛应用,关于其不良反应的报道也逐年增加。西妥昔单抗最常见的不良反应是皮肤毒性反应包括皮疹、口腔黏膜反应、甲沟炎、眼睑炎等,还有一些不良反应虽然发生率不高,但是仍应给予足够的重视如超敏反应、肺毒性、低镁血症等。了解西妥昔单抗导致不良反应的临床表现、有效的预防措施和及时的治疗方法对于患者能够顺利完成靶向治疗和提高患者在治疗期间的生活质量都是十分必要的。

关键词: 西妥昔单抗;;药物疗法;;抗肿瘤联合化疗方案;;药物不良反应

基金项目:

通讯作者: 韩佳容;

Email:

参考文献:

- [1]赵亮,方英立,周文.静脉滴注西妥昔单抗致过敏反应1例[J].中国新药与临床杂志,2011,30(3):236-237.
- [2]BLICK SK,SCOTT LJ.Cetuximab:a review of its use insquamous cell carcinoma of the head and neck and metastaticcolorectal cancer[J].Drugs,2007,67(17):2585-2607.
- [3]邵仲英,封宇飞,傅得兴.新型表皮生长因子受体抑制剂——西妥昔单抗[J].中国全科医学,2006,9(4):317-318.
- [4]MENDELSON J,BASELGA J.Status of epidermal growthfactor receptor antagonists in the biology and treatment of cancer[J].J Clin Oncol,2003,21(14):2787-2799.
- [5]LI T,PEREZ-SOLER R.Skin toxicities associated withepidermal growth factor receptor inhibitors[J].Target Oncol,2009,4(2):107-119.
- [6]FAKIH M,VINCENT M.Adverse events associated with anti-EGFR therapies for the

扩展功能

● 本文信息

[PDF\(328K\)](#)[参考文献](#)

● 服务与反馈

[引用本文](#)

● 本文关键词

相关文章

[西妥昔单抗](#)[药物疗法](#)[抗肿瘤联合](#)[化疗方案](#)[药物不良反](#)[应](#)

● 本文作者相关文章

[韩佳容](#)[邹梅娟](#)[程刚](#)

● 中国知网

- reatment of metastatic colorectal cancer[J].Curr Oncol,2010,17 Suppl 1:S18-S30.
- [7]MITCHELL EP,PEREZ-SOLER R,VAN CE,et al.Clinical presentation and pathophysiology of EGFR dermatologic toxicities[J].Oncology(Williston Park),2007,21(11 Suppl5):4-9.
- [8]MAKOTO K,MADOKA S,MEGUMU M,et al.Cetuximab-induced skin reactions are suppressed by cigarette smoking in patients with advanced colorectal cancer[J].Int J Clin Oncol,2012[Epub ahead of print].
- [9]OUWERKERK J,BDERS-DOETS C.Best practices in the management of toxicities related to anti-EGFR agents for metastatic colorectal cancer[J].Eur J Oncol Nurs,2010,14(4):337-349.
- [10]GARRETT CR,ENG C.Cetuximab in the treatment of patients with colorectal cancer[J].Expert Opin Biol Ther,2011,11(7):937-949.
- [11]GERBER PA,MELLER S,EAMES T,et al.Management of EGFR-inhibitor associated rash:a retrospective study in 49 patients[J].Eur J Med Res,2012,17:4.
- [12]SAIF MW,KIM R.Incidence and management of cutaneous toxicities associated with cetuximab[J].Expert Opin Drug Saf,2007,6(2):175-182.
- [13]王家晓,张玉梅.蒲腥解毒汤热湿敷治疗西妥昔单抗所致皮肤不良反应21例疗效观察[J].山东医药,2010,50(47):11.
- [14]龙小丽,黄艳,庞凡.金银花煎液外敷治疗西妥昔单抗致痤疮样皮疹[J].护理学杂志,2011,26(9):34.
- [15]GOBBO M,OTTAVIANI G,MUSTACCHI G,et al.Acneiform rash due to epidermal growth factor receptor inhibitors:high-level laser therapy as an innovative approach[J].Lasers Med Sci,2012,27(5):1085-1090.
- [16]PINTO C,BARONE CA,GIROLOMONI G,et al.Management of skin toxicity associated with cetuximab treatment in combination with chemotherapy or radiotherapy[J].Oncologist,2011,16(2):228-238.
- [17]HACHISUKA J,YUNOTANI S,SHIDAHARA S,et al.Effect of adapalene on cetuximab-induced painful periungual inflammation[J].J Am Acad Dermatol,2011,64(2):e20-e21.
- [18]CRIADO PR,LIMA AA.Blepharitis and trichomegaly induced by cetuximab[J].An Bras Dermatol,2010,85(6):919-920.
- [19]RAM 鯪 REZ-SORIA MP,ESPA 鯪 A-GREGORI E,AVI 鯪 鯪-MART 鯪 NEZ J,et al.Blepharitis related to cetuximab treatment in an advanced colorectal cancer patient[J].Arch Soc Esp Oftalmol,2008,83(11):665-668.
- [20]DRANKO S,KINNEY C,RAMANATHAN RK.Ocular toxicity related to cetuximab monotherapy in patients with colorectal cancer[J].Clin Colorectal Cancer,2006,6(3):224-225.
- [21]林少俊,宗井凤,廖希一,等.西妥昔单抗联合放化疗治疗进展期鼻咽癌的临床研究[J].中国癌症杂志,2009,19(2):133-137.
- [22]骆竹媚,李平,张洪,等.西妥昔单抗联合放化疗治疗晚期转移性恶性肿瘤的临床观察[J].肿瘤预防与治疗,2010,23(3):203-205,237.
- [23]冯惠霞,蒋红花,李桂容,等.晚期鼻咽癌局部同期放化疗联合西妥昔单抗治疗的护理观察[J].护士进修杂志,2012,27(10):898-899.
- [24]王芳,王汉渝,胡伟汉,等.调强放疗联合西妥昔单抗及顺铂治疗晚期鼻咽癌[J].中国

肿瘤生物治疗杂志,2008,15(5):489-493.

[25]MAS-VIDAL A,COTO-SEGURA P,GALACHE-OSUMA C,etal.Psoriasis induced by cetuximab:A paradoxical adverse effect[J].Australas J Dermatol,2011,52(1):56-68.

[26]翁羽骅.银屑病的护理[J].中国医药指南,2012,10(24):260-261.

[27]严颖,龚继芳,张小田.消化道肿瘤化疗及靶向治疗药物过敏反应的临床对策(附25例分析)[J].临床药物治疗杂志,2009,7(5):57-60.

[28]LENZ HJ.Management and preparedness for infusion andhypersensitivity reactions [J].Oncologist,2007,12(5):601-609.

[29]郭桂芳,夏良平,张蓓,等.西妥昔单抗联合化疗对53例晚期结直肠癌的疗效[J].癌症,2009,28(12):1317-1323.

[30]RESCH G,SCHABERL-MOSER R,KIER P,et al.Infusionreactions to the chimeric EGFR inhibitor cetuximab—change tothe fully human anti-EGFR monoclonal antibody panitumumab issafe[J].Ann Oncolo,2011,22(2):486-487.

[31]PETRELLI F,BORGONOVO K,CABIDDU M,et al.Risk ofanti-EGFR monoclonal