

Pathwise uniqueness for reflecting Brownian motion in certain planar Lipschitz domains

Richard F. Bass, *University of Connecticut*
Krzysztof Burdzy, *University of Washington*

Abstract

We give a simple proof that in a Lipschitz domain in two dimensions with Lipschitz constant one, there is pathwise uniqueness for the Skorokhod equation governing reflecting Brownian motion.

Full text: [PDF](#) | [PostScript](#)

Pages: 178-181

Published on: August 30, 2006

Bibliography

1. M. Barlow, K. Burdzy, H. Kaspi and A. Mandelbaum. Variably skewed Brownian motion *Electr. Comm. Probab.* 5 (2000), paper 6, pp. 57-66. [Math. Review 2001j:60146](#)
2. R. Bass, K. Burdzy and Z. Chen. Uniqueness for reflecting Brownian motion in lip domains *Ann. I. H. Poincar'e* 41 (2005) 197-235. [Math. Review 2005k:60174](#)
3. R. Bass and E.P. Hsu, Pathwise uniqueness for reflecting Brownian motion in Euclidean domains. *Probab. Th. Related Fields* 117 (2000) 183-200. [Math. Review 2001e:60164](#)
4. I. Karatzas and S.E. Shreve. *Brownian Motion and Stochastic Calculus*, 2nd Edition, Springer Verlag, New York, 1991. [Math. Review 92h:60127](#)
5. D. Revuz and M. Yor, *Continuous Martingales and Brownian Motion*, 3rd ed. Springer, Berlin, 1999. [Math. Review 2000h:60050](#)

Research Support Tool

[Capture Cite](#)
[View Metadata](#)
[Printer Friendly](#)

▼ [Context](#)

[Author Address](#)

▼ [Action](#)

[Email Author](#)
[Email Others](#)