



Mathematics > Statistics Theory

# Renorming divergent perpetuities

Paweł Hitczenko, Jacek Wesółowski

(Submitted on 14 Jul 2011)

We consider a sequence of random variables  $(R_n)$  defined by the recurrence  $R_n = Q_n + M_n R_{n-1}$ ,  $n \geq 1$ , where  $R_0$  is arbitrary and  $(Q_n, M_n)$ ,  $n \geq 1$ , are i.i.d. copies of a two-dimensional random vector  $(Q, M)$ , and  $(Q_n, M_n)$  is independent of  $R_{n-1}$ . It is well known that if  $E\{\ln|M|\} < 0$  and  $E\{\ln^+|Q|\} < \infty$ , then the sequence  $(R_n)$  converges in distribution to a random variable  $R$  given by  $R \stackrel{d}{=} \sum_{k=1}^{\infty} Q_k \prod_{j=1}^{k-1} M_j$ , and usually referred to as perpetuity. In this paper we consider a situation in which the sequence  $(R_n)$  itself does not converge. We assume that  $E\{\ln|M|\}$  exists but that it is non-negative and we ask if in this situation the sequence  $(R_n)$ , after suitable normalization, converges in distribution to a non-degenerate limit.

Comments: Published in at [this http URL](#) the Bernoulli ([this http URL](#)) by the International Statistical Institute/Bernoulli Society ([this http URL](#))

Subjects: **Statistics Theory (math.ST)**

Journal reference: Bernoulli 2011, Vol. 17, No. 3, 880-894

DOI: [10.3150/10-BEJ297](#)

Report number: IMS-BEJ-BEJ297

Cite as: [arXiv:1107.2753v1](#) [math.ST]

## Submission history

From: Paweł Hitczenko [[view email](#)]  
[v1] Thu, 14 Jul 2011 08:45:02 GMT (36kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

## Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

**math.ST**

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

Change to browse by:

[math](#)  
[stat](#)

## References & Citations

- [NASA ADS](#)

## Bookmark (what is this?)

