

Random walks with k-wise independent increments

Itai Benjamini, *The Weizmann Institute of Science*
Gady Kozma, *The Weizmann Institute of Science*
Dan Romik, *University of California, Berkeley*

Abstract

We construct examples of a random walk with pairwise-independent steps which is almost surely bounded, and for any m and k a random walk with k -wise independent steps which has no stationary distribution modulo m .

Full text: [PDF](#) | [PostScript](#)

Pages: 100-107

Published on: July 6, 2006

Bibliography

1. Alon, Noga; Spencer, Joel H. The probabilistic method. Optimization. *Wiley-Interscience [John Wiley & Sons], New York*, 2000. xviii+301 pp. ISBN: 0-471-37046-0 [MR1885388](#) (2003f:60003)
2. Janson, Svante. Some pairwise independent sequences for which the central limit theorem *Stochastics* 23 (1988), no. 4, 439--448. [MR0943814](#) (89e:60048)
3. Joffe, A. On a set of almost deterministic k -independent random variables. *Ann. Probability* 2 (1974), no. 1, 161--162. [MR0356150](#) (50 #8621)
4. Schipp, F.; Wade, W. R.; Simon, P.. Walsh series. *Adam Hilger, Ltd., Bristol*, 1990. x+560 pp. ISBN: 0-7503-0068-X [MR1117682](#) (92g:42001)
5. Wade, William R. Dyadic harmonic analysis. 313--350, *Contemp. Math.*, 208, *Amer. Math. Soc., Providence, RI*, 1997. [MR1467014](#) (98m:42046)

Research Support Tool

[Capture Cite](#)
[View Metadata](#)
[Printer Friendly](#)

▼ [Context](#)

[Author Address](#)

▼ [Action](#)

[Email Author](#)
[Email Others](#)