

Cytotoxic T lymphocyte responses to hepatitis B virus small surface antigen in BALB/c mice

SHI Li, LIU Shu-ren, YI Xue-rui, KONG Qiang-qing, ZHANG Yi-jun

Center for Hepatic Diseases of PLA, 458 Hospital of PLA, Guangzhou 510602, China

Abstract: Objective To observe cytotoxic T lymphocytes (CTL) responses induced by immunization of mice with hepatitis B small surface antigen (S-HBsAg). Methods BALB/c(H-2^d) mice were injected intraperitoneally with S-HBsAg at the doses ranging from 0-5 μ g with a booster injection 2 weeks later. Four weeks after the booster immunization, the specific cytotoxic reactivity of the spleen lymphocytes isolated from the immunized mice, following stimulation by S-HBsAg specific CTL epitope peptide in vitro, with ⁵¹Cr-labeled P815 cells treated with the peptide was observed. Results The percentage of specific release of ⁵¹Cr in mice immunized with S-HBsAg at the doses of 0, 0.65, 1.25, 2.5, and 5 μ g was 31.21%, 42.36%, 42.36%, 19.32%, 91.21%, 2.97%, 69.25%, 4.13% and 51.49%, 1.661% respectively. In mice receiving the only primary vaccination, the corresponding percentage was 27.34%, 4.25%, 32.27%, 5.35%, 56.28%, 4.35%, 44.34%, 8.65% and 40.76%, 56% respectively. Conclusion Immunization with S-HBsAg efficiently elicits CTL response in BALB/c mice in a dose-dependent manner and enhanced by booster immunization.

Key words: hepatitis B virus; small surface antigen; BALB/c mice; immunization; cytotoxic T lymphocytes

Hepatitis B virus (HBV) is the major etiological agent responsible for chronic liver disease, and chronic HBV infection often leads to hepatic cirrhosis and hepatocellular carcinoma. At present, the number of patients with chronic HBV infection has reached approximately 350 million worldwide, and interferon and lamivudine prevail in the treatment. In spite of the high cost of interferon therapy, adverse effect is a frequent clinical entity while only roughly 35% of the patients may have HBe/anti-HBe seroconversion with negative HBV DNA. Lamivudine possesses stronger antiviral effect, but the high rate of relapse after withdrawal of a 12-month treatment course remains a problem, with as low as about 20% of the patients having long-term HBe/anti-HBe seroconversion with negative HBV DNA and genotypic resistance. Recent clinical and experimental studies have demonstrated the efficacy of immunization with HBV vaccine and specific anti-HBV cytotoxic T lymphocyte (CTL) transfer in elimination of HBV and fighting pre-existing infections^[1-5], and in this study, we aim to test whether small surface antigen of HBV (S-HBsAg) may elicit CTL response in BALB/c mice and determine the dose of this vaccine that induce the strongest CTL response.

MATERIALS AND METHODS

Materials

The recombinant hepatitis B (HB) vaccine, Recombivax HB, was a yeast-derived HBV small surface antigen produced by Merck & Co, containing L^d-restricted CTL epitope S₂₈₋₃₉ (IPQSLSWWTSL). The specific peptide IPQSLDSWWTSL (S₂₈₋₃₉), which represents

a L^d-restricted CTL epitope of the S-HBsAg of H-2^d BALB/c mice, was synthesized by Saibaisheng Co. and dissolved at the concentrations of 10-50 mg/ml in the stock solution containing 70% acetonitrile and 0.1% TFA or DMSO, and diluted with culture medium before use.

H-2^d mastocytoma cell line P815 were purchased from Chinese Cell Culture Center.

Methods

Immunization of mice Female BALB/c (H-2^d) mice of 6-10 weeks old were bred under standard pathogen-free conditions. The mice were grouped to receive intraperitoneal injections with Recombivax HB at different doses, with a booster injection 2 weeks later. Details of grouping and the doses injected are listed below:

Group 0 (n=10), in which the mice were injected with 5 μ g adjuvant (aluminum hydroxide); Group 0.65 (n=10), in which the mice received injection of 0.65 μ g Recombivax HB; Group 1.25 (n=10), in which Recombivax HB was injected at the dose of 1.25 μ g; Group 2.5 (n=10), in which Recombivax HB at the dose of 2.5 μ g was used; Group 5 (n=10), in which the dose of Recombivax HB was 5 μ g.

CTL assays

P815 cells (1 \times 10⁷) were treated with specific peptide and labeled with 100 μ g Na₂⁵¹CrO₄ for 1 h at 37 $^{\circ}$ C, to serve as the target cells. The spleen cells as the effector cells were suspended in complete DMEM supplemented with 10% fetal calf serum (FCS), followed by treatment with 2-mercaptoethanol (final concentration of 5 \times 10⁻⁵ mol/L), murine interleukin (IL) -2 (final concentration of 5 U/ml). After 5 days of in vitro culture

and washing twice, the cytotoxic effector lymphocytes were harvested and cocultured in triplicates with the target P815 cells at the ratio of 50 颐 with 1伊0⁴ cells in each well. After a 4-hour incubation at 37 益 and centrifugation, the supernant (100 滋) was collected for assessment of the radioactivity with 酌counter.

Two weeks after the primary inoculation, the spleen lymphocytes isolated from half of the mice were incubated with ⁵¹Cr-labeled P815 cells, and the percentage of specific ⁵¹Cr release (PSR) of the spleen lymphocytes for only the primary immunization was calculated; four weeks after the booster immunization, the spleen lymphocytes isolated from the rest of the mice were incubated in the presence or absence of ⁵¹Cr-labelled P815 cells, and PSR of the spleen lymphocytes was calculated.

PSR was calculated by (experimental release - spontaneous release)/ (maximum release - spontaneous release)伊100%, where the experimental release was defined as the mean cpm released by the target cells in the presence of effector cells. Maximum release was the mean cpm measured after lysis of the target cells with 5% Triton X-100. Spontaneous release was the mean cpm present in the culture medium of the target cells alone. The purpose of calculating PSR was to eliminate the possible interference of the background factors.

Statistical analysis

A non-parametric Mann-Whitney U test was used to compare the results between different groups. A P value of less than 0.05 or 0.01 was considered statistically significant or very statistically significant, respectively.

RESULTS

In the CTL assay, the ratio of spontaneous release to maximum release was 28.76% 依3.45% for the target cells.

In BALB/c mice with booster immunization, the PSR of the spleen lymphocytes of the mice in Groups 0, 0.65, 1.25, 2.5 and 5 was 31.21% 依2.23%, 42.36% 依19.32%, 91.21% 依2.97%, 69.25% 依4.13%, and 51.49% 依1.661% respectively, which was the highest in Group 1.25 and lowest in Group 0. PSR was the lowest in Group 0.65 among the mice with booster immunization.

In mice with only primary immunization, PSR of the spleen lymphocytes in Groups 0, 0.65, 1.25, 2.5, and 5 was 27.34% 依4.25%, 32.27% 依5.35%, 56.28% 依24.35%, 44.34% 依8.65% and 40.76% 依6% respectively.

For the target cells without treatment with the specific peptide, the PSR of the spleen lymphocytes from

the mice with booster immunization were 20.56% 依17.28%, 22.37% 依8.29%, 34.29% 依1.23%, 21.27% 依20.30%, 32.38% 依6.56%, respectively.

DISCUSSION

HBV is a noncytopathic DNA virus, and adult onset of HBV infection usually results in self-limited acute hepatitis followed by viral clearance, with only 5% of the infected adults developing chronic infection. In patients with self-limited acute HBV infection, the virus can be spontaneously cleared from the serum, accompanied by increased cellular immune responses, especially CTL response. In the majority of the patients with chronic hepatitis, HBV-specific CTL response was thought to be low so as to give rise to HBV persistence. The therapy of HBV infection depends mainly on host cellular immune response. CTL play important roles in the elimination of the virus-infected cells, and in HBV-infected patients, CTL is involved in the reduction of viral load. Although immunization with S-HBsAg vaccine has been acknowledged to induce humoral immune response and protect against HBV attack, the questions whether this vaccine also elicits cellular immune response, and if so, what the optimal dose is to induce the strongest CTL response, have not been answered 啍4 暂.

To investigate cell-mediated immune responses to S-HBsAg, in vivo primed spleen cells from immunized mice were stimulated in vitro with a specific peptide and tested for CTL activity against P815 cells treated with the specific peptide. We found that the PSR in the vaccinated mice at the doses of 0.65, 1.25, 2.5, and 5 滋 was all significantly higher than that in the control group without vaccination, and also higher than those in the parallel groups with the target cells not treated with the specific peptide, indicating specific CTL response to S-HBsAg vaccination. In the experimental groups, PSR tended to increase as the dose of vaccination increased from 0.65 to 1.25 滋, but not with the doses from 2.5 to 5 滋. PSR in Group 1.25 was significantly higher than those in other groups (P<0.05). The difference in PSR between Groups 0.65 and 5 was not significant (P>0.05), suggesting that the CTL response is dependent on the dose of immunization. The PSR in mice with booster immunization was significantly higher than that in the mice with only primary inoculation, indicating that the booster immunization enhanced the CTL response.

Specific CTLs are the major cells participating in the cellular immune response against acute, chronic, and persistent infections with intracellular pathogens

such as viruses, and function as a crucial defense against the spread of these pathogens through both cytolysis of infected cells they recognized and noncytolytic mechanisms such as cytokine production (interferon and tumor necrosis factor α). CTLs specific for the epitopes in HBV contribute to the control of HBV infection, which were readily detectable in patients with HBV clearance (self-limited acute hepatitis recovering from acute HBV infection exhibits a polyclonal and multispecific CTL response to HBsAg) but not in cases of chronic infection. It is now clear that therapeutic vaccination with immunogenic peptides depends heavily on CTL induction. Vaccination strategies favoring the induction of stable cell-mediated responses might protect against subsequent viral infections. Patients with HBV infection benefit from injection of S-HBsAg through induction of viral clearance by enhancement of suboptimal cellular immune responses. Injection of this vaccine induces marked cellular immune responses to HBsAg, significantly decreasing HBV viral load and increasing the rate of HBe/anti-HBe seroconversion. The immunotherapeutic effect of the vaccination in chronically HBV-infected patients to enhance HBV-specific cellular immune responses may be of value in the control of persistent infection.

CTLs recognize the target cells through the CTL epitope and the major histocompatibility complex- II determinants on the surface of the target cells. In this study, the S-HBsAg vaccine contained HBV Ld-restricted CTL epitope S₂₈₋₃₉ (IPQSLOSWSL). P815 cells treated with the specific peptide expressed H-2^d MHC-II restricted determinants similar with the BALB/c. The in vivo primed spleen cells from the immunized BALB/c (H-2^d) mice stimulated in vitro with specific peptide ex-

hibited specific cytotoxic reactivity against P815 cells, and high-dose vaccination resulted in substantial specific CTL activity of the spleen cells, which was markedly enhanced by a booster vaccination^[11].

REFERENCES

- 1 Rizzetto M, Zanetti AR. Progress in the prevention and control of viral hepatitis type B: closing remarks. *J Med Virol*, 2002, 67(3): 463-6.
- 2 Wong DKH, Cheung AM, O'Rourke K, et al. Effect of alpha-interferon in patients with hepatitis Be antigen-positive chronic hepatitis B: a meta-analysis. *Ann Intern Med*, 1993, 119: 312-23.
- 3 Ohkoshi S, Ogata N, Ichida T. The long-term clinical outcome of 1-year treatment of chronic hepatitis B with lamivudine-5 years observation. *Hepatol Res*, 2003, 27(1):13-7.
- 4 Lagget M, Rizzetto M. Current pharmacotherapy for the treatment of chronic hepatitis B. *Expert Opin Pharmacother*, 2003, 4(10): 1821-7.
- 5 Stanislas Pol, Bertrand Nalpas, Francoise Driss, et al. Efficacy and limitations of a specific immunotherapy in chronic hepatitis B. *Hepatol*, 2001, 34: 917-921.
- 6 Du DW, Jia ZS, Li GY, et al. HBV DNA vaccine with adjuvant cytokines induced specific immune responses against HBV infection. *World J Gastroenterol*, 2003, 9(1): 108-11.
- 7 Michel ML. Towards immunotherapy for chronic hepatitis B virus infections. *Vaccine*, 2002, 19, 20 (Suppl 4): A83-8.
- 8 Chakravarty R, Neogi M, Roychowdhury S, et al. Presence of hepatitis B surface antigen mutant G145R DNA in the peripheral blood leukocytes of the family members of an asymptomatic carrier and evidence of its horizontal transmission. *Virus Res*, 2002, 90 (1-2): 133-41.
- 9 Francois G, Hallauer J, Van Damme P. Hepatitis B vaccination: how to reach risk groups. *Vaccine*, 2002, 21(1-2): 1-4.
- 10 Williams A. Reduction in the hepatitis B related burden of disease --measuring the success of universal immunisation programs. *Commun Dis Intell*, 2002, 26(3): 458-60.
- 11 Kasahara S, Ando K, Saito K, et al. Lack of tumor necrosis factor alpha induces impaired proliferation of hepatitis B virus-specific cytotoxic T lymphocytes. *J Virol*, 2003, 77(4): 2469-76.

BALB/c小鼠腹腔接种 HBV 表面 S 抗原疫苗引起的 CTL 反应

施 理 袁 树 人 袁 学 瑞 袁 祥 平 袁 张 宜 俊 袁 58 医院全军肝病中心 袁 东 广州 510602

摘要 目的 研究 BALB/c 小鼠免疫接种 HBV 表面 S 抗原疫苗能否引起特异性细胞毒 T 淋巴细胞 CTL 反应。方法 分别给 BALB/c 小鼠腹腔内接种 0~6 周 S-HBsAg 疫苗,两周后加强一次,两周后分离小鼠脾细胞,体外用 S-HBsAg 特异性 CTL 表位多肽刺激,用特异性多肽⁵¹Cr 标记的 P815 细胞作为靶细胞,用⁵¹Cr 释放实验检测 CTL 反应。结果 分别接种 0 周、2.5 周、5 周 S-HBsAg 的小鼠,其脾淋巴细胞 CTL 特异性释放率分别为 31.21%、23.23%、2.36%、9.32%、1.21%、2.97%、9.25%、4.13%、1.49%、1.66%。只接受一次免疫接种的小鼠,其脾淋巴细胞 CTL 特异性释放率分别为 27.34%、4.25%、2.27%、5.35%、6.28%、4.35%、4.34%、8.65%、0.76%。结论 BALB/c(H-2^d)小鼠腹腔内接种 S-HBsAg 引起剂量依赖性特异性 CTL 反应,加强免疫能够提高 CTL 反应。关键词 HBV 表面 S 抗原疫苗, BALB/c 小鼠,免疫接种,细胞毒 T 淋巴细胞

中图分类号 R392.12; R512.62 文献标识码 A 文章编号 000-2588(2003)12-1242-03