

中立型时滞系统在 C_1 空间中的稳定性

章毅 钟守铭
(电子科技大学)

摘 要

本文中研究了中立型时滞系统在 C_1 空间中的稳定性, 获得了半线性和非线性中立型时滞系统在 C_1 空间中稳定性的一些充分条件.

关键词——中立型; 时滞; 空间; 稳定性.

一、引 言

在时滞系统的稳定性理论中, 对于中立型时滞系统的稳定性的研究进展很慢^{[1]-[9]}, 这主要是由于问题本身的困难性及缺少方法所致. 文[2]和[3]研究了中立型系统在通常意义下的稳定性. 文[4]-[7]则在 C_1 空间中对中立型时滞系统的稳定性进行了研究, 获得了稳定、渐近稳定性方面的定理. 众所周知, 这在通常意义下研究稳定性更加困难. 在文[8]、[9]中, 笔者提出了中立型时滞系统的零解在 C_1 空间中的大范围指数稳定性问题. 本文将利用另一种方法来继续研究中立型时滞系统在 C_1 空间中的大范围指数稳定性, 进一步改进了过去的研究结果.

我们总是假定本文所考虑的系统解是存在、唯一和连续的.

二、半线性系统

研究如下的中立型系统:

$$\left. \begin{aligned} \dot{x}(t) &= A(t)x(t) + f[t, x(t - \Delta(t)), \dot{x}(t - \Delta(t))], \\ x(t) &= \varphi(t), \quad \dot{x}(t) = \dot{\varphi}(t), \quad t_0 - \Delta \leq t \leq t_0, \end{aligned} \right\} \quad (1)$$

其中 $x = \text{col}(x_1, x_2, \dots, x_n)$; $A(t)$ 为 $n \times n$ 阶连续函数矩阵, 且有 $\sup_{t \geq t_0} \|A(t)\| \leq a < +\infty$. $f: R \times R^n \times R^n \rightarrow R^n$, $f(t, 0, 0) \equiv 0$; 时滞 $\Delta(t)$ 是非负连续有界函数, 即 $0 \leq \Delta(t) \leq \Delta$, Δ 是某一常数. $\varphi(t)$, $\dot{\varphi}(t)$ 在 $t_0 - \Delta \leq t \leq t_0$ 上连续, 记

$$\|R\| = \sup_{t_0 - \Delta \leq t \leq t_0} [\|\varphi(t)\| + \|\dot{\varphi}(t)\|].$$

定义. 如果存在常数 $\lambda > 0$, 且对任意的 $\alpha > 0$, 存在 $K(\alpha) > 0$, 使得当 $\|R\| \leq \alpha$ 时, 有:

$$\|x(t)\| + \|\dot{x}(t)\| \leq K(\alpha)\|R\|e^{-\lambda(t-t_0)}, \quad t \geq t_0,$$

则称系统(1)的零解在 C_1 空间中大范围指数稳定.

设 $y(s, t)$ 满足

$$\begin{cases} \frac{\partial y(s, t)}{\partial t} = A(t)y(s, t), \\ Y(s, s) = I, \quad (I \text{ 是单位矩阵}). \end{cases}$$

定理 1. 若系统(1)满足以下条件:

(i) $\|Y(s, t)\| \leq e^{-r(s-t)}, (t, s \geq t_0), r > 0;$

(ii) $\|f[t, x(t - \Delta(t)), \dot{x}(t - \Delta(t))]\| \leq b\|x(t - \Delta(t))\| + c\|\dot{x}(t - \Delta(t))\|,$

其中 $b \geq 0, c > 0;$

(iii) $\frac{b + ac}{r} + c < 1;$

则系统(1)的零解在 C_1 空间中大范围指数稳定.

证. 构造函数

$$f(\lambda) = \frac{e^{\lambda\Delta}(b + ac)}{r - \lambda} + e^{\lambda\Delta}c,$$

显然, $f(\lambda)$ 在原点的某一足够小的区域内是连续的, 因为:

$$f(0) = \frac{b + ac}{r} + c < 1,$$

故必有足够小的 $\lambda_1 (\lambda_1 < r)$, 使得:

$$f(\lambda_1) = \frac{e^{\lambda_1\Delta}(b + ac)}{r - \lambda_1} + e^{\lambda_1\Delta}c \stackrel{\text{def}}{=} \delta < 1.$$

对系统(1)利用参数变易法, 可得:

$$x(t) = y(t_0, t)x(t_0) + \int_{t_0}^t y(s, t)f[s, x(s - \Delta(s)), \dot{x}(s - \Delta(s))]ds,$$

代入(1)式得到:

$$\begin{aligned} \dot{x}(t) = & A(t)Y(t_0, t)x(t_0) + \int_{t_0}^t A(t)Y(s, t)f[s, x(s - \Delta(s)), \dot{x}(s - \Delta(s))]ds \\ & + f[t, x(t - \Delta(t)), \dot{x}(t - \Delta(t))]. \end{aligned}$$

利用条件 (i)、(ii) 可推得:

$$\|x(t)\| \leq e^{-r(t-t_0)}\|R\| + \int_{t_0}^t e^{-r(t-s)}[b\|x(s - \Delta(s))\| + c\|\dot{x}(s - \Delta(s))\|]ds$$

$$\begin{aligned} \|\dot{x}(t)\| \leq & e^{-r(t-t_0)}a\|R\| + \int_{t_0}^t ae^{-r(t-s)}[b\|x(s - \Delta(s))\| + c\|\dot{x}(s - \Delta(s))\|]ds \\ & + [b\|x(t - \Delta(t))\| + c\|\dot{x}(t - \Delta(t))\|]. \end{aligned}$$

(a) 若 $b \neq 0$, 令

$$P(t) = [b\|x(t)\| + c\|\dot{x}(t)\|]e^{\lambda_1(t-t_0)}, \quad t \geq t_0 - \Delta,$$

不难推得:

$$P(t) \leq (b+c+ac)\|R\| + e^{\lambda_1 \Delta}(b+ac) \int_{t_0}^t e^{-r(t-s)} P(s - \Delta(s)) ds + ce^{\lambda_1 \Delta} P(t - \Delta(t)), \quad t \geq t_0. \quad (2)$$

现将证明, 对一切 $t \geq t_0$, 有:

$$P(t) \leq \frac{2(b+c+ac)}{1-\delta} \|R\| \stackrel{\text{def}}{=} M.$$

注意到当 $t_0 - \Delta \leq t \leq t_0$ 时, 有:

$$P(t) \leq (b+c)\|R\| < (b+c+ac)\|R\| < M, \quad (3)$$

于是, 假如上面的结论不成立, 则必有 $t_1 > t_0$, 使得:

$$P(t_1) = M, \quad P(t) < M, \quad t_0 - \Delta \leq t < t_1. \quad (4)$$

将 $P(t) \leq M$, $t_0 - \Delta \leq t \leq t_1$ 代入(2)式右端, 有:

$$P(t) \leq (b+c+ac)\|R\| + \delta M, \quad t_0 \leq t \leq t_1,$$

结合(3)式, 则有:

$$P(t) \leq (b+c+ac)\|R\| + \delta M, \quad t_0 - \Delta \leq t \leq t_1.$$

下面用数学归纳法证明下式成立:

$$P(t) \leq (b+c+ac)\|R\| [1 + \delta + \delta^2 + \dots + \delta^{n-1}] + \delta^n M, \quad t_0 - \Delta \leq t \leq t_1. \quad (5)$$

当 $n=1$ 时, 显然(5)式成立. 设 $n=k$ 时, (5)式成立, 即有:

$$P(t) \leq (b+c+ac)\|R\| [1 + \delta + \delta^2 + \dots + \delta^{k-1}] + \delta^k M \stackrel{\text{def}}{=} M_k, \quad t_0 - \Delta \leq t \leq t_1.$$

将 $P(t) \leq M_k$, $t_0 - \Delta \leq t \leq t_1$ 代入(2)式右端, 则有:

$$P(t) \leq (b+c+ac)\|R\| + \delta M_k, \quad t_0 \leq t \leq t_1.$$

结合(3)式, 则有:

$$\begin{aligned} P(t) &\leq (b+c+ac)\|R\| + \delta M_k \\ &= (b+c+ac)\|R\| (1 + \delta + \delta^2 + \dots + \delta^k) + \delta^{k+1} M, \\ &\quad t_0 - \Delta \leq t \leq t_1. \end{aligned}$$

即当 $n=k+1$ 时, (5)式亦成立, 从而证明了对一切自然数 n , (5)式成立. 特别地有:

$$P(t_1) \leq (b+c+ac)\|R\| [1 + \delta + \delta^2 + \dots + \delta^{n-1}] + \delta^n M.$$

令 $n \rightarrow \infty$ 时, 注意到 $\delta < 1$, 于是得到:

$$P(t_1) \leq \frac{(b+c+ac)}{1-\delta} \|R\| < M,$$

这与(4)式中 $P(t_1) = M$ 相矛盾. 这样便证明了不等式

$$P(t) < \frac{2(b+c+ac)}{1-\delta} \|R\|, \quad t \geq t_0,$$

成立, 由此式可推得:

$$\|x(t)\| + \|\dot{x}(t)\| \leq \frac{2(b+c+ac)}{(1-\delta)\min(b,c)} \|R\| e^{-\lambda_1(t-t_0)}, \quad t \geq t_0.$$

此式表明系统(1)的零解在 C_1 空间中大范围指数稳定.

(b) 若 $b = 0$, 则条件 (iii) 成为 $ac/r + c < 1$. 此时, 必有足够小的 $\varepsilon > 0$, 使得 $(\varepsilon + ac)/r + c < 1$.

令 $P(t) = (\varepsilon\|x(t)\| + c\|\dot{x}(t)\|)e^{\lambda_1(t-t_0)}$, $t \geq t_0 - \Delta$, 可推得:

$$P(t) \leq (\varepsilon + c + ac)\|R\| + e^{\lambda_1\Delta}(\varepsilon + ac) \int_{t_0}^t e^{-r(s-s)}P(s - \Delta(s))ds \\ + ce^{\lambda_1\Delta}P(t - \Delta(t)), \quad t \geq t_0.$$

然后类似于前面的推导, 可得到:

$$\|x(t)\| + \|\dot{x}(t)\| \leq \frac{2(\varepsilon + c + ac)}{(1 - \delta)\min(\varepsilon, c)} \|R\| e^{-\lambda_1(t-t_0)}, \quad t \geq t_0.$$

所以系统(1)的零解在 C_1 空间中大范围指数稳定. 证毕.

考虑系统:

$$\left. \begin{aligned} \dot{x}(t) &= A(t)x(t) + D(t)x(t - \Delta(t)) + f[t, x(t - \Delta(t)), \dot{x}(t - \Delta(t))], \\ x(t) &= \varphi(t), \quad \dot{x}(t) = \dot{\varphi}(t), \quad t_0 - \Delta \leq t \leq t_0. \end{aligned} \right\} \quad (6)$$

其中 $D(t)$ 是 $n \times n$ 阶连续函数矩阵, 且有 $\sup_{t \geq t_0} \|D(t)\| \leq d < +\infty$. 其它条件与系统(1)一样.

设 $y_1(s, t)$ 满足

$$\begin{cases} \frac{\partial y_1(s, t)}{\partial t} = [A(t) + D(t)]y_1(s, t), \\ y_1(s, s) = I, \quad (I \text{ 是单位矩阵}) \end{cases}$$

定理 2. 若系统(6)满足以下条件:

(i) $\|y_1(s, t)\| \leq e^{-r(t-s)}$, $(t, s \geq t_0)$, $r > 0$;

(ii) $\|f[t, x(t - \Delta(t)), \dot{x}(t - \Delta(t))]\| \leq b \sup_{t-\Delta \leq \theta \leq t} \|x(\theta)\| + c \sup_{t-\Delta \leq \theta \leq t} \|\dot{x}(\theta)\|$,

其中 $b \geq 0$, $c > 0$;

(iii) $\frac{b + ac}{r} + \Delta d + c < 1$;

则系统(6)的零解在 C_1 空间中大范围指数稳定.

证. 只要注意到:

$$\|x(t - \Delta(t)) - x(t)\| \leq \Delta \sup_{t-\Delta \leq \theta \leq t} \|\dot{x}(\theta)\|,$$

类似定理 1 的证明, 可推证本定理.

推论. 考虑系统:

$$\begin{cases} \dot{x}(t) = ax(t) + bx(t - \Delta(t)) + c\dot{x}(t - \Delta(t)), \\ x(t) = \varphi(t), \quad \dot{x}(t) = \dot{\varphi}(t), \quad t_0 - \Delta \leq t \leq t_0. \end{cases}$$

若下面任一条件满足:

(i) $a < 0$, $\left| \frac{b}{a} \right| + 2|c| < 1$,

(ii) $a < 0$, $\Delta|b| + 2|c| < 1$,

则其零解在 C_1 空间中大范围指数稳定.

三、非线性系统

考虑系统:

$$\left. \begin{aligned} \dot{x}(t) &= F[t, x(t), x(t - \Delta(t)), \dot{x}(t - \Delta(t))], \\ x(t) &= \varphi(t), \quad \dot{x}(t) = \dot{\varphi}(t), \quad t_0 - \Delta \leq t \leq t_0, \end{aligned} \right\} \quad (7)$$

其中 $F: R \times R^n \times R^n \times R^n \rightarrow R^n$, $F(t, 0, 0, 0) \equiv 0$. 时滞 $\Delta(t)$ 是一个非负有界连续函数, 即 $0 \leq \Delta(t) \leq \Delta$, Δ 是某一常数. $\varphi(t)$ 、 $\dot{\varphi}(t)$ 在 $t_0 - \Delta \leq t \leq t_0$ 上连续, 记

$$\|R\| = \sup_{t_0 - \Delta \leq t \leq t_0} [\|\varphi(t)\| + \|\dot{\varphi}(t)\|].$$

$$\begin{aligned} \text{条件 I: } \|F(t, x_1, y_1, z_1) - F(t, x_2, y_2, z_2)\| \\ \leq a\|x_1 - x_2\| + b\|y_1 - y_2\| + c\|z_1 - z_2\|, \end{aligned}$$

其中 $a > 0$, $b \geq 0$, $c > 0$.

条件 II: 存在 $V(t, x)$ 满足下列不等式:

$$(i) \quad (\alpha\|x\|)^2 \leq V(t, x) \leq (\beta\|x\|)^2, \quad \alpha > 0, \beta > 0;$$

$$(ii) \quad \left| \frac{\partial V(t, x)}{\partial x} \right| \leq 2\delta \sqrt{V(t, x)}, \quad \delta > 0;$$

$$(iii) \quad \frac{\partial V(t, x)}{\partial x} F[t, x(t), 0, 0] + \frac{\partial V(t, x)}{\partial t} \leq -2rV, \quad r > 0.$$

$$\text{条件 III: } \frac{\delta(b + ac)}{\alpha r} + c < 1.$$

定理 3. 若条件 I、II、III 成立, 则系统 (7) 的零解是在 C_1 空间中大范围指数稳定.

证. 设 $x(t)$ 是 (7) 的解, 记 $V(t) = V(t, x(t))$, 则由条件 I、II 可得:

$$\begin{aligned} \dot{V}(t) &= \frac{\partial V(t, x)}{\partial x} F[t, x(t), 0, 0] + \frac{\partial V(t, x)}{\partial t} \\ &\quad + \frac{\partial V(t, x)}{\partial x} \{F[t, x(t), x(t - \Delta(t)), \dot{x}(t - \Delta(t))] - F[t, x(t), 0, 0]\} \\ &\leq -2rV(t) + 2\delta\sqrt{V(t)}(b\|x(t - \Delta(t))\| + c\|\dot{x}(t - \Delta(t))\|). \end{aligned}$$

令: $V_1(t) = \sqrt{V(t) + \|R\|^2 e^{-2r(t-t_0)}}$, $t \geq t_0$, 则有:

$$V_1(t) \geq \sqrt{V(t)} \geq \alpha\|x\|, \quad t \geq t_0; \quad V_1(t_0) \leq \sqrt{\beta^2 + 1} \|R\|. \quad (8)$$

并且不难计算得:

$$\dot{V}_1(t) \leq -2V_1(t) + \delta(b\|x(t - \Delta(t))\| + c\|\dot{x}(t - \Delta(t))\|),$$

从而有:

$$\frac{d[V_1 e^{-r(t-s)}]}{ds} \leq \delta e^{-r(t-s)}(b\|x(s - \Delta(s))\| + c\|\dot{x}(s - \Delta(s))\|). \quad (9)$$

将上式从 t_0 到 t 两端积分, 得到:

$$V_1(t) \leq V_1(t_0) e^{-r(t-t_0)} + \delta \int_{t_0}^t e^{-r(t-s)}(b\|x(s - \Delta(s))\| + c\|\dot{x}(s - \Delta(s))\|) ds.$$

由(8)式有:

$$\|x(t)\| \leq K\|R\| + \frac{\delta}{\alpha} \int_{t_0}^t e^{-r(t-s)} (b\|x(s - \Delta(s))\| + c\|\dot{x}(s - \Delta(s))\|) ds, \quad t \geq t_0,$$

其中 $K = \sqrt{\beta^2 + 1}/\alpha$.

利用条件 I 可推得:

$$\|\dot{x}(t)\| \leq aK\|R\| + \frac{a\delta}{\alpha} \int_{t_0}^t e^{-r(t-s)} (b\|x(s - \Delta(s))\| + c\|\dot{x}(s - \Delta(s))\|) ds \\ + (b\|x(t - \Delta(t))\| + c\|\dot{x}(t - \Delta(t))\|), \quad t \geq t_0.$$

令: $P(t) = b\|x(t)\| + c\|\dot{x}(t)\|$, $t \geq t_0 - \Delta$, 则有:

$$P(t) \leq (b + ac + c)(k + 1)\|R\| + \frac{\delta(b + ac)}{\alpha} \int_{t_0}^t e^{-r(t-s)} P(s - \Delta(s)) ds \\ + cP(t - \Delta(t)), \quad t \geq t_0.$$

以下的证明类似定理 1, 此不赘述.

条件 IV: 存在 $V(t, x)$ 满足下列不等式:

$$(i) \quad (\alpha\|x\|)^2 \leq V(t, x) \leq (\beta\|x\|)^2, \quad \alpha > 0, \beta > 0;$$

$$(ii) \quad \left| \frac{\partial V(t, x)}{\partial x} \right| \leq 2\delta\sqrt{V(t, x)}, \quad \delta > 0;$$

$$(iii) \quad \frac{\partial V(t, x)}{\partial x} F[t, x(t), x(t), 0] + \frac{\partial V(t, x)}{\partial t} \leq -2rV, \quad r > 0.$$

条件 V: $\frac{\delta ac}{\alpha r} + \Delta b + c < 1$.

定理 4. 若条件 I、IV、V 成立, 则系统 (7) 的零解是在 C_1 空间中大范围指数稳定.

参 考 文 献

- [1] 钱学森、宋健, 工程控制论, 科学出版社, 1980.
- [2] 秦元勋、刘永清、王联, 带有时滞的动力系统的运动稳定性, 科学出版社, 1963.
- [3] Hale, J.K., Theory of Functional Differential Equations, Springer-Verlag, New York, 1977.
- [4] Эльстольц Л.Э., Вестн. Моск. ун-та., 5(1959), 65—71.
- [5] 斯力更, 数学学报, 17 (1974), 3, 197—204.
- [6] 李森林, 科学通报, 23 (1978), No. 2, 88—93.
- [7] Liao Xiaoxin, SCIENTIA SINICA, A (1986), No. 3, 225—240.
- [8] 斯力更, 章毅, 科学通报, 31 (1986), No. 20, 1527—1530.
- [9] 章毅, 科学通报, 31 (1986), No. 5.

THE STABILITY IN C_1 SPACE FOR NEUTRAL DELAY SYSTEMS

ZHANG YI ZHONG SHOUMING

(University of Electronic Science and Technology of China)

ABSTRACT

In this paper, the stability in C_1 space for neutral delay systems is studied. Some new sufficient conditions for stability in C_1 space are obtained.

Key words — Neutral type; Delay; Space; Stability.