

三七环二肽成分和人参内酰胺成分*

谭宁华¹, 王双明^{1, 2}, 杨亚滨¹, 何敏¹

(1 中国科学院昆明植物研究所植物化学与西部植物资源持续利用国家重点实验室, 云南 昆明 650204;

2 天津天士力制药集团有限公司药物研究所, 天津 300142)

Cyclodipeptides of *Panax notoginseng* and Lactams of *Panax ginseng* *

TAN Ning-Hua¹, WANG Shuang-Ming^{1, 2}, YANG Ya-Bin¹, HE Ming¹

(1 State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany,

Chinese Academy of Sciences, Kunming 650204, China;

2 Tianjin Tianshili Pharmaceutical Group Co. Ltd, Tianjin 300142, China)

Abstract: From the roots of *Panax notoginseng* fourteen cyclodipeptides **1** – **14** were isolated including one new compound (**1**), seven new natural compounds (**4** – **10**) and six known compounds (**2** – **3**, **11** – **14**) together with one known other compound **15**. The chemical structure of **1** was elucidated as cyclo-(Leu-Thr) based on spectral methods. From the roots of *Panax ginseng* five known lactams (**16** – **20**) including pyroglutamic acid were isolated together with butyric diacid, daucosterol and sucrose. The primary bioactivity test showed that pyroglutamic acid and its n-butyl derivative have weak Ca²⁺ antagonistic activity.

Key words: *Panax notoginseng*; *Panax ginseng*; Araliaceae; cyclodipeptides; lactams

关键词: 三七; 人参; 五加科; 环二肽; 内酰胺

中图分类号: Q 946 文献标识码: A 文章编号: 0253 – 2700(2003)03 – 0366 – 03

Panax notoginseng (Bur.) F. H. Chen and *Panax ginseng* C. A. Meyer (Araliaceae) are two famous Traditional Chinese Medicines. There are a lot of chemical studies on them, especially saponins. As one part of our investigation on cyclopeptides from higher plants with the TLC chemical detection method for cyclopeptides (Zhou and Tan, 2000) some cyclopeptides and lactams were detected in the EtOAc fractions of *P. notoginseng* and *P. ginseng* respectively, leading to isolate fifteen compounds including fourteen cyclodipeptides **1** – **14** from the roots of *P. notoginseng* and five known lactams including pyroglutamic acid (**16**), methyl pyroglutamate (**17**), ethyl pyroglutamate (**18**), isobutyl pyroglutamate (**19**), n-butyl pyroglutamate (**20**) from the roots of *P. ginseng*. Among them

* Foundationitem: This work was supported by grants from the National Natural Science Foundation of China (39500182) and Natural Science Fund of Yunnan Province (95C088Q)

Received date: 2002 – 11 – 03, Accepted date: 2003 – 01 – 09

作者简介: 谭宁华 (1963 –) 女, 重庆人, 博士、研究员, 主要从事天然产物化学和生物活性筛选研究。

Tel: +86 – 871 – 5223800. Fax: +86 – 871 – 5223228. E-mail: nhtan@mail.kib.ac.cn

compound **1** is one new compound, compounds **4–10** are new natural compounds, compounds **2–3** and **11–15** are known compounds. Compounds **2** and **11**, **3** and **4**, **10** and **15**, **12** and **13** are mixtures with 2:1, 1:1, 1:1, 2:1 ratios, respectively. Compounds **17–20** may be artifacts of compound **16** during isolation. The primary bioactivity test showed that pyroglutamic acid and its n-butyl derivative have weak Ca^{2+} antagonistic activity. Herein we report the structure elucidation of compound **1**.




Table 1 NMR data of **1** (in DMSO, 400 MHz for δ_{H} , 100 MHz for δ_{C})

No.	δ_{H}	δ_{C}
1	8.21 (1H, s) ^f	
2	4.00 (1H, m) ^g	52.8 (d) ^h
3	1.68 (1H, m), 1.55 (1H, m)	45.0 (t)
4	1.80 (1H, m)	23.3 (d)
5	0.84 (3H, d, 6.6)	23.1 (q)
6	0.82 (3H, d, 6.5)	21.5 (q)
7		168.6 (s) ⁱ
8	8.02 (1H, s) ^f	
9	5.02 (1H, d, 5.4) ^g	60.5 (d) ^h
10	4.00 (1H, m)	66.9 (d)
11	1.06 (3H, d, 6.5)	20.1 (q)
12		166.8 (s) ⁱ

^{a-d} Assignments may be reversed.

two amides NH at δ 8.02, 8.21, three methines CH at δ 4.00 – 5.02, one methine CH and one methene CH₂ at δ 1.55 – 1.80, three methyls CH₃ at δ 0.82 – 1.06, respectively. These facts indicated that **1** is a cyclodipeptide and composed of Leu(1eq) and Thr(1eq). The NMR data are shown in the Table 1. Therefore, the structure of **1**, a new cyclodipeptide, was elucidated as cyclo-(Leu-Thr).

With the same method as **1**, other thirteen cyclodipeptides were determined by spectral methods as following: **2** [cyclo-(Leu-Ile)], **3** [cyclo-(Leu-Val)], **4** [cyclo-(Ile-Val)], **5** [cyclo-(Leu-Ser)], **6** [cyclo-(Leu-Tyr)], **7** [cyclo-(Val-Pro)], **8** [cyclo-(Ala-Pro)], **9** [cyclo-(Phe-Tyr)], **10** [cyclo-(Phe-Ala)], **11** [cyclo-(Phe-Val)], **12** [cyclo-(Leu-Ala)], **13** [cyclo-(Ile-Ala)], **14** [cyclo-(Val-Ala)].

Acknowledgment: We are grateful to all members of the analytical group for spectrum measurement including NMR, MS, IR, UV in the State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences.

References:

Zhou J, Tan NH, 2000. Application of a new TLC chemical method for detection of cyclopeptides in plants [J]. *Chinese Science Bulletin*, 2000, **45** (20): 1825—1831

Compound 1 Colorless needles (CH₃OH), gave a negative ninhydrin reaction but positive after hydrolysis with 6 mol/L HCl (Zhou and Tan, 2000). Its molecular formula was determined as C₁₀H₁₈O₃N₂ by means of DEPT spectrum and FAB-MS in which its quasimolecular ion peak at m/z 215 (M + 1)⁺. The ¹³C NMR spectrum showed the presence of two amides CO at δ 166.8, 168.6, four methines CH at δ 66.9, 60.5, 52.8, 23.3, one methene CH₂ at δ 45.0, three methyls CH₃ at δ 23.1, 21.5, 20.1, respectively. The ¹H NMR spectrum showed the presence of