

Revista Mexicana de Astronomía y Astrofísica

Revista Mexicana de Astronomía y Astrofísica
Universidad Nacional Autónoma de México
rmaa@astroscu.unam.mx
ISSN (Versión impresa): 0185-1101
MÉXICO

2003

A. Rodríguez González / G. Tenorio Tagle / S. A. Silich
ON THE HIGH METALLICITY PRODUCED BY CSNRS
Revista Mexicana de Astronomía y Astrofísica, número 015
Universidad Nacional Autónoma de México
Distrito Federal, México
p. 274

Red de Revistas Científicas de América Latina y el Caribe, España y Portugal

Universidad Autónoma del Estado de México

reDalyC
LA BIBLIOTECA CIENTÍFICA EN LÍNEA
<http://redalyc.uaemex.mx>

ON THE HIGH METALLICITY PRODUCED BY CSNRs

A. Rodríguez-González,¹ G. Tenorio-Tagle,¹ and S. A. Silich¹

We present semi-analytic models for the hydrodynamic and chemical evolution of compact supernova remnants (cSNRs). Explosions of two progenitor stars of 15 and 25 M_{\odot} are explored. We find the metallicity of the cSNRs to be high ($1.0 \leq Z/Z_{\odot} \leq 6.0$) if oxygen is used as a tracer. These values are close to the metallicities of highly redshifted QSOs (Ferland et al. 1996; Hamann & Ferland 1999).

Spectra of highly redshifted QSOs present broad line emission. The metallicities of the associated fast-moving gas derived from the spectral analyses are surprisingly high for such young objects and fall into the range $Z \sim 1$ to $9 Z_{\odot}$ (Hamann & Ferland 1999). This implies an effective contamination of the interstellar gas within a short timescale at the pre-QSO phase. QSO gas metal enrichment is usually related to violent star formation (Ferland et al. 1996). However, this model is limited by the short time required to produce enough metals and mix them with the ambient medium. We analyze another approach based on the possible association of QSOs with cSNRs (Terlevich et al. 1992).

We developed a semi-analytical model for cSNR evolution and examined the explosion of 15 M_{\odot} and 25 M_{\odot} progenitors into a dense (10^7 cm^{-3}) circumstellar medium (CSM) (Rodríguez-González 2002). The density and the velocity of the ejecta were taken from Franco et al. (1991):

$$\rho_{\text{ej}}(r \geq R_c, t) = \frac{M_{\text{ej}}}{4\pi r^3 \ln[R(t)/R_c]},$$

$$V_{\text{ej}}(r \geq R_c, t) = \frac{r - R_c}{R(t) - R_c} V_m,$$

and $\rho_{\text{ej}}(r < R_c, t) = V_{\text{ej}}(r < R_c, t) = 0$, where $R(t) = R_m + V_m t$, R_m is the initial radius, V_m is the maximum ejecta velocity, and R_c is the inner radius of the ejecta.

In both cases, soon after the explosion cooling becomes important and dense shells form after ~ 1 yr of evolution. Then, we calculate the characteristic diffusion time $t_d = L^2/D$ needed to mix the ejected




Fig. 1. The cSNR metallicity as function of time if oxygen is used as a tracer.

metals with the swept-up CSM. The thickness, L , of the cSNR has been taken from our hydrodynamical model and two different approximations for the diffusion coefficient, D , have been used. The metallicity of the cSNR has been calculated from

$$Z_A(t)/Z_{\odot} = \frac{M_{A,\text{ej}}(t)/Z_{A,\odot} + Z_{\text{CSM}}M_{\text{CSM}}(t)}{M_{\text{CSM}}(t) + \chi M_{\text{ej}}(t)},$$

where $M_{\text{CSM}}(t)$ and $\chi M_{\text{ej}}(t)$ are masses of the swept-up CSM and the ejecta, respectively, and $M_{A,\text{ej}}(t)$ is the mass of the element A that has been accumulated within a shell.

The calculations show that effective metal diffusion occurs soon after dense shell formation. The resultant shell metallicity was found to be within the range $1.0 \leq Z/Z_{\odot} \leq 6.0$ if oxygen is used as a tracer and $0.65 \leq Z/Z_{\odot} \leq 1.3$ if iron is used as a tracer of the gas metallicity. We speculate that the observed QSO metallicities could be related to the cSNRs.

The authors feel honored to have attended this meeting and acknowledge support from CONACYT grant 36132-E.

REFERENCES

- Ferland, G. J., et al. 1996, ApJ, 461, 683
 Franco, J., Tenorio-Tagle, G., Bodenheimer, P., Różyczka, M. 1991, PASP, 103, 803
 Hamann, F., & Ferland, G. J. 1999, ARA&A, 37,487
 Rodríguez-González, A. 2002, MSc Thesis, INAOE
 Terlevich, R. J., Tenorio-Tagle, G., Franco, J., & Melnick, J. 1992, MNRAS, 255, 713

¹Instituto Nacional de Astrofísica Óptica y Electrónica, Apdo. Postal 51, 72000 Puebla, Puebla, México (ary@inaoep.mx).