

ON THE COMPLETE INTEGRABILITY OF NONLINEAR DYNAMICAL SYSTEMS ON DISCRETE MANIFOLDS WITHIN THE GRADIENT-HOLONOMIC APPROACH

YAREMA A. PRYKARPATSKY, NIKOLAI N. BOGOLUBOV (JR.), ANATOLIY K. PRYKARPATSKY,
AND VALERIY H. SAMOYLENKO

ABSTRACT. A gradient-holonomic approach for the Lax type integrability analysis of differential-discrete dynamical systems is devised. The asymptotical solutions to the related Lax equation are studied, the related gradient identity is stated. The integrability of a discrete nonlinear Schredinger type dynamical system is treated in detail.

1. PRELIMINARY NOTIONS AND DEFINITIONS

Consider an infinite dimensional discrete-manifold $M \subset l_2(\mathbb{Z}; \mathbb{C}^m)$ for some integer $m \in \mathbb{Z}_+$ and a nonlinear dynamical system on it in the form

$$(1.1) \quad du/dt = K[u],$$

where $u \in M$ and $K : M \rightarrow T(M)$ is a Frechet smooth nonlinear local functional on M and

$t \in \mathbb{R}$ is the evolution parameter. As an example of dynamical system (1.1) on a discrete manifold $M \subset l_2(\mathbb{Z}; \mathbb{C}^2)$ one can consider the following well-known [5, 10] discrete nonlinear Schrödinger equation:

$$(1.2) \quad \left. \begin{aligned} d\psi_n/dt &= i(\psi_{n+1} - 2\psi_n + \psi_{n-1}) - i\psi_n^* \psi_n (\psi_{n+1} + \psi_{n-1}) \\ d\psi_n^*/dt &= -i(2\psi_n^* - \psi_{n+1}^* - \psi_{n-1}^*) + i\psi_n^* \psi_n (\psi_{n+1}^* + \psi_{n-1}^*) \end{aligned} \right\} := K_n[\psi, \psi^*],$$

having interesting applications in diverse physics plasma investigations.

For studying the integrability properties of differential-difference dynamical system (1.1) we will develop below a gradient-holonomic scheme before devised in [6, 14, 12, 7] for nonlinear dynamical systems defined on spatially one-dimensional functional manifolds and extended in [11] on the case of discrete manifolds.

Denote by (\cdot, \cdot) the standard bi-linear form on the space $T^*(M) \times T(M)$ naturally induced by that existing in the Hilbert space $l_2(\mathbb{Z}; \mathbb{C}^m)$. Having denoted by $\mathcal{D}(M)$ smooth functionals on M , for any functional $\gamma \in \mathcal{D}(M)$ one can define the gradient $\text{grad } \gamma[u] \in T^*(M)$ as follows:

$$(1.3) \quad \text{grad } \gamma[u] := \gamma'^* [u] \cdot 1,$$

where the dash-sign "′" means the corresponding Frechet deritative and the star-sign "∗" means the conjugation naturally related with the bracket on $T^*(M) \times T(M)$.

Definition 1.1. A linear smooth operator $\vartheta : T^*(M) \times T(M)$ is called implectic, on the manifold M , if the bi-linear bracket

$$(1.4) \quad \{\cdot, \cdot\}_\vartheta := (\langle \text{grad}(\cdot), \vartheta \text{grad}(\cdot) \rangle)$$

is Poissonian [1, 2] on the space of functionals $\mathcal{D}(M)$.

This means, in particular, that bracket (1.3) satisfies the standard Jacobi identity on $\mathcal{D}(M)$.

Definition 1.2. A linear smooth operator $\vartheta : T^*(M) \times T(M)$ is called Nötherian subject to the nonlinear dynamical system (1.1), if the following condition

$$(1.5) \quad L_K \vartheta = \vartheta' K - \vartheta K'^* - K' \vartheta = 0$$

1991 *Mathematics Subject Classification.* 35A30, 35G25, 35N10, 37K35, 58J70, 58J72, 34A34; PACS: 02.30.Jr, 02.30.Hq.

Key words and phrases. gradient identity, conservation laws, asymptotical analysis, implectic structures, Lax type representation, finite-dimensional reduction, Liouville integrability.

holds identically on the manifold M , where we denoted by L_K the corresponding Lie-derivative along the vector field $K : M \rightarrow T(M)$.

Assume now that the mapping $\vartheta : T^*(M) \times T(M)$ is invertible, that is there exists the inverse mapping $\vartheta^{-1} := \Omega : T^*(M) \times T(M)$, and called symplectic. It then follows easily from (1.5) then the condition

$$(1.6) \quad L_K \Omega = \Omega' K + \Omega K' + K'^* \Omega = 0$$

hold identically on M . Having now assumed that the manifold $M \subset l_2(\mathbb{Z}; \mathbb{C}^2)$ is endowed with a smooth implectic structure $\vartheta : T^*(M) \times T(M)$ one can define the Hamiltonian system

$$(1.7) \quad du/dt := -\vartheta \text{grad } H[u],$$

corresponding to a Hamiltonian function $H \in \mathcal{D}(M)$. As a simple corollary of definition (1.7) one obtains that the dynamical system

$$(1.8) \quad -\vartheta \text{grad } H[u] := K[u]$$

satisfies the Nötherian conditions (1.5). Keeping in mind the study of the integrability problem [2, 4, 12, 8] subject to discrete dynamical system (1.1), we need to construct a priori given set of invariant with respect to it functions suitable on M called conservation laws commuting to each other with respect to the Poisson bracket (1.3). The following Lax criterion [3, 6, 12] proves to be very useful.

Lemma 1.3. *Any symmetric with respect to the bracket (\cdot, \cdot) smooth solution $\varphi \in T^*(M)$ to the Lax equation*

$$(1.9) \quad L_K \varphi = d\varphi/dt + K'^* \varphi = 0, \quad \varphi' = \varphi'^*$$

corresponds to the conservation law

$$(1.10) \quad \gamma := \int_0^1 d\lambda(\varphi[u\lambda], u).$$

The expression easily obtains from the well-known Volterra homology expressions:

$$(1.11) \quad \gamma = \int_0^1 \frac{d\gamma[u\lambda]}{d\lambda} d\lambda = \int_0^1 d\lambda(\gamma'[u\lambda] \cdot u, 1) = \int_0^1 d\lambda(\gamma'^*[u\lambda] \cdot 1, u) = \int_0^1 d\lambda(\text{grad } \gamma[u\lambda], u)$$

and

$$(1.12) \quad (\text{grad } \gamma[u])' = (\text{grad } \gamma[u])'^*,$$

holding identically on M , since then one ensues that there exists a function $\gamma \in \mathcal{D}(M)$ such that

$$(1.13) \quad L_K \gamma = 0 \text{ and } \text{grad } \gamma[u] = \varphi[u]$$

for any $u \in M$. The Lax lemma naturally arises from the following generalized Nöther type lemma.

Lemma 1.4. *Let a smooth element $\varphi \in T^*(M)$ satisfy the Nöther condition*

$$(1.14) \quad L_K \varphi = d\varphi/dt + K'^* \varphi = \nabla \mathcal{L}_\varphi,$$

where we denoted $\nabla \mathcal{L}_\varphi := \text{grad } \mathcal{L}_\varphi$ for some smooth functional $\mathcal{L}_\varphi \in \mathcal{D}(M)$. Then the following Hamiltonian representation

$$(1.15) \quad K = -\vartheta \text{grad } H_\vartheta,$$

where

$$(1.16) \quad \vartheta := \varphi' - \varphi'^*, \quad H_\vartheta := (\varphi, K) - \mathcal{L}_\varphi,$$

and the Hamiltonian function $H_\vartheta \in \mathcal{D}(M)$.

It is easy to see that Lemma (1.3) follows from Lemma (1.4) if the conditions $\varphi' = \varphi'^*$ and $\mathcal{L}_\varphi = 0$ are imposed in (1.16).

Assume now that equation (1.14) allows an additional not symmetric smooth solution $\phi \in T^*(M)$:

$$(1.17) \quad L_K \phi = d\phi/dt + K'^* \phi = \text{grad } \mathcal{L}_\phi.$$

This means that our system (1.1) is bi-Hamiltonian:

$$(1.18) \quad -\vartheta \text{ grad } H_\vartheta = K = -\eta \text{ grad } H_\eta,$$

where, by definition,

$$(1.19) \quad \eta := \phi' - \phi'^*, \quad H_\eta = (\phi, K) - \mathcal{L}_\phi.$$

Definition 1.5. One says that two implectic structures $\vartheta, \eta : T^*(M) \rightarrow T(M)$ on M are compatible [9, 17, 6, 8], if for any $\lambda, \mu \in \mathbb{R}$ the linear combination $\lambda\vartheta + \mu\eta : T^*(M) \rightarrow T(M)$ will be too implectic on M .

It is easy to derive that this condition is satisfied if, for instance, there exist the inverse operator $\vartheta^{-1} : T(M) \rightarrow T^*(M)$ and the expression $\eta(\vartheta^{-1}\eta) : T^*(M) \rightarrow T(M)$ is also implectic on M .

Concerning the integrability problem posed for the infinite-dimensional dynamical system (1.1) on the discrete manifold M it is, in general, necessary, but not enough [4, 6, 12], to prove the existence of an infinite hierarchy of commuting to each other with respect to the Poissonian structure (1.3) conservation laws.

Since in the case of the Lax type integrability almost of (1.1) there exist compatible implectic structures and related hierarchies of conservation laws, we will further constrain our analysis by devising an integrability algorithm under the a priori assumption that a given nonlinear dynamical system (1.1) on the manifold M is Lax type integrable. This means that it possesses a related Lax type representation in the following, generally written form:

$$(1.20) \quad \Delta f_n := f_{n+1} = l_n[u; \lambda] f_n,$$

where $f := \{f_n \in \mathbb{C}^r : n \in \mathbb{Z}\} \subset l_2(\mathbb{Z}; \mathbb{C}^r)$ for some integer $r \in \mathbb{Z}_+$ and $l_n[u; \lambda] \in \text{End } \mathbb{C}^r$ in (1.20) is a local matrix-valued functional on M , depending on the "spectral" parameter $\lambda \in \mathbb{C}$, invariant with respect to our dynamical system (1.1).

Taking into account that the Lax representation (1.20) is "local" with respect to the discrete variable $n \in \mathbb{Z}$, we will assume for convenience, that our manifold $M := M_{(N)} \subset l_\infty(\mathbb{Z}; \mathbb{C}^m)$ is periodic with respect to the discrete index $n \in \mathbb{Z}$, that is for any $n \in \mathbb{Z}$ and $\lambda \in \mathbb{C}$

$$(1.21) \quad l_n[u; \lambda] = l_{n+N}[u; \lambda]$$

for some integer $N \in \mathbb{Z}_+$. In this case the smooth functionals on $M(N)$ can be represented as

$$(1.22) \quad \gamma := \sum_{n=0}^{N-1} \gamma_n[u]$$

for some local "Frechet smooth densities" $\gamma_n : M_{(N)} \rightarrow \mathbb{C}$, $n = \overline{0, N-1}$.

2. THE INTEGRABILITY ANALYSIS: GRADIENT-HOLONOMIC SCHEME

Consider the representation (1.20) and define its fundamental solution $F_{n,n_0}^{(\lambda)} \in \text{Aut}(\mathbb{C}^r)$, $n \in \mathbb{Z}$, satisfying the condition $F_{n,n_0}^{(\lambda)} \Big|_{n=n_0} = \mathbf{1}$ for some arbitrarily fixed index $n_0 \in \mathbb{Z}$ and all $\lambda \in \mathbb{C}$:

$$(2.1) \quad F_{n+1,n_0}^{(\lambda)} = l_n[u; \lambda] F_{n,n_0}^{(\lambda)}.$$

Then the matrix function

$$(2.2) \quad S_{n_0}(\lambda) := F_{n_0+N,n_0}(\lambda) \in \text{Aut}(\mathbb{C}^r)$$

is called the monodromy matrix for the linear equation (1.21) and satisfies at $n_0 = n \in \mathbb{Z}$ the following Novikov-Lax type relationship:

$$(2.3) \quad S_{n+1}(\lambda) l_n = l_n S_n(\lambda).$$

It easy to compute that $S_n(\lambda) := \prod_{k=\overline{0, N-1}} l_{n+k}[u; \lambda]$ owing to the periodical condition (1.21).

Construct the following generating functional:

$$(2.4) \quad \bar{\gamma}(\lambda) := \text{tr} S_n(\lambda)$$

and assume that there exists its asymptotical expansion

$$(2.5) \quad \bar{\gamma}(\lambda) \simeq \sum_{j \in \mathbb{Z}_+} \bar{\gamma}_j \lambda^{-j+j_0}$$

as $|\lambda| \rightarrow \infty$ for some fixed $j_0 \in \mathbb{Z}_+$. Then, owing to the evident condition

$$(2.6) \quad \frac{d}{dn} \bar{\gamma}(\lambda) = 0$$

for all $n \in \mathbb{Z}$, where we put, by definition, the "discrete" derivative

$$(2.7) \quad d/dn := \Delta - 1,$$

we obtain that all functionals $\bar{\gamma}_j \in \mathcal{D}(M_{(N)})$, $j \in \mathbb{Z}_+$, are independent of the discrete index $n \in \mathbb{Z}$.

Assume now additionally that the following natural condition holds: the gradient vector

$$(2.8) \quad \varphi(\lambda) := \text{grad} \bar{\gamma}(\lambda) = \text{tr}(l'^* S_n(\lambda))$$

satisfies, owing to (2.3), for all $\lambda \in \mathbb{C}$ the next "spectral"-gradient relationship:

$$(2.9) \quad z(\lambda) \vartheta \varphi(\lambda) = \eta \varphi(\lambda),$$

where $z : \mathbb{C} \rightarrow \mathbb{C}$ is some meromorphic mapping, ϑ and $\eta : T^*(M_{(N)}) \rightarrow T(M_{(N)})$ are compatible implectic on the manifold $M_{(N)}$ and Nötherian operators with respect to the dynamical system (1.1).

As a corollary of the above condition one follows easily that the generating functional $\bar{\gamma} \in \mathcal{D}(M_{(N)})$ is a conservation law to (1.1), there by giving rise, owing to (2.5), to the existence of the infinite hierarchy of conservation laws $\bar{\gamma}_j \in \mathcal{D}(M_{(N)})$, $j \in \mathbb{Z}_+$, on the periodic manifold $M_{(N)}$.

The latter property, as characteristic concerning the Lax type integrability of the nonlinear dynamical system (1.1), can be effectively implemented into the direct scheme of our analysis if to take into account the defining relationships (1.20) and (2.2). Namely, the following theorem holds.

Theorem 2.1. . *The Lax equation (1.9) allows the following asymptotical as $|\lambda| \rightarrow \infty$ periodical solution $\varphi(\lambda) \in T^*(M_{(N)})$, where for all $n \in \mathbb{Z}$*

$$(2.10) \quad \varphi_n(\lambda) \simeq a_n(\lambda) \exp[\omega(t; \lambda)] \prod_{j=0}^n \sigma_j(\lambda),$$

$$(2.11) \quad \begin{aligned} a_n(\lambda) & : = (1, a_{(1),n}[u; \lambda], a_{(2),n}[u; \lambda], \dots, a_{(m-1),n}[u; \lambda])^\tau, \\ a_{(k),n}(\lambda) & \simeq \sum_{s \in \mathbb{Z}_+} a_{(k),n}^{(s)}[u] \lambda^{-s+\bar{a}}, \sigma_j(\lambda) \simeq \sum_{s \in \mathbb{Z}_+} a_j^{(s)}[u] \lambda^{-s+\bar{\sigma}}, \end{aligned}$$

$k = \overline{1, m-1}$ and $\omega(t; \cdot) : \mathbb{C} \rightarrow \mathbb{C}$, $t \in \mathbb{R}$, is some dispersion function. Moreover the functional $\gamma(\lambda) := \sum_{n=\overline{0, N}} \ln \sigma_n[u; \lambda] \in \mathcal{D}(M_{(N)})$ is a generating function of conservation laws to the dynamical system (1.1).

Proof. The condition $L_K \vartheta = 0 = L_K \eta$ gives rise to the equality $L_K \varphi = 0$, coinciding with (1.9). The latter, owing to Lemma (1.3) and (2.8) means that the functional (2.4) is a conservation law to our dynamical system (1.1). Based now on the expression (2.2) and equation (1.20) we arrive at the solution representation (2.10) to the Lax equation (1.9). Making use of the periodicity of the solution (2.10) we obtain that the functional

$$(2.12) \quad \gamma(\lambda) := \sum_{n=\overline{0, N}} \ln \sigma_j[u; \lambda] \in \mathcal{D}(M_{(N)})$$

is a conservation law to (1.1), finishing the proof. \square

Thus, if the start the Lax type integrability analysis of a priori given nonlinear dynamical system (1.1), it is necessary, as the first step, to study the asymptotical solutions (2.10) to the corresponding Lax equation (1.9) and construct a related hierarchy of conservation laws in the form

$$(2.13) \quad \gamma_j := \sum_{j=0, N-1} \gamma_j[u],$$

taking into account the expansions of (2.11) and the definition (2.12).

Remark 2.2. It is easy to observe that owing to the arbitrariness of the period $N \in \mathbb{Z}_+$ of the manifold $M_{(N)}$, all of the finite-sum expressions obtained above are generalized to the corresponding infinite dimensional manifold $M \subset l_2(\mathbb{Z}; \mathbb{C}^m)$, if the corresponding infinite series persist to be convergent.

Since our dynamical system (1.1) under the above conditions is a bi-Hamiltonian flow on the manifold $M_{(N)}$, as the next step of the related compatible implectic (or symplectic) structures, satisfying, respectively, the equality (1.5) or (1.6). Before doing this, we need to formulate the following lemma.

Lemma 2.3. *All functionals (2.13) are commuting to each other with respect to both implectic structures $\vartheta, \eta : T^*(M_{(N)}) \rightarrow T(M_{(N)})$, satisfying the "spectral"-gradient relationship(2.14).*

Proof. Based on the representations (2.10) and (2.8) one obtains that there holds the asymptotical as $|\lambda| \rightarrow \infty$ relationship

$$(2.14) \quad \ln \bar{\gamma}(\lambda) \simeq \gamma(\lambda).$$

Since the generative function of conservation laws $\bar{\gamma}(\lambda) \in \mathcal{D}(M_{(N)})$ is owing to the "spectral"-gradient relationship (2.9) and the Poisson bracket definition (1.3), commuting the same, evidently, holds to each other, owing to the relationship (2.14), for the generalizing function $\gamma(\lambda) \in \mathcal{D}(M_{(N)})$, finishing the proof. \square

Proceed now to constructing the related to dynamical system (1.1) implectic structures $\vartheta, \eta : T^*(M_{(N)}) \rightarrow T(M_{(N)})$. Note here, that these implectic structures are Nötherian also for the whole hierarchy of dynamical systems

$$(2.15) \quad du/dt_j := -\vartheta \text{ grad } \gamma_j := K^{(j)}[u],$$

where $t_j \in \mathbb{R}, j \in \mathbb{Z}_+$, are the corresponding evolution parameters, and which commute to each other on the manifold $M_{(N)}$. The latter makes it possible to apply Lemma (1.4) to any of the dynamical systems (2.15), if the related vector fields commuting with (1.1) are supposed to be found before.

To solve analytically the equation (1.14) subject to an element $\varphi \in T^*(M_{(N)})$, we can, in the case of a polynomial dynamical system (1.1), make use of the well known asymptotical small parameter method [14]. Applying this approach it is necessary to take into account the following expansions at zero - element $\bar{u} = 0 \in M_{(N)}$ with respect to the small parameter $\mathbb{C} \ni \mu \rightarrow \infty$:

$$(2.16) \quad \begin{aligned} u & : = \mu u^{(1)}, \quad \varphi[u] = \varphi^{(0)} + \mu \varphi^{(1)}[u] + \mu^2 \varphi^{(2)}[u] + \dots, \\ d/dt & = d/dt_0 + \mu d/dt_1 + \mu^2 d/dt_2 + \dots, \\ K[u] & = \mu K^{(1)}[u] + \mu^2 K^{(2)}[u] + \dots, \\ K'[u] & = K'_0 + \mu K'_1[u] + \mu^2 K'_2[u] + \dots, \\ \nabla L[u] & = \nabla L^{(0)} + \mu \nabla L^{(1)}[u] + \mu^2 \nabla L^{(2)}[u] + \dots \end{aligned}$$

having solved the corresponding linear nonuniform functional equations

$$(2.17) \quad \begin{aligned} d\varphi^{(0)}/dt_0 + K_0'^* \varphi^{(0)} & = \nabla L^{(0)}, \\ d\varphi^{(1)}/dt_0 + K_0'^* \varphi^{(1)} & = \nabla L^{(1)} - K_0'^* \varphi^{(0)}, \\ d\varphi^{(2)}/dt_0 + K_0'^* \varphi^{(2)} & = \nabla L^{(2)} - K_1'^* \varphi^{(1)} - K_2'^* \varphi^{(0)} \end{aligned}$$

by means the standard Fourier transform applied to the suitable N -periodical functions, one can obtain the related K_0 -implectic structure as follows

$$(2.18) \quad \vartheta^{-1} = \varphi^{(0),'} - \varphi^{(0),'*} + \mu(\varphi^{(1),'} - \varphi^{(1),'*}) + \dots$$

and put in (2.18) at the end $\mu = 1$.

Another direct way of obtaining an implectic operator $\vartheta : T^*(M_{(N)}) \rightarrow T(M_{(N)})$ for (1.1) is to solve by means of the same asymptotical small parameter approach the Nötherian equation (1.5), having reduced it to the following set of linear nonuniform equations:

$$(2.19) \quad \begin{aligned} \frac{d}{dt_0}(\vartheta_0\varphi^{(0)}) &= K'_0(\vartheta_0\varphi^{(0)}), \\ \frac{d}{dt_0}(\vartheta_1\varphi^{(0)}) &= K'_0(\vartheta_1\varphi^{(0)}) + \vartheta_0K'_1{}^*\varphi^{(0)} + K'_1\vartheta_0\varphi^{(0)}, \\ \frac{d}{dt_0}(\vartheta_2\varphi^{(0)}) &= K'_0(\vartheta_2\varphi^{(0)}) - \varphi^{(0)'}K^1 + \vartheta_0K'_2{}^*\varphi^{(0)} + \\ &\quad + \vartheta_1K'_1{}^*\varphi^{(0)} + \vartheta_2K'_0{}^*\varphi^{(0)} + K'_1\vartheta_1\varphi^{(0)} + K'_2\vartheta_0\varphi^{(0)}. \end{aligned}$$

Based now on the analytical expressions for actions $\vartheta_j : \varphi^{(0)} \rightarrow \vartheta_j\varphi^{(0)}$, $j \in \mathbb{Z}_+$, one can easily retrieve them in operator form, giving rise to the implectic operator

$$(2.20) \quad \vartheta = \vartheta_0 + \mu\vartheta_1 + \mu^2\vartheta_2 + \dots$$

if to put at the end of calculations $\mu = 1$. Similarly one can construct the second implectic operator $\eta : T^*(M_{(N)}) \rightarrow T(M_{(N)})$ for the nonlinear dynamical system (1.1).

Resuming up all this analysis described above, we can formulate the following proposition.

Proposition 2.4. *Let a nonlinear dynamical system (1.1) on a discrete manifold M allow a nontrivial asymptotical as $|\lambda| \rightarrow \infty$ symmetric solution $\varphi \in T^*(M)$ to the Lax equation (1.9) in the form (2.10), generating an infinite hierarchy of nontrivial functionally independent conservation laws (2.12). Then this dynamical system is a Lax type integrable bi-Hamiltonian flow on M with respect to two compatible implectic structures $\vartheta, \eta : T^*(M) \rightarrow T(M)$, whose adjoint Lax type representation*

$$(2.21) \quad d\Lambda/dt = [K'^*, \Lambda],$$

where $\Lambda := \vartheta^{-1}\eta$ is the so-called recursion operator, can be transformed to the standard discrete Lax type form as

$$(2.22) \quad \frac{dl_n}{dt} = [p_n(l), l_n] + \frac{d}{dn}p_n(l)$$

for all $n \in \mathbb{Z}$, where $p_n(l) \in \text{End}\mathbb{C}^r$ describes the related to the discrete linear Lax type problem (1.20) temporal evolution

$$(2.23) \quad df_n/dt = p_n(l)f_n$$

for $f_n \in \mathbb{C}^r$, $n \in \mathbb{Z}$.

Remark 2.5. Based on the property that all Hamiltonian flows (2.15) commute to each other and with dynamical system (1.1) and using the fact that they possess the same implectic and compatible (ϑ, η) -pair, the analytical algorithm described above can be successfully applied to any other flow, commuting with (1.1).

Concerning the discrete linear Lax type problem (1.20), it can be constructed by means of the gradient-holonomic algorithm, first devised in [6, 7, 12] for studying the integrability nonlinear dynamical systems on functional manifolds. Namely, making use of the preliminary found analytical expressions for the related compatible implectic structures $\vartheta, \eta : T^*(M) \rightarrow T(M)$ on the manifold M and using the fact that the recursion operator $\Lambda := \vartheta^{-1}\eta : T^*(M) \rightarrow T^*(M)$ satisfies the dual Lax type commutator equality

$$(2.24) \quad d\Lambda/dt = [\Lambda, K'^*],$$

one can retrieve the standard "spectral" Lax representation for (2.24) by means of suitably derived algebraic relationships. As a corollary of Proposition (2.4), formulated above, one can claim that the existence of a nontrivial asymptotic as $|\lambda| \rightarrow \infty$ solution to the Lax equation (1.9) can serve as an effective Lax type integrability criterion for a given nonlinear dynamical system (1.1) on the manifold M .

3. THE BOGOYAVLENSKY-NOVIKOV FINITE-DIMENSIONAL REDUCTION

Assume that our dynamical system (1.1) on the periodic manifold $M_{(N)}$ is Lax type integrable and possesses two compatible implectic structures $\vartheta, \eta : T^*(M_{(N)}) \rightarrow T(M_{(N)})$. Thus, we have the nonlinear finite-dimensional dynamical system

$$(3.1) \quad du_n/dt := K_n[u] = -\vartheta \operatorname{grad} H_n[u]$$

for indexes $n = \overline{0, N-1}$ owing to its N -periodicity. The finite dimensional dynamical system (3.1) can be equivalently considered as that on the finite-dimensional space $M_{(N)} \simeq (\mathbb{C}^m)^N$ parametrized by any integer index $n \in \mathbb{Z}$, and whose Liouville integrability is of the next our analysis. To proceed with studying the flow (3.1) on the manifold $M_{(N)}$, we will make use of the Bogoyavlensky-Novikov [15, 4] reduction scheme, first adapted for discrete dynamical systems in [11, 6].

Let $\Lambda(M_{(N)}) := \bigoplus_{j \in \mathbb{Z}_+} \Lambda^j(M_{(N)})$ be the standard Grassmann algebra [2, 6, 12] of differential forms on the manifold $\overline{M}_{(N)}$. Then the following differential complex

$$(3.2) \quad \Lambda^0(M_{(N)}) \xrightarrow{d} \Lambda^1(M_{(N)}) \rightarrow \dots \xrightarrow{d} \Lambda^j(M_{(N)}) \xrightarrow{d} \Lambda^{j+1}(M_{(N)}) \xrightarrow{d} \dots,$$

where $d : \Lambda(M_{(N)}) \rightarrow \Lambda(M_{(N)})$ is the external differentiation, is finite and exact. Since the discrete "derivative" $D_n := \Delta - 1$ commutes with the differentiation $d : \Lambda(M_{(N)}) \rightarrow \Lambda(M_{(N)})$, $[D_n, d] = 0$ for all $n = \overline{0, N-1}$, and for any element $a \in \Lambda^0(M_{(N)})$

$$(3.3) \quad \operatorname{grad} \left(\sum_{n=0}^{N-1} \frac{d}{dn} a_n[u] \right) = 0$$

one can formulate the following Gelfand-Dikiy type [16] lemma.

Lemma 3.1. *4.1 . Let $\mathcal{L}[u] \in \Lambda^0(M_{(N)})$ be a Frechet smooth local Lagrangian functional on the manifold $M_{(N)}$. Then there exists a 1-form $\alpha^{(1)} \in \Lambda^1(M_{(N)})$ such that the equality*

$$(3.4) \quad d\mathcal{L}_n[u] = \langle \operatorname{grad} \mathcal{L}_n[u], du_n \rangle + D_n \alpha^{(1)}[u]$$

is satisfied for all $n \in \mathbb{Z}$.

Proof. We can easily observe that

$$(3.5) \quad \begin{aligned} d\mathcal{L}_n[u] &= \sum_{n=0}^{N-1} \left\langle \frac{\partial \mathcal{L}_n[u]}{\partial u_{n+j}}, du_{n+j} \right\rangle = \sum_{n=0}^{N-1} \left\langle \frac{\partial \mathcal{L}_n[u]}{\partial u_{n+j}}, \Delta^j du_n \right\rangle = \\ &= \left\langle \sum_{n=0}^{N-1} \Delta^{-j} \frac{\partial \mathcal{L}_n[u]}{\partial u_{n+j}}, du_n \right\rangle + D_n \left(\sum_{n=0}^{N-1} \langle p_j, du_{n+j} \rangle \right), \end{aligned}$$

where, by definition, for $j = \overline{0, N-1}$

$$(3.6) \quad p_j := \sum_{k=0}^{N-1} \Delta^{-k} \frac{\partial \mathcal{L}_n[u]}{\partial u_{n+j+k+1}}.$$

Having denoted the expression

$$(3.7) \quad \operatorname{grad} \mathcal{L}_n[u] := \sum_{j=0}^{N-1} \Delta^{-j} \frac{\partial \mathcal{L}_n[u]}{\partial u_{n+j}},$$

one obtains the result (3.4), where

$$(3.8) \quad \alpha^{(1)}[u] := \sum_{j=0}^{N-1} \langle p_j, du_{n+j} \rangle$$

is the corresponding 1-form on the manifold $M_{(N)}$. □

Applying now the d -differentiation to expression (3.4) we obtain that

$$(3.9) \quad -D_n \omega^{(2)}[u] = \langle d \operatorname{grad} \mathcal{L}_n[u], \wedge du_n \rangle$$

for any $n \in \mathbb{Z}$, where the 2-form

$$(3.10) \quad \omega^{(2)}[u] := d\alpha^{(1)}[u]$$

is nondegenerate on $M_{(N)}$, if the Hessian $\partial_n^2 \mathcal{L}[u]/\partial^2 u_n$ is nondegenerate too.

Assume now that the submanifold

$$(3.11) \quad \bar{M}_{(N)} := \{\text{grad } \mathcal{L}_n[u] = 0; u \in M\},$$

where, by definition, the Lagrangian functional

$$(3.12) \quad \mathcal{L} := -\gamma_{N+1} + \sum_{k=0}^N c_k \gamma_k$$

and $\gamma_k \in \mathcal{D}(M)$, $k = \overline{0, N+1}$, are the suitable nontrivial conservation laws for the dynamical system (1.1), constructed before. As a result of (3.11) and (3.9) we obtain that closed 2-form $\omega_n^{(2)} \in \Lambda^2(\bar{M}_{(N)})$ is invariant with respect to the index $n \in \mathbb{Z}$ on the manifold $\bar{M}_{(N)}$. Moreover, the submanifold (3.11) is also invariant both with respect to the index $n \in \mathbb{Z}$ and the evolution parameter $t \in \mathbb{R}$. Really, for any $n \in \mathbb{Z}$ the Lie derivative

$$(3.13) \quad L_K \text{grad } \mathcal{L}_n = (\text{grad } \mathcal{L}_n)' K + K'^* (\text{grad } \mathcal{L}_n) = 0,$$

since owing to the functional $\mathcal{L} \in \mathcal{D}(M_{(N)})$ is a conservation law for the dynamical system (1.1), whose gradient satisfies the Lax condition (1.9). Applying now the Lie derivative L_K to the expression $D_n \text{grad } \mathcal{L}_n[u]$, $n \in \mathbb{Z}$, we easily obtain, owing to (3.13), that if $D_n \text{grad } \mathcal{L}_n[u] = 0$, $n \in \mathbb{Z}$,

at $t = 0$, then

$$(3.14) \quad D_n \text{grad } \mathcal{L}_n[u] = 0$$

for all $t \in \mathbb{R}$ and $n \in \mathbb{Z}$. Thus, we have reduced via Bogoyavlensky-Novikov the initially Lax type given integrable dynamical system (1.1) upon the invariant submanifold $\bar{M}_{(N)}$.

A question arises: how are related the dynamical system (1.1), reduced upon the submanifold $M_{(N)}$, and the dynamical system (1.1), reduced upon the finite dimensional submanifold $\bar{M}_{(N)} \subset M$. To analyze these reductions we will consider the following equality:

$$(3.15) \quad \langle \text{grad } \mathcal{L}_n[u], K_n[u] \rangle = -D_n h^{(t)}[u],$$

for some local functional $h^{(t)}[u] \in \Lambda^0(M)$, following from the conditions (3.3) and (1.9):

$$(3.16) \quad \begin{aligned} \text{grad } \langle \text{grad } \mathcal{L}_n[u], K_n[u] \rangle &= (\text{grad } \mathcal{L}_n[u])' K_n[u] + K_n'^* [\text{grad } \mathcal{L}_n[u]] = \\ &= (\text{grad } \mathcal{L}_n[u])' K_n[u] + K_n'^* [\text{grad } \mathcal{L}_n[u]] = L_K \text{grad } \mathcal{L}_n[u] = 0, \end{aligned}$$

giving rise to (3.15). Since on the submanifold $\bar{M}_{(N)}$ the gradient $\text{grad } \mathcal{L}_n[u] = 0$ for all $n \in \mathbb{Z}$, we obtain from (3.15) that the local functional $h^{(t)}[u] \in \Lambda^0(\bar{M}_{(N)})$ does not depend on the index $n \in \mathbb{Z}$.

The properties of the manifold $\bar{M}_{(N)}$, described above, make it possible to consider it as a symplectic manifold endowed with the symplectic structure $\omega^{(2)} \in \Lambda^2(\bar{M}_{(N)})$, given by expressions (3.8) and (3.10). From this point of view we can proceed to studying the integrability properties of the dynamical system (1.1) reduced upon the invariant finite-dimensional manifold $\bar{M}_{(N)} \subset M$.

First, we observe that the vector field d/dt on $\bar{M}_{(N)}$ is canonically Hamiltonian with respect to the symplectic structure $\omega^{(2)} \in \Lambda^2(\bar{M}_{(N)})$:

$$(3.17) \quad -i_{\frac{d}{dt}} \omega^{(2)}(u, p) = dh^{(t)}(u, p),$$

where $h^{(t)}(u, p) := h^{(t)}(u)$ and $(u, p)^\tau \in \bar{M}_{(N)}$ are canonical variables induced on the manifold $\bar{M}_{(N)}$ by the Liouville 1-form (3.8). Really, from the expression (3.15) one obtains that

$$d i_{\frac{d}{dt}} \langle \text{grad } \mathcal{L}_n[u], du_n \rangle = -D_n dh^{(t)}[u],$$

which, being supplemented with the identity (3.9) in the form

$$i_{\frac{d}{dt}} d \langle \text{grad } \mathcal{L}_n[u], du_n \rangle = -D_n i_{\frac{d}{dt}} \omega^{(2)}[u],$$

give rise to the following:

$$(3.18) \quad \frac{d}{dt} \langle \text{grad } \mathcal{L}_n[u], du_n \rangle = -D_n (dh^{(t)}[u] + i_{\frac{d}{dt}} \omega^{(2)}[u]),$$

Since $\text{grad } \mathcal{L}_n[u] = 0$ on $\bar{M}_{(N)}$, from (3.18) one follows the result (3.17).

The same one can claim subject to any of Hamiltonian systems (2.15), commuting with (1.1) on the manifold M . Moreover, owing to the functional independence of invariants $\gamma_j \in \mathcal{D}(M_{(N)})$, $j = \overline{0, N+1}$, entering into the Lagrangian functional (3.12), we can construct the set of functionally independent functions $h_j \in \mathcal{D}(\bar{M}_{(N)})$, $j = \overline{0, N}$, as follows:

$$(3.19) \quad \langle \text{grad } \mathcal{L}_n[u], \vartheta \text{ grad } \gamma_j[u] \rangle = \frac{dh_j[u]}{dn},$$

It is easy check that these functions $h_j \in \mathcal{D}(\bar{M}_{(N)})$, $j = \overline{0, N}$, are commuting to each other and to the Hamiltonian function $h^{(t)} \in \mathcal{D}(\bar{M}_{(N)})$ with respect to the symplectic structure $\omega^{(2)} \in \Lambda^2(\bar{M}_{(N)})$. Thus, if the dimension $\dim \bar{M}_{(N)} = 2(N+1)$, the discrete dynamical system (1.1) reduced on the finite-dimensional sub-manifold $\bar{M}_{(N)} \subset M$ will be Liouville integrable.

4. EXAMPLE: THE DIFFERENTIAL-DIFFERENCE NONLINEAR SCHRÖDINGER DYNAMICAL SYSTEM AND ITS INTEGRABILITY

The entitled above dynamical system given by (1.2) on the discrete infinite-dimensional manifold $M \subset l_2(\mathbb{Z}; \mathbb{C}^2) \subset l_\infty(\mathbb{Z}; \mathbb{C}^2)$. Its Lax type integrability was stated in [5, 10, 13] making use of the simplest discretization of the standard Zakharov-Shabat spectral problem for the well-known nonlinear Schrödinger equation.

In this Section we will try to apply the gradient-holonomic integrability analysis devised above to the system (1.2). From the very beginning we will show the existence of an infinite hierarchy of functionally independent conservation laws, having solved the determining Lax equation (1.9) in the asymptotical form (2.10). The following lemma holds.

Lemma 4.1. *The expression*

$$(4.1) \quad \varphi_n := \left(\begin{array}{c} 1 \\ a_n(\lambda) \end{array} \right) \prod_{j=0}^n \sigma_j(\lambda) \exp[it(2 - \lambda - \lambda^{-1})],$$

where, by definition,

$$(4.2) \quad \begin{aligned} \sigma_j(\lambda) &\simeq \frac{\lambda}{h_j[\psi^*, \psi]} \left(1 - \sum_{s \in \mathbb{Z}_+} \sigma_j^{(s)}[\psi^*, \psi] \lambda^{-s-1} \right), \\ a_n(\lambda) &\simeq \sum_{s \in \mathbb{Z}_+} a_n^{(s)}[\psi^*, \psi] \lambda^{-s}, \end{aligned}$$

is an asymptotical solution as $|\lambda| \rightarrow \infty$ to the Lax determining equation (1.9)

$$(4.3) \quad d\varphi_n/dt + K^{t,*} \varphi_n = 0$$

for all $n \in \mathbb{Z}$ with the operator $K^{t,*} : T^*(M) \rightarrow T^*(M)$ of the form:

$$(4.4) \quad K^{t,*} = \left(\begin{array}{cc} i\Delta^{-1}D_n^2 - i\psi_n^*(\psi_{n+1} + \psi_{n-1}) - & i\psi_n^*(\psi_{n+1}^* + \psi_{n-1}^*) \\ -i(\Delta + \Delta^{-1}) \cdot \psi_n^* \psi_n & \\ -i\psi_n(\psi_{n+1} + \psi_{n-1}) & -i\Delta^{-1}D_n^2 + i\psi_n(\psi_{n+1}^* + \psi_{n-1}^*) + \\ & +i(\Delta + \Delta^{-1}) \cdot \psi_n^* \psi_n \end{array} \right).$$

Proof. To prove Lemma 4.1 it is enough to find the corresponding coefficients of the asymptotical expansions (4.2). To do this we will consider the following equations easily obtained from (1.9):

$$(4.5) \quad \begin{aligned} &D_n^{-1}[-\ln h_n + \ln(1 - \sum_{s \in \mathbb{Z}_+} \sigma_n^{(s)} \lambda^{-s-1})]_{t+} \\ &+ i\lambda[h_{n+1}^{-1}(1 - \psi_n^* \psi_n)(1 - \sum_{s \in \mathbb{Z}_+} \sigma_n^{(s)} \lambda^{-s-1}) - 1] + \\ &+ \frac{i}{\lambda} \left[\frac{(1 - \psi_{n-1}^* \psi_{n-1})h_n}{(1 - \sum_{s \in \mathbb{Z}_+} \sigma_n^{(s)} \lambda^{-s-1})} - 1 \right] - i\psi_n^*(\psi_{n+1} + \psi_{n-1}) + \\ &\quad + i\psi_n^*(\psi_{n+1}^* + \psi_{n-1}^*) \sum_{s \in \mathbb{Z}_+} a_n^{(s)} \lambda^{-s}; \end{aligned}$$

$$\begin{aligned}
 & \left(\sum_{s \in \mathbb{Z}_+} a_n^{(s)} \lambda^{-s} \right) D_n^{-1} \left[-\ln h_n + \ln \left(1 - \sum_{s \in \mathbb{Z}_+} \sigma_n^{(s)} \lambda^{-s-1} \right) \right]_t + 4i \left(\sum_{s \in \mathbb{Z}_+} a_n^{(s)} \lambda^{-s} \right) + \\
 & + \left[i \lambda h_{n+1} (\psi_{n+1}^* \psi_{n+1} - 1) \left(\sum_{s \in \mathbb{Z}_+} a_{n+1}^{(s)} \lambda^{-s} \right) \left(\sum_{s \in \mathbb{Z}_+} a_{n+1}^{(s)} \lambda^{-s} \right) - \sum_{s \in \mathbb{Z}_+} a_n^{(s)} \lambda^{-s} \right] + \\
 & + \frac{d}{dt} \sum_{s \in \mathbb{Z}_+} a_n^{(s)} \lambda^{-s} + \frac{i}{\lambda} \left[\frac{(\psi_n^* - \psi_{n-1} - 1) \left(\sum_{s \in \mathbb{Z}_+} a_{n+1}^{(s)} \lambda^{-s} \right) h_n}{\left(1 - \sum_{s \in \mathbb{Z}_+} \sigma_n^{(s)} \lambda^{-s-1} \right)} - \sum_{s \in \mathbb{Z}_+} a_n^{(s)} \lambda^{-s} \right] + \\
 & - i \psi_n (\psi_{n+1} + \psi_{n-1}) + i \psi_n (\psi_{n+1}^* + \psi_{n-1}^*) \sum_{s \in \mathbb{Z}_+} a_n^{(s)} \lambda^{-s}.
 \end{aligned}$$

Having equated the coefficients of (4.5) at the same degrees of the parameter $\lambda \in \mathbb{C}$, we obtain step-by-step the functional expression expression for $h_n, \sigma_n^{(s)}$ and $a_n^{(s)}, n \in \mathbb{Z}, s \in \mathbb{Z}_+$:

$$\begin{aligned}
 (4.6) \quad h_n &= (1 - \psi_n^* \psi_n), a_n^{(0)} = 0, a_n^{(1)} = \beta, \\
 \sigma_n^{(0)} &= \psi_{n-1}^* (\psi_n + \psi_{n-2}) - i \Delta^{-1} D_n^2 (\ln h_n)_t, \\
 \sigma_n^{(1)} &= i \frac{d}{dt} \sigma_{n-1}^{(0)} + (h_{n-1} h_{n-2} - 1) + a_{n-1}^{(1)} \psi_{n-1}^* (\psi_n + \psi_{n-2}), \\
 a_n^{(2)} &= -3a_{n-1}^{(1)} + i \frac{d}{dt} \sigma_{n-1}^{(1)} - i a_{n-1}^{(1)} D_n^{-1} (\ln h_{n-1})_t + \\
 & + a_n^{(1)} \sigma_n^{(0)} - \psi_{n-1} (\psi_n^* + \psi_{n-2}^*) a_{n-1}^{(1)}, \\
 \frac{d}{dt} h_n &= i D_n (\psi_{n-1}^* \psi_n - \psi_n^* \psi_{n-1}), \dots,
 \end{aligned}$$

whence

$$\begin{aligned}
 (4.7) \quad \sigma_n^{(0)} &= -(\psi_n^* \psi_{n-1} + \psi_{n-1}^* \psi_{n-2}), \\
 \sigma_n^{(1)} &= i \frac{d}{dt} \sigma_{n-1}^{(0)} + (1 - \psi_{n-1}^* \psi_{n-1}) (1 - \psi_{n-2}^* \psi_{n-2}) + \beta \psi_{n-1}^* (\psi_n + \psi_{n-2}), \dots,
 \end{aligned}$$

and so on. Thus, having stated the corresponding iterative equations solvable for all $s \in \mathbb{Z}_+$, we can claim that expression (4.1) is a true asymptotical solution to the Lax equation (4.3). \square

Recalling now that the expression

$$(4.8) \quad \gamma(\lambda) := - \sum_{n=0}^N \ln h_n + \sum_{n=0}^N \ln \left(1 - \sum_{s \in \mathbb{Z}_+} \sigma_n^{(s)} \lambda^{-s-1} \right)$$

as $|\lambda| \rightarrow \infty$ is a generating function of conservation laws for the dynamical system (1.2), one finds that functionals

$$\begin{aligned}
 (4.9) \quad \bar{\gamma}_0 &= \prod_{n=0}^N (1 - \psi_n^* \psi_n), \gamma_0 = - \sum_{n=0}^N \sigma_n^{(0)}, \\
 \gamma_1 &= - \sum_{n=0}^N (\sigma_n^{(1)} + \frac{1}{2} \sigma_n^{(0)} \sigma_n^{(0)}), \\
 \gamma_2 &= - \sum_{n=0}^N (\sigma_n^{(2)} + \frac{1}{3} \sigma_n^{(0)} \sigma_n^{(0)} \sigma_n^{(0)} + \sigma_n^{(0)} \sigma_n^{(1)}), \dots,
 \end{aligned}$$

and so on, make up an infinite hierarchy of exact conservative quantities for the discrete nonlinear Schrödinger dynamical system (1.2).

Make here some remarks concerning the complete integrability of the discrete nonlinear Schrödinger dynamical system (1.2). First we can easily enough state, making use of the standard asymptotical small parameter approach [6, 12, 7], that the Nöther equation (1.5) possesses [11, 10] the following exact implectic operator solution:

$$(4.10) \quad \vartheta = \begin{pmatrix} 0 & i h_n \\ -i h_n & 0 \end{pmatrix},$$

$n \in \mathbb{Z}$, subject to which the dynamical system (1.2) is Hamiltonian:

$$(4.11) \quad \frac{d}{dt} \begin{pmatrix} \psi_n \\ \psi_n^* \end{pmatrix} = -\vartheta \operatorname{grad} H_\vartheta[\psi^*, \psi]_n$$

on the periodic manifold $M_{(N)}$, where the Hamiltonian function

$$(4.12) \quad H_\vartheta := \sum_{n=0}^N \ln h_n^2 - \sum_{n=0}^N (\psi_n^* \psi_{n+1} - \psi_n^* \psi_{n+1}) = 2 \ln \bar{\gamma}_0 - \frac{1}{2}(\gamma_0 + \gamma_0^*).$$

By means of similar, but more cumbersome calculations, one can find the second implectic operator solution to the Nöther equation (1.5) in the following matrix form:

$$(4.13) \quad \eta = \begin{pmatrix} (h_n - \psi_n D_n^{-1} \psi_n) \Delta & (\psi_n^2 + \psi_n D_n^{-1} \psi_n) \Delta^{-1} \\ \psi_n^* D_n^{-1} \psi_n^* \Delta & -(1 + \psi_n^* D_n^{-1} \psi_n) \Delta^{-1} \end{pmatrix} \times \\ \times \begin{pmatrix} \psi_n D_n^{-1} \psi_n & (h_n - \psi_n D_n^{-1} \psi_n^*) \\ 1 + \psi_n^* D_n^{-1} \psi_n & -(\psi_n^* + \psi_n^* D_n^{-1} \psi_n^*) \end{pmatrix},$$

where the operation $D_n^{-1}(\dots) := \frac{1}{2}[\sum_{k=0}^{n-1} (\dots)_k - \sum_{k=n}^N (\dots)_k]$ is quasi-skew-symmetric with respect to the usual bi-linear form on $T^*(M_{(N)}) \times T(M_{(N)})$, satisfying the operator identity $(D_n^{-1})^* = -\Delta^{-1} D_n^{-1} \Delta$, $n \in \mathbb{Z}$.

The implectic operators (4.10) and (4.13) are compatible, that makes it possible to construct by means of the algebraic gradient-holonomic algorithm the related Lax type representation for the dynamical system (1.2). The corresponding result is as follows: the discrete linear spectral problem

$$(4.14) \quad \Delta f_n = l_n[\psi^*, \psi; \lambda] f_n,$$

where $f \in l_\infty(\mathbb{Z}; \mathbb{C}^2)$ and for $n \in \mathbb{Z}$

$$(4.15) \quad l_n[\psi^*, \psi; \lambda] = \begin{pmatrix} \lambda & \psi_n \\ \psi_n^* & \lambda^{-1} \end{pmatrix},$$

allows the linear Lax type isospectral evolution

$$(4.16) \quad df_n/dt = p_n(l) f_n$$

for some matrix $p_n(l) \in \operatorname{End} \mathbb{C}^2$, $n \in \mathbb{Z}$, equivalent to the Hamiltonian flow

$$(4.17) \quad df_n/dt = \{H_\vartheta, f_n\}_\vartheta,$$

where $\{.,.\}_\vartheta$ is the corresponding to (4.10) Poissonian structure on the manifold $M_{(N)}$. The equivalence of (4.10) and (4.17) can be easily enough demonstrated, if to construct the corresponding to (4.14) monodromy matrix $S_n(\lambda)$, $n \in \mathbb{Z}$, for all $\lambda \in \mathbb{C}$ and to calculate the Hamiltonian evolution

$$(4.18) \quad \frac{d}{dt} S_n(\lambda) = \{H_\vartheta, S_n(\lambda)\}_\vartheta = [p_n(l), S_n(\lambda)],$$

giving rise to the same matrix $p_n(l) \in \operatorname{End} \mathbb{C}^2$, $n \in \mathbb{Z}$, as that entering equation (4.16).

Thus, we have shown that the nonlinear discrete Schrödinger dynamical system (1.2) is Lax type integrable bi-Hamiltonian flow on the manifold $M_{(N)}$. Since the solution $\varphi(\lambda) \in T^*(M_{(N)})$, constructed above, satisfies the gradient-like relationship

$$(4.19) \quad \lambda \vartheta \varphi(\lambda) = \eta \varphi(\lambda)$$

for all for $\lambda \in \mathbb{C}$, we derive that the found above conservation laws are commuting to each other with respect to both Poisson brackets $\{.,.\}_\vartheta$ and $\{.,.\}_\eta$. The latter gives rise to the classical Liouville integrability [2, 14] of the discrete nonlinear Schrödinger dynamical system (1.2) on the periodic manifold $M_{(N)}$. The detail analysis of the integrability procedure via the mentioned before Bogoyavlensky- Novikov reduction [15, 4] and the explicit construction of solutions to the dynamical system (1.2) are planned to be presented in a separate work.

5. CONCLUSION

The gradient-holonomic scheme of direct studying Lax type integrability of differential-difference nonlinear dynamical systems devised in this work appears to be effective enough for applications in the one-dimensional case, similar to the case [6, 14, 12, 7] of nonlinear dynamical systems defined on spatially one-dimensional functional manifolds. The developed algorithm, developing that before suggested in [11, 10], makes it possible to construct simply enough an infinite hierarchy of conservation laws as well as to calculate their compatible co-symplectic structures. As it was also shown, the reduced via the Bogoyavlensky-Novikov approach integrable Hamiltonian dynamical systems on the corresponding invariant periodic submanifolds generate finite dimensional Liouville integrable Hamiltonian systems with respect to the canonical Gelfand-Dikiy type symplectic structures. As an example almost complete integrability analysis of the nonlinear discrete Schrödinger dynamical system was presented in details.

Subject to different not-direct approaches to studying the integrability of differential-difference dynamical systems on discrete manifolds it is worth to mention the works [18, 19, 20, 21, 22, 27] based on the inverse spectral transform and related Lie-algebraic methods, where there are constructed *a priori* Lax type integrable Hamiltonian flows and possessing infinite hierarchies of conservation laws. Concerning these approaches, many of their important analytical properties were constructively absorbed by the gradient-holonomic scheme of this work and realized directly as an algorithm.

We would also like here to mention the interesting differential-algebraic approaches [23, 28, 29] proposed for direct studying the integrability both of differential and differential-difference dynamical systems. So, in [23, 25, 24] there was analyzed by means of simple enough differential-algebraic tools a generalized (owing to D. Holm and M. Pavlov) Riemann type hydrodynamical hierarchy of dynamical systems

$$(5.1) \quad D_t^N u = 0, \quad D_t := \partial/\partial t + uD_x, D_x := \partial/\partial x,$$

on a smooth functional manifold $M \subset C^\infty(\mathbb{R}; \mathbb{R})$ for any integer $N \in \mathbb{Z}_+$ and proved their both bi-Hamiltonian structure and Lax type integrability. Having replaced in equations (5.1) the spatial differentiation $D_x, x \in \mathbb{R}$, by its discrete analog $D_n := \Delta - 1, n \in \mathbb{Z}$, one can similarly construct a generalized Riemann type hierarchy of the following discrete dynamical systems

$$(5.2) \quad \mathfrak{D}_t^N u_n = 0, \quad \mathfrak{D}_t := \partial/\partial t + u_n D_n, \quad n \in \mathbb{Z},$$

for any integer $N \in \mathbb{Z}_+$ on a suitable discrete manifold $M \subset l_2(\mathbb{Z}; \mathbb{R})$, and whose integrability properties are important from practical point of view. In particular, it would be interesting both to apply the devised in this work the direct gradient-holonomic integrability approach to hierarchy (5.2) and to find its correct differential-difference analog using the already known [23, 24, 26] corresponding Lax type representations. As one can easily check, a one of the discrete analogs of the corresponding linear Lax type "spectral" problem for (5.1) at $N = 2$ has the form

$$(5.3) \quad \Delta f_n = l_n[u, z; \lambda] f_n, \quad l_n[u, z; \lambda] = (\mathbf{1} - \mathcal{A}_{n+1})^{-1}(\mathbf{1} - \mathcal{A}_n),$$

where, by definition,

$$(5.4) \quad \mathcal{A}_n := \begin{pmatrix} -\lambda u_n & -z_n \\ 2\lambda^2 n & \lambda u_n \end{pmatrix}$$

and $z_n := \mathfrak{D}_n u_n$ for any $n \in \mathbb{Z}$. Remark here that the linear "spectral" problem (5.3) is non-uniform, exactly depending on the variable index $n \in \mathbb{Z}$.

6. ACKNOWLEDGEMENTS

The second author (N.B.) is very indebted to the Abdus Salam International Centre for Theoretical Physics, Trieste, Italy, for the kind hospitality during his ICTP-2010 research scholarship. A.P. is cordially indebted to Profs. Denis Blackmore (NJIT, Newark, USA) and Maksim Pavlov (Lebedev Physics Institute of RAS, Moscow, Russia) for valuable comments and remarks concerning the gradient-holonomic scheme of integrability analysis and its application to the generalized (owing to D. Holm and M. Pavlov) Riemann type hydrodynamical hierarchy of dynamical systems. Authors are also cordially appreciated to Mrs. Natalia Prykarpatska (Lviv, Ukraine) and

Dilys Grilli (Trieste, Publications Office, ICTP) for professional help in editing and preparing the manuscript for publication.

REFERENCES

- [1] Abraham R., Marsden J. Foundation of mechanics. *The Benjamin/Cummings Publ. Co*, Masachusets, 1978
- [2] Arnold V.I. Mathematical methods of classical mechanics. NY, *Springer*, 1978
- [3] Lax P.D. Periodic solutions of the KdV equation // *Commun. Pure and Appl. Math.*, 1975, 28, N421, p. 141-188
- [4] Novikov S.P. (Editor) Theory of Solitons. Moscow, "Nauka", 1980 (in Russian)
- [5] Ablowitz M., Ladik J. Nonlinear differenmtial-difference equations. *J. Math. Phys.*,1975, 16, N3, p.598-603
- [6] Prykarpatsky A. and Mykytyuk I. Algebraic integrability of nonlinear dynamical systems on manifolds: classical and quantum aspects. *Kluwer Academic Publishers*, the Netherlands, 1998
- [7] Hentosh O., Prytula M. and Prykarpatsky A. Differential-geometric and Lie-algebraic foundations of investigating nonlinear dynamical systems on functional manifolds, 2nd edn. Lviv: Lviv University Publication, 2006 (in Ukrainian)
- [8] Blaszak M. Multi-Hamiltonian theory of dynamical systems. Berlin-Heidelberg: Springer, 1998
- [9] Magri F. A geometric approach to the nonlinear solvable equations. *Lect. Notes Phys.*, 1980, N120, p. 233-263
- [10] Mitropolsky Yu.A. and Samoilenko V.H. Differential-difference dynamical systems, associated with the difference Dirac operator, and their complete integrability. *Ukr. Math. J.*, 1985, 37, N2, p. 180-186 (in Russian)
- [11] Prykarpatsky A.K. Elements of the integrability theory of discrete dynamical systems. *Ukr. Math. J.*, 39, N1, p. 298-301 (in Russian)
- [12] Blackmore D., Prykarpatsky A.K. and Samoilenko V. Nonlinear dynamical systems of mathematical physics: spectral and differential-geometric integrability analysis. NY, World Scientific, 2010
- [13] Bogolubov N.N. (Jr.) and Prykarpatsky A.K. Inverse periodic problem for a discrete approximation of the Nonlinear Schrödinger equation. *Dokl. AN SSSR*, 1982, 265, N5, p. 1103-1108
- [14] Mitropolsky Yu.A., Bogolubov N.N. (Jr.), Prykarpatsky A.K. and Samoilenko V.H. Integrable dynamical systems: differential-geometric and spectral aspects. K.: "Naukova Dumka", 1987 (in Russian)
- [15] Bogoyavlensky O.I. and Novikov S.P. On the connection of Hamiltonian formulations of stationary and non-stationary problems. *Funk. Anal. Appl.*, 1976, 10, N1, p. 9-13 (in Russian)
- [16] Gelfand I.M. and Dikii L.A. Integrable nonlinear equations and the Liouville theorem. *Funk. Anal. Appl.*, 1979, 13, N1, p. 8-20 (in Russian)
- [17] Fuchssteiner B. and Fokas A. Symplectic structures, Backlund transformations and hereditary symmetries. *Physica D*, 1981, 4, N1, p. 47-66
- [18] Kupershmidt B.A. Discrete Lax equations and differential-difference calculus. // *Asterisque*, 1985, vol. 123, p. 5-212.
- [19] Deift P. Li L.-C., Tomei C. Loop groups, discrete versions of some classical integrable systems and rank-2 extensions. *Memoirs of the AMS*, 1992, vol. 100, No 479, p. 1-101
- [20] Moser J., Veselov A. P. Discrete versions of some classical integrable systems and factorization of matrix polynomials. Zurich, ETH, Preprint, 1989, 76p.
- [21] Baez J. C., Gillam J. W. An algebraic approach to discrete mechanics. *Lett. Math. Phys.*, 1994, vol. 31, No 3, p. 205-212.
- [22] Blaszak M., Szum A. and Prykarpatsky A. Central extension approach to integrable field and lattice-field systems in (2+1)-dimensions. *Reports on Mathematical Physics*, 1999.-37, N5, p. 27-32
- [23] Prykarpatsky A.K., Artemovych O.D., Popowicz Z. and Pavlov M.V. Differential-algebraic integrability analysis of the generalized Riemann type and Korteweg-de Vries hydrodynamical equations. *J. Phys. A: Math. Theor.* 43 (2010) 295205 (13pp)
- [24] Golenia J., Pavlov M., Popowicz Z. and Prykarpatsky A. On a nonlocal Ostrovsky-Whitham type dynamical system, its Riemann type inhomogeneous regularizations and their integrability. *SIGMA*, 2010, 6, p. 1-13
- [25] Golenia J., Bogolubov N. (jr.), Popowicz Z., Pavlov M. and Prykarpatsky A. A new Riemann type hydrodynamical hierarchy and its integrability analysis. *Preprint ICTP*, IC/2009/095, 2009
- [26] Pavlov M. The Gurevich-Zybin system. *J. Phys. A: Math. Gen.* 2005, 38, p. 3823-3840
- [27] Hentosh O., Prykarpatsky A. and Samoilenko V.H. The Lie-Algebraic structure of Lax type integrable nonlocal differential-difference equations. *Journal of Nonlinear Oscillations*. 2000, v.3, N1, p.84-94
- [28] Hardouin C. and Singer M.F. Differential Galois theory of linear difference equations. *Math. Ann.*, 2008, 342, p. 333-377
- [29] Wilson G. On the quasi-Hamiltonian formalism of the KdV equation. *Phys. Lett.* 132, p. 445-50

THE IVAN FRANKO STATE PEDAGOGICAL UNIVERSITY, DROHOBYCH, LVIV REGION, UKRAINE, AND THE DEPARTMENT OF APPLIED MATHEMATICS AT THE AGRARIAN UNIVERSITY OF KRAKOW, POLAND
E-mail address: yarpry@gmail.com

THE V.A. STEKLOV MATHEMATICAL INSTITUTE OF RAS, MOSCOW, RUSSIAN FEDERATION AND THE ABDUS SALAM INTERNATIONAL CENTRE OF THEORETICAL PHYSICS, TRIESTE, ITALY
E-mail address: nikolai.bogolubov@hotmail.com

THE DEPARTMENT OF MINING GEODESY AT THE AGH UNIVERSITY OF SCIENCE AND TECHNOLOGY, CRACÓW 30059, POLAND, AND THE IVAN FRANKO STATE PEDAGOGICAL UNIVERSITY, DROHOBYCH, LVIV REGION, UKRAINE
E-mail address: pryk.anat@ua.fm, prykanat@agh.edu.pl

THE DEPARTMENT OF MECHANICS AND MATHEMATICS AT THE T. SHEVCHENKO NATIONAL UNIVERSITY, KYIV, UKRAINE
E-mail address: vsam@univ.kiev.ua