

<http://www.geojournals.cn/dzxb/ch/index.aspx>

广东河源白石冈岩体：一个高分异的 I 型花岗岩

邱检生, 胡建, 王孝磊, 蒋少涌, 王汝成, 徐夕生

南京大学地球科学系, 成矿作用国家重点实验室, 210093

内容提要:广东河源白石冈岩体位于近东西向展布的佛冈花岗岩带的东端, 主体岩性为中粗粒黑云母花岗岩, 主要组成矿物为石英(25%~35%)、微纹长石(45%~50%)、斜长石($An=20\sim30$, 15%~20%)和黑云母(5%~10%)。锆石 U-Pb 定年结果表明其形成年龄为 148.5 ± 1.6 Ma, 属晚侏罗世岩浆活动的产物。化学成分上, 该岩体铝弱过饱和, A/NKC 值主要变化于 1.0~1.1 之间; 富硅, 富钾($K_2O/Na_2O = 1.31\sim 1.70$), 全碱含量中等偏低($K_2O+Na_2O = 7.44\%\sim 8.48\%$), 碱铝指数(AKI 值)为 0.75~0.88, 可归为高钾钙碱性岩系。微量和稀土元素组成上, 岩体富 Rb、Th、U、Pb 和轻稀土, 贫 Ba、Sr、P、Ti, Rb/Sr 比值高, K/Rb 比值低, 铕负异常显著($\delta Eu = 0.05\sim 0.28$), Nb、Ta、Zr、Hf 等高场强元素含量及 $10^4\times Ga/Al$ 比值(2.43~3.26)较之典型 A 型花岗岩均偏低。岩体的 $\epsilon_{Nd}(t)$ 值为 $-5.99\sim -7.51$, T_{DM} 值偏低(1.42~1.54 Ga), 综合地球化学资料指示其应属高分异的 I 型花岗岩。结合对区域动力地质背景的全面分析, 表明白石冈岩体形成于后造山的伸展引张环境, 是由底侵的幔源基性岩浆及其诱发的长英质岩浆在深部岩浆房混合, 并经高程度分离结晶的产物。

关键词: 高分异 I 型花岗岩; 壳幔相互作用; 地球化学; 岩石成因; 广东白石冈岩体

南岭花岗岩类分布十分广泛, 约占全区总面积的 1/5(袁忠信等, 1992)。花岗岩的成因及其产出构造背景是该区地质研究的核心问题, 长期以来倍受关注。近年来的研究表明, 南岭自早中生代开始即处于岩石圈伸展—减薄的引张构造背景(赵振华等, 1998; Chen et al., 2002; Li et al., 2003; Wang et al., 2003; 范蔚茗等, 2003), 一些具低 T_{DM} 值和高 $\epsilon_{Nd}(t)$ 值花岗岩带的发现指示它们的形成受到了岩石圈上地幔过程的制约, 是壳幔相互作用的结果(Gilder et al., 1996; Chen et al., 1998; Hong et al., 1998; Sewell et al., 2000; Shen et al., 2000), 这些新认识极大地推动了南岭花岗岩研究的进展。白石冈岩体位于广义的规模巨大的佛冈花岗岩带的东端, 处于近东西向的佛冈—丰良深断裂与北东向邵武—河源深断裂的交汇部位(图 1), 是区内壳幔相互作用形成的花岗质岩石的典型代表。本文拟报道该岩体的锆石 U-Pb 年龄, 并结合元素—同位素地球化学资料, 探讨其成因和形成的构造背景, 这一研究对于进一步揭示南岭地区的构造演化和壳幔相互作用过程均具有重要意义。

1 岩体地质与岩相学特征

白石冈岩体侵入的最新地层为下侏罗统蓝塘群砂页岩(图 1), 与围岩呈突变接触, 接触面倾向围岩, 出露面积约 440 km^2 。结合与区域范围内相关岩石的对比, 广东省地质矿产局(1988)将其归为燕山第三期($155\pm 5\sim 137\pm 5$ Ma)岩浆活动的产物。岩体岩性较均一, 相带分异不明显, 主体岩性为中粗粒黑云母花岗岩, 仅在局部与围岩接触处粒度有变细的现象, 同时围岩也不同程度的热变质, 岩体中缺乏或很少含岩石包体。

组成岩石的矿物主要为石英(25%~35%)、微纹长石(45%~50%)、斜长石($An=20\sim 30$, 15%~20%)和黑云母(5%~10%), 副矿物有锆石、磷灰石、萤石和钛铁氧化物等。微纹长石多不同程度泥化, 条纹主要呈细脉状和树枝状, 部分薄片微纹长石具发育的格子双晶, 偶尔可见石英与微纹长石呈文象交生, 指示岩体定位深度较浅。岩体中有时还出现少量白云母, 根据显微镜下白云母粒度较细, 形态多为半自形—他形, 主要呈细小鳞片状分布在长石表面, 或与黑云母呈反应关系等特征, 表明它们应属

注: 本文为教育部新世纪优秀人才支持计划和国家自然科学基金项目(编号 40132010, 40221301)联合资助成果。

收稿日期: 2004-11-10; 改回日期: 2005-04-08; 责任编辑: 刘淑春。

作者简介: 邱检生, 男, 1965 年生, 博士, 教授。主要从事花岗岩、火山岩及壳幔演化与成岩成矿关系研究。通讯地址: 210093, 南京大学地球科学系; Email: jsqiuwyh@public1.ppt.js.cn。




图1 白石冈岩体地质略图

Fig. 1 Sketch geological map of the Baishigang pluton

图 a 据赵子杰等(1987)修改;图 b 据 1:200000 紫金幅区调报告①;1—第四系;2—第三系;3—下侏罗统蓝塘群;4—泥盆系;5—下古生界;6—加里东期花岗岩;7—印支期花岗岩;8—燕山第二期花岗岩;9—燕山第三期花岗岩;10—燕山第四期花岗岩;11—采样点;12—实测及推测断裂:①—广州—从化断裂,②—邵武—河源断裂,③—紫金—博罗断裂,④—莲花山断裂,⑤—丽水—海丰断裂,⑥—佛冈—丰良断裂

Figure (a) is modified after Zhao Zijie et al. (1987) and Figure (b) is after regional survey report of Zijin area (1:200000)①;1—Quaternary System; 2—Tertiary System; 3—Lantang Group of Lower Jurassic; 4—Devolian System; 5—Lower Palaeozoic Group; 6—Caledonian granite; 7—Indosinian granite; 8—the second stage granite of Yanshanian; 9—the third stage granite of Yanshanian; 10—the fourth stage granite of Yanshanian; 11—sample localities; 12—measured or inferred faults; ①—Guangzhou-Conghua fault, ②—Shaowu-Heyuan fault, ③—Zijin-Boluo fault, ④—Lianhuashan fault, ⑤—Lishui-Haifeng fault, ⑥—Fogang-Fengliang fault

次生白云母(孙涛等,2002)。

2 分析方法

锆石 U-Pb 同位素定年在国土资源部天津地质矿产研究所用 VG-354 质谱仪测定,详细的分析过程见陆松年等(1991),全流程本底 Pb 为 $5 \times 10^{-11} \text{g}$, U 为 $2 \times 10^{-12} \text{g}$ 。主量元素在南京大学现代分析中心用 XRF 方法测定,相对标准样品的偏差,高含量氧化物小于 2%,低含量氧化物小于 10%。微量元素(包括稀土元素)在南京大学成矿作用国家重点实验室用 ICP-MS 方法分析,精度优于 10%。Nd 同位素组成在南京大学现代分析中心用 VG354 质谱测定,详细的分析方法见王银喜等(1988),在本文样品分析过程中,该仪器测定的 La Jolla 标样的 $^{143}\text{Nd}/^{144}\text{Nd} = 0.511854 \pm 6(2\sigma)$,Nd 同位素比值采用 $^{146}\text{Nd}/^{144}\text{Nd} = 0.7219$ 进行质量分馏校正,Sm、Nd 实验室全流程本底为 $5 \times 10^{-11} \sim 7 \times 10^{-11} \text{g}$ 。

3 年代学

前人对白石冈岩体进行过 K-Ar 年龄测定,所报道的 2 个年龄值分别为 139 Ma 和 142 Ma(广东省地质矿产局,1988)。由于 K-Ar 体系的抗扰动性差,加之封闭温度低(约 $200 \pm 50^\circ\text{C}$,郑永飞等,1997),使得定年结果常低于岩体的实际年龄。而锆石 U-Pb 计时体系的封闭温度($700 \pm 50^\circ\text{C}$,Dodson et al., 1985)与花岗质岩石的结晶温度相近,被认为是花岗质岩石定年最为可靠的方法。由表 1 所列的白石冈岩体锆石 U-Pb 定年结果可以看出,除 4 号锆石点的 $^{207}\text{Pb}/^{206}\text{Pb}$ 表面年龄偏高外,其余的表面年龄在测试误差范围内基本一致。在 $^{207}\text{Pb}/^{235}\text{U}-^{206}\text{Pb}/^{238}\text{U}$ 图解中,4 颗锆石点均投影在 U-Pb 谐和线上(图 2),说明样品中放射成因铅基本没有扩散丢失。由于半衰期差异,锆石中放射成因 ^{207}Pb 的丰度比放射成因 ^{206}Pb 的丰度约低一个数量级,因而对年轻锆石来说, $^{206}\text{Pb}/^{238}\text{U}$ 年龄值精度较高

表 1 白石岗岩体锆石 U-Pb 同位素定年结果
Table 1 U-Pb dating results for zircons from the Baishigang pluton

点号	样品情况		浓度(μg/g)		样品普通 铅含量 (ng)	同位素原子比率					表面年龄(Ma)		
	锆石特征	重量 (μg)	U	Pb		²⁰⁶ Pb/ ²⁰⁴ Pb	²⁰⁸ Pb/ ²⁰⁶ Pb	²⁰⁶ Pb/ ²³⁸ U	²⁰⁷ Pb/ ²³⁵ U	²⁰⁷ Pb/ ²⁰⁶ Pb	²⁰⁶ Pb/ ²³⁸ U	²⁰⁷ Pb/ ²³⁵ U	²⁰⁷ Pb/ ²⁰⁶ Pb
1	浅紫红色透明长柱状自形晶	30	2419	58	0.097	1139	0.07566	0.02355 (21)	0.1593 (26)	0.04906 (64)	150.1	150.1	150.5
2	浅紫红色透明长柱状自形晶	40	3144	73	0.150	1283	0.06110	0.02325 (19)	0.1573 (17)	0.04909 (31)	148.1	148.4	151.9
3	紫红色透明短柱状自形晶	40	6397	146	0.200	1925	0.05527	0.02320 (6)	0.1570 (8)	0.04907 (20)	147.9	148.1	151.0
4	浅紫红色透明细长柱自形晶	30	1300	32	0.054	1077	0.1223	0.02327 (40)	0.1581 (45)	0.04928 (105)	148.3	149.1	161.3

注:1~4 号点 ²⁰⁶Pb/²³⁸U 表面年龄加权平均值:148.5±1.6 Ma;括号内的数字为 2σ 绝对误差,例如:0.02355(21)表示 0.02355±0.00021(2σ)。




图 2 白石岗岩体锆石 U-Pb 年龄谐和图
Fig. 2 U-Pb concordia diagram for zircons of the Baishigang pluton

(Compston et al., 1992; Shen et al., 2000)。由 4 个锆石颗粒计算出白石岗岩体的 ²⁰⁶Pb/²³⁸U 表面年龄统计权重平均值为 148.5±1.6 Ma, 这一年龄代表了该岩体的侵位年龄,因此,白石岗岩体应属晚侏罗世岩浆活动的产物。

4 地球化学

4.1 主量元素

表 2 列出白石岗岩体代表性岩石样品的化学成份、CIPW 标准矿物及主要岩石化学参数。其主量元素特征可归结为:① 富硅, SiO₂ = 73.37%~76.05%, D.I. = 87.9~94.1, 反映岩体经历了高程度分异演化作用。② 铝弱过饱和, Al₂O₃ = 12.00%~13.16%, A/NKC 值主要变化于 1.0~1.1 之间, 绝大多数样品的 CIPW 标准矿物中均出现刚玉分子, 但含量多在 1% 以下, 与典型的强过铝 S 型花岗岩(A/NKC > 1.1, CIPW 标准矿物中刚玉分子

表 2 白石岗岩体岩石化学成分(%), CIPW 标准矿物及主要岩石化学参数

Table 2 Petrochemical compositions (%), CIPW-normative minerals and predominant petrochemical parameters of the Baishigang pluton

样号	BSG -1	BSG -2	BSG -3	BSG -5	BSG -7	BSG -8	BSG -9	BSG -10	BSG -14
SiO ₂	73.37	76.05	74.37	73.66	74.61	76.05	74.3	74.81	75.34
TiO ₂	0.24	0.07	0.12	0.23	0.21	0.15	0.10	0.18	0.10
Al ₂ O ₃	13.16	12.47	13.02	13.06	12.41	12.00	13.11	12.81	12.68
TF ₂ O ₃	2.37	1.18	1.69	2.42	2.26	1.78	1.30	1.99	1.39
MnO	0.05	0.03	0.04	0.04	0.04	0.03	0.04	0.04	0.05
MgO	0.38	0.08	0.14	0.26	0.19	0.13	0.20	0.16	0.11
CaO	1.32	0.91	0.87	1.31	1.16	1.10	1.06	1.27	0.85
Na ₂ O	2.92	3.33	3.39	3.11	2.86	3.22	3.19	3.09	3.36
K ₂ O	4.71	5.02	5.03	4.77	4.86	4.22	5.29	4.60	4.86
P ₂ O ₅	0.07	0.01	0.03	0.05	0.03	0.02	0.02	0.03	0.01
SO ₃	0.02	0.04	0.02	0.04	0.03	0.04	0.05	/	0.04
烧失	0.69	0.90	0.65	0.92	0.44	0.61	0.49	0.66	0.43
总量	99.30	100.09	99.40	99.80	99.10	99.30	99.10	99.60	99.10
Q	34.51	35.79	33.86	34.64	37.21	39.20	33.46	36.74	35.70
Or	28.31	29.95	30.14	28.52	29.15	25.29	31.73	27.49	29.11
Ab	25.08	28.39	29.03	26.57	24.51	27.57	27.34	26.38	28.76
An	6.24	4.25	4.20	6.28	5.66	5.41	5.22	6.19	4.21
C	1.01	/	0.47	0.49	0.39	0.17	0.24	0.49	0.36
ALK	7.63	8.35	8.42	7.88	7.72	7.44	8.48	7.69	8.22
A.R.	2.35	2.98	2.91	2.53	2.46	2.93	2.64	2.56	2.97
K/N	1.61	1.51	1.48	1.53	1.70	1.31	1.66	1.49	1.45
AKI	0.75	0.88	0.85	0.79	0.80	0.82	0.84	0.79	0.85
A/NKC	1.07	0.99	1.03	1.03	1.03	1.01	1.02	1.04	1.03
D.I.	87.9	94.1	93.0	89.7	90.9	92.1	92.5	90.6	93.6

注:Q—石英; Or—钾长石; Ab—钠长石; An—钙长石; C—刚玉; ALK = K₂O+Na₂O; A.R.—碱度率; K/N = K₂O/Na₂O; AKI = (Na₂O+K₂O) / Al₂O₃(分子比); A/NKC = Al₂O₃/(Na₂O+K₂O+CaO)(分子比); D.I.—分异指数。

含量 > 1%, Chappell et al., 2001) 有一定差别, 也不同于华南陆壳改造系列花岗岩(刘昌实等, 1990)。在 ACF 三角图解中, 样品点均投影在 I 型与 S 型花岗岩交界区(图 3)。③ 碱含量中等偏低, K₂O+Na₂O

= 7.44%~8.48%, A. R. 值变化于 2.35~2.98, AKI 值介于 0.75~0.88 之间, 与区内时代相近的花山和佛冈主体花岗岩相似(花山: 0.83~0.84, 朱金初等, 1989; 佛冈: 0.72~0.88, 陈小明等, 2002), 但较之 A 型花岗岩的平均值(0.95, Whalen et al., 1987)明显偏低。按洪大卫等(1987)提出的碱性、偏碱性和钙碱性花岗岩 AKI 值分界线(≥ 1.0 、 $0.9 \sim 1.0$ 和 < 0.9), 则岩体属钙碱性花岗岩。④ 富钾, $K_2O = 4.22\% \sim 5.29\%$, $K_2O/Na_2O = 1.31 \sim 1.70$, 其总体特征与 Barbarin (1999)划分的富钾钙碱性花岗岩类(KCG)相似。⑤ 铁、镁、钙、钛含量低, 贫磷, P_2O_5 含量均在 0.10% 以下, 且 P_2O_5 与 SiO_2 含量之间有较显著的负消长演化关系(图 4), 这也明显不同于典型的 S 型花岗岩, 后者常具有较高的 P_2O_5 含量, 且随分异作用的进行 P_2O_5 有递增的演化趋势(Chappell, 1999)。其 P_2O_5 及 CIPW 标准矿物中刚玉分子含量均与澳大利亚 Lachlan 褶皱带的 I 型花岗岩(Chappell, 1999)相似, 结合岩石分异演化程度高的特点, 表明白石冈岩体应属分异的 I 型花岗岩。

4.2 稀土及微量元素

白石冈岩体代表性岩石样品的稀土及微量元素测定结果列于表 3。由表中数据可看出, 岩体的稀土总量偏高, $\Sigma REE = 114.19 \times 10^{-6} \sim 398.78 \times 10^{-6}$; 富轻稀土, $LREE/HREE = 3.12 \sim 14.77$, $(La/Yb)_N = 2.79 \sim 21.21$, 其中轻稀土的分馏较之重稀土明显, $(La/Sm)_N$ 及 $(Gd/Yb)_N$ 比值分别为 1.69~5.47 和 1.04~2.50; 铕亏损强烈, $\delta Eu = 0.05 \sim 0.28$ 。随




图 3 白石冈岩体 ACF 图
Fig. 3 ACF diagram of the Baishigang pluton




图 4 白石冈岩体 SiO_2 - P_2O_5 关系图
Fig. 4 SiO_2 vs. P_2O_5 diagram of the Baishigang pluton
图中 I 型和 S 型花岗岩 P_2O_5 随 SiO_2 递增的变异趋势据 Chappell (1999)
The variation trends of P_2O_5 with increasing SiO_2 for I- and S-type granites are after Chappell (1999)




图 5 白石冈岩体稀土元素球粒陨石标准化配分型式
Fig. 5 Chondrite-normalized REE distribution patterns of the Baishigang pluton
球粒陨石值据 Boynton(1984)
The chondrite values are after Boynton (1984)

着分异指数的增大, 轻、重稀土比值及 δEu 值具有明显降低的演变趋势, 指示成岩过程中存在富轻稀土矿物(如磷灰石、褐帘石、独居石等)和斜长石的分异结晶作用。岩石的稀土元素球粒陨石标准化配分型式呈明显的右倾斜形(图 5), 与佛冈主体黑云母花岗岩相似(陈小明等, 2002), 但明显不同于典型 S 型花岗岩常表现出的“海鸥型”稀土配分型式。

微量元素组成上, 白石冈岩体富 Rb、Th、U、Pb, 贫 Ba、Sr、P、Ti(图 6), Rb/Sr 比值高(2.36~

表 3 白石岗岩体稀土及微量元素含量($\times 10^{-6}$)Table 3 REE and trace element abundances of the Baishigang pluton ($\times 10^{-6}$)

样号	BSG-1	BSG-2	BSG-3	BSG-5	BSG-7	BSG-8	BSG-9	BSG-10	BSG-14
La	63.47	16.82	45.84	93.09	75.13	58.93	23.54	45.59	39.81
Ce	125.10	37.80	101.55	183.52	155.84	124.12	47.80	93.89	90.84
Pr	13.24	5.02	12.22	19.66	17.62	14.60	5.71	11.00	11.40
Nd	47.65	20.47	45.73	65.60	59.93	52.49	21.03	39.78	43.69
Sm	9.13	6.25	11.63	10.70	10.38	10.91	5.51	8.25	11.31
Eu	0.62	0.11	0.28	0.91	0.59	0.44	0.38	0.51	0.23
Gd	7.35	7.14	11.85	9.17	9.10	10.89	6.18	8.52	10.71
Tb	0.90	1.12	1.68	1.03	1.08	1.45	1.01	1.11	1.48
Dy	5.32	7.86	11.25	6.31	6.83	10.25	7.36	7.65	10.33
Ho	1.03	1.62	2.33	1.26	1.53	2.16	1.54	1.56	2.13
Er	2.90	4.71	7.12	3.64	4.68	6.31	4.77	4.58	6.26
Tm	0.40	0.66	0.99	0.48	0.68	0.91	0.73	0.65	0.92
Yb	2.49	4.06	6.52	2.96	4.07	5.33	4.79	4.15	6.27
Lu	0.37	0.56	0.93	0.43	0.58	0.77	0.71	0.63	0.87
ΣREE	279.98	114.19	259.92	398.78	348.03	299.55	131.06	227.87	236.25
LREE/HREE	12.48	3.12	5.09	14.77	11.20	6.87	3.84	6.90	5.06
(La/Yb) _N	17.16	2.79	4.74	21.21	12.43	7.46	3.31	7.40	4.28
(La/Sm) _N	4.37	1.69	2.48	5.47	4.55	3.40	2.69	3.48	2.21
(Gd/Yb) _N	2.38	1.42	1.47	2.50	1.80	1.65	1.04	1.66	1.38
δEu	0.22	0.05	0.07	0.28	0.18	0.12	0.20	0.19	0.06
Cs	20.66	11.97	15.70	5.58	10.57	10.50	20.63	8.98	11.34
Rb	275.10	384.86	464.76	217.69	279.99	288.68	450.35	308.46	441.62
Sr	116.44	18.48	40.15	104.64	55.85	45.97	51.86	62.10	29.30
Ba	457.25	39.93	162.02	781.91	250.11	193.88	131.49	293.18	30.82
U	8.94	11.29	16.74	7.34	9.82	10.24	26.59	15.16	20.13
Th	45.23	34.25	46.38	35.63	49.30	57.21	43.52	38.81	59.24
Pb	35.69	32.38	43.08	29.27	30.75	98.87	43.70	39.22	40.57
Ga	19.13	17.44	22.45	18.33	18.35	18.88	16.89	19.28	18.95
Sc	5.46	2.99	4.54	5.27	3.25	4.18	2.55	4.05	3.66
Y	28.96	43.33	68.01	32.28	50.65	58.26	44.91	41.39	60.20
Nb	17.01	24.49	23.83	14.38	17.17	21.32	18.58	18.35	22.65
Ta	2.76	3.70	4.56	1.44	1.76	2.49	3.66	2.92	5.93
Zr	144.01	73.16	122.72	204.95	226.07	123.34	102.58	169.33	101.96
Hf	4.52	3.62	5.17	5.91	7.19	4.33	3.48	6.14	3.97
Rb/Sr	2.36	20.82	11.58	2.08	5.01	6.28	8.68	4.97	15.07
K/Rb	142.07	108.23	89.81	181.82	144.03	121.30	97.47	123.74	91.32
Nb/Ta	6.15	6.61	5.22	9.99	9.74	8.57	5.08	6.28	3.82
$10^4 \times \text{Ga}/\text{Al}$	2.75	2.64	3.26	2.65	2.79	2.97	2.43	2.84	2.82

20.82), K/Rb 比值低(绝大多数样品均小于 150, 表 3), 上述微量元素组成特征同样指示岩体经历了高程度的分异演化, 其 Ba、Sr、P、Ti 亏损应是成岩过程中斜长石、磷灰石和钛铁矿等的分离结晶所致。岩体的 Nb、Ta、Zr、Hf 等高场强元素含量及 Ga/Al 比值均较之典型 A 型花岗岩(Whalen et al., 1987) 偏低, 在 $(\text{Zr} + \text{Nb} + \text{Ce} + \text{Y}) \cdot 10^4 \times \text{Ga}/\text{Al}$ 关系图上, 样品点主要落在分异的 I 型花岗岩区(Eby, 1990, 图 7), 其 Nb/Ta 比值(3.82~9.99)与分异花岗岩的相应值(2.3~9.9, Dostal et al., 2000)十分接近, 这

一特征也说明白石岗岩体应属分异的 I 型花岗岩。

4.3 Nd 同位素

由于白石岗岩体具高的 Rb/Sr 比值, 使得年龄对 I_{Sr} 值的校正十分敏感, 在这种情况下, 岩石的 Sr 同位素组成往往不具有明确的成岩意义, 而 Nd 同位素由于其强的抗扰动性则能有效地示踪岩浆源区性质, 为此, 本文仅报道其 Nd 同位素组成的测定结果。

由表 4 所列数据可看出, 白石岗岩体的 Nd 同位素组成相对均一, $\epsilon_{\text{Nd}}(t) = -5.99 \sim -7.51$, 与




图6 白石冈岩体微量元素相对于原始地幔标准化蛛网图

Fig. 6 Primitive mantle-normalized spidergrams for trace elements of the Baishigang pluton
原始地幔值据 Sun & McDonough (1989)

The primitive mantle values are from Sun & McDonough (1989)




图7 白石冈岩体 $(Zr+Nb+Ce+Y)-10^4 \times Ga/Al$ 关系图

Fig. 7 $(Zr+Nb+Ce+Y)$ vs. $10^4 \times Ga/Al$ diagram of the Baishigang pluton

OGT—I, S 和 M 型花岗岩区; FG—分异的 I 型花岗岩区;
底图据 Eby (1990).

OGT—Field for I-, S- and M-type granitoids; FG—field for fractionated I-type granitoids; the base map is after Eby (1990)

区内时代相近的佛冈主体黑云母花岗岩 ($t=167$ Ma, $\epsilon_{Nd}(t) = -6.20 \sim -8.93$, 陈小明等, 2002; 包志伟等, 2003) 及龙窝花岗岩闪长岩 ($t=169.1$ Ma, $\epsilon_{Nd}(t) = -6.53 \sim -8.89$, 邱检生等, 2004a) 相似, 也可与千里山第一阶段斑状黑云母花岗岩 ($t=152$ Ma, $\epsilon_{Nd}(t) = -6.39 \sim -7.59$, 毛景文等, 1995) 相对比。在 $t-\epsilon_{Nd}(t)$ 关系图上, 样品点投影于华南元古宙地壳演化域上方 (图 8), 说明它们并非单纯起

源于基底变质岩的部分熔融, 成岩过程中应有幔源组份的参与。




图8 白石冈岩体 $\epsilon_{Nd}(t)-t$ 关系图

Fig. 8 $\epsilon_{Nd}(t)-t$ diagram of the Baishigang pluton

A—华南成熟度较低的古宙地壳; B—华南成熟度较高的元古宙地壳; DM—亏损地幔; CHUR—球粒陨石均一储库; 底图据沈渭洲等 (1993); 1—白石冈 (本文资料); 2—千里山 (毛景文等, 1995); 3—佛冈 (陈小明等, 2002; 包志伟等, 2003); 4—龙窝 (邱检生等, 2004a), 为了清晰地对比 Nd 同位素组成, 图中将各岩体的数据点作了适当分开, 因此, 样品点对应的横坐标并不代表各岩体的精确形成年龄

A—Immature Proterozoic crust in south China; B—mature Proterozoic crust in south China; DM—depleted mantle; CHUR—chondritic uniform reservoir; the base map is after Shen Weizhou et al. (1993); 1—Baishigang (this paper); 2—Qianlishan (Mao Jingwen et al., 1995); 3—Fogang (Chen Xiaoming et al., 2002; Bao Zhiwei et al., 2003); 4—Longwo (Qiu Jiansheng et al., 2004a); Data points of each plutons are properly separated for clarity and thus no accurate age differences are implied

白石冈岩体的二阶段 Nd 模式年龄变化于 1.42 ~ 1.54 Ga (表 4), 较之华夏地块基底变质岩的 Nd 模式年龄 (主要为 1.8 ~ 2.2 Ga, 陈江峰等, 1999) 显著偏低。近年来, 在华南相继识别出多条具低 T_{DM} 值的花岗岩带, 这些具低 T_{DM} 值的花岗岩理论上可通过二种方式形成, 其一为起源于年轻地壳物质的部分熔融, 其二是成岩过程中有不同比例地幔物质的加入。由于本区基底岩石的 T_{DM} 值主要为古、中元古代 (沈渭洲等, 1993; 陈江峰等, 1999), 不支持年轻地壳的存在, 因此, 目前普遍认为这些具低 T_{DM} 值的花岗岩是地幔物质参与成岩过程的重要表现 (Gilder et al., 1996; Chen et al., 1998; Hong et al., 1998; Shen et al., 2000)。白石冈岩体空间上大致位于陈江峰等 (1999) 划分的 S_3 低 T_{DM} 值花岗岩带上, 无疑指示地幔物质参与了成岩过程。

5 讨论

表 4 白石岗岩体 Sm-Nd 同位素组成

Table 4 Sm-Nd isotopic compositions of the Baishigang pluton

样号	Sm($\times 10^{-6}$)	Nd($\times 10^{-6}$)	Sm/Nd	$^{147}\text{Sm}/^{144}\text{Nd}$	$^{143}\text{Nd}/^{144}\text{Nd}(\pm 2\sigma)$	T_{Nd}	$\epsilon_{\text{Nd}}(t)$	$T_{2\text{DM}}(\text{Ga})$	$f(\%)$
BSG-2	6.62	21.05	0.314	0.1847	0.512264 ± 4	0.512085	-7.07	1.49	42.6
BSG-3	12.08	46.17	0.262	0.1539	0.512270 ± 18	0.512120	-6.37	1.45	46.3
BSG-5	10.24	64.87	0.158	0.09868	0.512232 ± 6	0.512136	-6.06	1.43	47.8
BSG-9	5.85	22.04	0.265	0.1585	0.512216 ± 7	0.512062	-7.51	1.54	40.2
BSG-10	9.29	40.11	0.232	0.1255	0.512246 ± 10	0.512124	-6.30	1.45	46.6
BSG-14	10.94	42.79	0.256	0.1566	0.512292 ± 6	0.512140	-5.99	1.42	48.2

注:为减少 $^{147}\text{Sm}/^{144}\text{Nd}$ 变化对Nd模式年龄计算产生的影响,表中所列 $T_{2\text{DM}}$ 年龄统一采用二阶段模式计算, f 为根据刘昌实等(1990)方法计算的岩浆源区中地幔物质的混入比例。

5.1 成岩过程

Nd 同位素示踪指示白石岗岩体成岩过程中有地幔组份的参与,即岩体属壳幔混源型花岗岩。壳幔物质的混合既可以发生在岩浆源区,也可以发生在幔源岩浆上侵途经地壳的过程中。由于岩体的Nd同位素组成相对均一, $\epsilon_{\text{Nd}}(t)$ 值与相应样品的 SiO_2 及Nd含量之间缺乏明显的线性演化关系,加之岩体岩性较均匀,各类岩石包体均不发育,说明成岩过程中壳幔物质的混合应发生在岩浆源区,因为深部岩浆房中的对流混合有利于岩浆均一化(Martin et al., 1994)。

选择华南陆壳和亏损地幔作为混合端元,利用刘昌实等(1990)和 Faure(1986)提供的二端元参数,将其Sr、Nd同位素组成校正到岩石形成时($t=148.5\text{ Ma}$)的初始值,则有华南陆壳: $\text{Nd}=28 \times 10^{-6}$, $\epsilon_{\text{Nd}}(t)=-13.28$; 亏损地幔: $\text{Nd}=14 \times 10^{-6}$, $\epsilon_{\text{Nd}}(t)=9.69$ 。再按简单的二元混合模拟计算,可以得出白石岗岩体成岩过程中地幔物质的卷入比例变化于40.2%~48.2%之间(表4)。如果这一比例地幔与地壳物质混合形成的岩浆直接结晶,显然难以形成目前所观测到的主量和微量元素地球化学特征。岩体富硅,Rb/Sr比值高,K/Rb比值低,Sr、Ba、P、Ti、Nb、Eu等元素显著亏损,这些特征充分说明岩浆经历了高分异的分离结晶作用。根据造岩矿物中上述元素分配系数的大小(Arth, 1976; Hanson, 1978; Green et al., 1986), Sr、Ba、Eu的亏损指示成岩过程中发生了斜长石和钾长石的分离结晶,而P和Ti、Nb的亏损则分别与磷灰石及含钛矿物(如钛铁矿、榍石等)的分离结晶有关(Raith, 1995; Wu et al., 2003)。岩石的轻稀土含量及轻、重稀土比值随分异演化程度增高而明显降低的演变趋势,应是成岩过程中富轻稀土矿物(如磷灰石、褐帘石、独居石等)的分离结晶所致。按Rayleigh分离结晶方程,经计算得到由矿物分离结晶作用形成的




图 9 白石岗岩体副矿物分离结晶作用过程判别图解
Fig. 9 Discrimination diagram showing the fractional crystallization process by separation of accessory minerals for the Baishigang pluton

Allan—褐帘石; Mon—独居石; Ap—磷灰石; Zr—锆石; 褐帘石和锆石分配系数数据 Mahood 等(1983); 独居石分配系数数据 Yurimoto 等(1990); 磷灰石的分配系数数据 Arth (1976)和 Fujimaki (1986); 图中黑点代表白石岗黑云母花岗岩样品的投影点,分异趋势线上的数字代表分离结晶作用的程度,计算过程中假定分异程度最低的样品为初始熔体

Allan—Allanite; Mon—monazite; Ap—apatite; Zr—zircon; partition coefficients are from Mahood et al. (1983) for allanite and zircon; Yurimoto et al. (1990) for monazite, Arth (1976) and Fujimaki (1986) for apatite. The solid circles in this figure are sample plot points of the Baishigang pluton, and the data shown on the differentiation tendency lines represent the fractionation degree of the accessory minerals. During the calculation, the least fractionated sample is supposed to be the parent melt

La-(La/Yb)_N 变异图解(图9),从白石岗岩体样品点在图中的分布来看,独居石和褐帘石的分离结晶是控制成岩过程中稀土元素变异的主要因素。

5.2 构造意义

对中国东南部中生代岩浆作用构造背景的认识是正确理解区内花岗岩成因的关键。自上世纪70年代以来,随着板块构造学说的兴起,许多学者认为华南中生代属于安第斯型活动大陆边缘,岩浆活动与

古太平洋板块向欧亚大陆的俯冲有关(郭令智等,1983;王鸿祯等,1983;任纪舜等,1990)。Zhou等(2000)提出自中侏罗世开始,古太平洋板向中国东南大陆俯冲消减,由此诱发玄武质岩浆的底侵,并促使中下地壳部分熔融,这一机制是产生区内各类花岗质岩浆的主要原因。近年来,越来越多的学者主张南岭地区自早中生代开始(175 Ma±)即处于软流圈上隆的陆内岩石圈伸展引张构造背景(Li et al., 2003; Wang et al., 2003; 范蔚茗等,2003)。可见对华南中生代岩浆作用动力地质背景的认识尚存在分歧。

南岭中生代花岗岩基本均呈EW向展布,大致可分为3条岩浆活动带,自北向南依次为诸广山—青障山带、大东山—贵东带及佛冈—新丰江带(Shu et al., 2004)。已有资料表明,自燕山早期开始,古太平洋板块沿NW向朝欧亚大陆俯冲(Maruyama et al., 1986),随后俯冲方向逐渐向NNW向过渡(Engelbreton et al., 1985),按此俯冲方向,则南岭内陆不太可能形成一系列呈东西向展布的岩浆活动带,说明南岭中生代岩浆活动并非完全受控于古太平洋构造域。近年来,在南岭地区相继厘定了一系列印支期花岗岩(Xu et al., 2003; 邱检生等, 2004b; 张文兰等,2004),表明该区中生代岩浆作用受到过特提斯构造域的制约,南岭地区深部表现出的EW向粤北幔坳—佛冈幔坡—广州幔隆和近EW向均匀单一完整的重力梯度带(舒良树等,2002)应是特提斯构造动力体系影响的印记。

白石冈岩体形成于燕山早期,化学成分上富钾,属高钾钙碱性岩石。Liegeois等(1998)认为高钾钙碱性岩石的岩浆源区通常与先期的俯冲作用有关,它们主要形成于同碰撞岩石圈加厚之后的伸展垮塌向非造山板内的过渡阶段。大量研究资料表明,华南印支期以挤压逆冲推覆和地壳叠置加厚为主要特征




图 10 白石冈岩体主量元素构造环境判别图解

Fig. 10 Tectonic discrimination diagram using major elements for the Baishigang pluton. IAG—岛弧花岗岩类; CAG—大陆弧花岗岩类; CCG—大陆碰撞花岗岩类; POG—后碰撞花岗岩类; RRG—与裂谷有关的花岗岩类; CEUG—与大陆造陆抬升有关花岗岩类; 底图据 Maniar 等(1989). IAG—Island arc granitoids; CAG—continental arc granitoids; CCG—continental collision granitoids; POG—post-collisional granitoids; RRG—rift-related granitoids; CEUG—continental epeirogenic uplift granitoids. The base map is after Maniar et al. (1989)

(Chen, 2001; Wang et al., 2002),主体变形作用的时间发生在 258~192 Ma(范蔚茗等,2003)。至燕山早期,则发育有 A 型花岗岩类(176~178 Ma)和双峰式火山岩(158~179 Ma)组合(Chen et al., 2002),并在燕山早期发现有具 OIB 特征的板内拉斑或碱性玄武岩(175~178 Ma, 赵振华等,1998; 陈培荣等,1999),上述岩石组合的出现是软流圈上涌和岩石圈伸展减薄的最直接证据,因此,南岭地区燕山早期的火成岩组合应形成于板内环境,主要受控于印支造山运动之后的后造山大陆裂解的地球动力学背景(Chen et al., 2002)。

采用相关的构造环境判别图投影也表明,白石冈岩体的投影点均落在后碰撞(post-collisional)或后碰撞伸展(post-collision extension)花岗岩区(图 10, 图 11)。尽管高场强元素(Nb、Ta、Zr、Hf、Ti 等)的强烈亏损是与俯冲有关岩浆岩的典型特征(Kelemen et al., 1990; Stolz et al., 1996),但白石冈岩体形成于板内环境,其 Nb、Ta、Ti 等高场强元素亏损的




图 11 白石岗岩体(Y+Nb)-Rb关系图

Fig. 11 (Y+Nb)-Rb diagram of the Baishigang pluton. The base map is after Pearce et al. (1984), and the range of the post-CEG is after Forster et al. (1997); VAG—volcanic arc granites; ORG—ocean ridge granites; WPG—within plate granites; syn-COLG—syn-collision granites; post-CEG—post-collision extension granites

底图据 Pearce 等(1984); 图中 post-CEG 区域据 Forster 等(1997)资料圈定; VAG—火山弧花岗岩; ORG—洋脊花岗岩; WPG—板内花岗岩; syn-COLG—同碰撞花岗岩; post-CEG—后碰撞伸展花岗岩

特征应是从源岩中继承而来的,而不应被视为产于俯冲环境的标志,因为典型地壳熔体高场强元素含量很低(Ryerson et al., 1987)。基于以上的讨论可以设想,自燕山早期以来,南岭地区由于后造山拉张裂解作用,导致软流圈上涌和岩石圈伸展减薄,由此诱发幔源基性岩浆底侵于地壳下部。在断裂引起的减压作用和幔源基性岩浆底侵带来足够热量的影响下,促使了地壳物质的部分熔融形成长英质岩浆。幔源基性岩浆与长英质岩浆在深部岩浆房混合产生混染的母岩浆,后者再经高分异的演化,即形成本文所讨论的白石岗岩体,这一模式较好地解释了目前所观测到的岩体的地质地球化学特征。

6 结论

(1) 白石岗岩体锆石 U-Pb 年龄为 148.5 ± 1.6 Ma, 属晚侏罗世岩浆活动的产物。

(2) 白石岗岩体铝弱过饱和,富硅,富钾,富 Rb、Th、U、Pb 和轻稀土,贫 Ba、Sr、P、Ti, Rb/Sr 比值高, K/Rb 比值低, 铕负异常显著。其 Nb、Ta、Zr、Hf 等高场强元素含量及 Ga/Al 比值均较之典型 A 型花岗岩偏低, 并具偏高的 $\epsilon_{Nd}(t)$ 值 ($-5.99 \sim$

-7.51) 和偏低的 T_{DM} 值 ($1.42 \sim 1.54$ Ga), 综合地球化学资料指示岩体应属高分异的 I 型花岗岩。

(3) 白石岗岩体形成于后造山的伸展引张环境, 是由底侵的幔源基性岩浆及其诱发的长英质岩浆在深部岩浆房混合, 并经高程度分离结晶的产物。

致谢: 本研究得到周新民教授的指导, 并承审阅论文初稿, 谨此致以诚挚谢意!

注 释

① 广东省地质局. 1965. 紫金幅区域地质测量报告(1:200000).

参 考 文 献

- 包志伟, 赵振华. 2003. 佛冈铝质 A 型花岗岩的地球化学及其形成环境初探. 地质地球化学, 31(1): 52~61.
- 陈江峰, 郭新生, 汤加富, 周泰禧. 1999. 中国东南地壳增长与 Nd 同位素模式年龄. 南京大学学报(自然科学), 35(6): 649~658.
- 陈培荣, 孔兴功, 倪琦生, 章邦桐, 刘昌实. 1999. 赣南燕山早期双峰式火山岩的厘定和意义. 地质论评, 45(增刊): 734~741.
- 陈小明, 王汝成, 刘昌实, 胡欢, 张文兰, 高剑锋. 2002. 广东从化佛冈(主体)黑云母花岗岩定年和成因. 高校地质学报, 8(3): 293~307.
- 范蔚茗, 王岳军, 郭锋, 彭头平. 2003. 湘赣地区中生代镁铁质岩作用与岩石圈伸展. 地学前缘, 10(3): 159~169.
- 广东省地质矿产局. 1988. 广东省区域地质志. 北京: 地质出版社, 477~509.
- 郭令智, 施央申, 马瑞士. 1983. 西太平洋中生代活动大陆边缘和岛弧构造的形成及演化. 地质学报, 57(1): 11~21.
- 洪大卫, 郭文歧, 李戈晶, 康炜, 徐海明. 1987. 福建沿海晶洞花岗岩带的岩石学和成因演化. 北京: 北京科学技术出版社, 1~132.
- 刘昌实, 朱金初, 沈渭洲, 徐士进. 1990. 华南陆壳改造系列花岗岩类型划分和成岩物质来源. 地质学报, 64(1): 43~52.
- 陆松年, 李惠民. 1991. 蓟县长城系大红峪组火山岩的单颗粒锆石 U-Pb 法准确定年. 中国地质科学院院报, 22: 137~146.
- 毛景文, 李红艳, 裴荣富. 1995. 湖南千里山花岗岩体的 Nd-Sr 同位素及岩石成因研究. 矿床地质, 14(3): 235~242.
- 邱检生, 胡建, McInnes B I A, 蒋少涌, 徐夕生, Allen C M. 2004a. 广东龙窝花岗岩闪长质岩体的年代学、地球化学及岩石成因. 岩石学报, 20(6): 1363~1374.
- 邱检生, McInnes B I A, 徐夕生, Allen C M. 2004b. 赣南大吉山五里亭岩体的锆石 ELA-ICP-MS 定年及其与钨成矿关系的新认识. 地质论评, 50(2): 125~133.
- 任纪舜, 陈廷愚, 牛宝贵, 刘志刚, 刘凤仁. 1990. 中国东部及邻区大陆岩石圈的构造演化与成矿. 北京: 科学出版社, 1~205.
- 沈渭洲, 朱金初, 刘昌实, 徐士进, 凌洪飞. 1993. 华南基底变质岩的 Sm-Nd 同位素及其对花岗岩类物质来源的制约. 岩石学报, 9(2): 115~124.
- 舒良树, 周新民. 2002. 中国东南部晚中生代构造作用. 地质论评, 48(3): 249~260.
- 孙涛, 陈培荣, 周新民, 王汝成, 王志成. 2002. 南岭东段强过铝质花岗岩中白云母研究. 地质论评, 48(5): 518~525.
- 王鸿楨, 杨森楠, 李思田. 1983. 中国东南部及邻区中、新生代盆地发育及大陆边缘区的构造发展. 地质学报, 57(3): 213~223.

- 王银喜, 杨杰东, 陶仙聪, 李惠民. 1988. 化石、矿物和岩石样品的 Sm-Nd 同位素实验方法及其应用. 南京大学学报(自然科学版), (2): 297~308.
- 袁忠信, 张宗清. 1992. 南岭花岗岩类岩石 Sm、Nd 同位素特征及岩石成因探讨. 地质论评, 38(1): 1~15.
- 张文兰, 华仁民, 王汝成, 李惠民, 陈培荣. 2004. 江西大吉山五里亭花岗岩单颗粒锆石 U-Pb 同位素年龄及其地质意义探讨. 地质学报, 78(3): 352~358.
- 赵振华, 包志伟, 张伯友. 1998. 湘南中生代玄武岩类地球化学特征. 中国科学(D辑), 28(增刊): 7~14.
- 赵子杰, 马大铨, 林惠坤, 袁春林, 张小豪. 1987. 广东龙窝、佛岗花岗岩类岩体地球化学特征及其形成的地质背景. 见: 宜昌地质矿产研究所编. 南岭地质矿产科研报告集(一). 武汉: 武汉地质学院出版社, 28~68.
- 郑永飞, 魏春生, 王峥嵘, 黄耀生, 张宏. 1997. 大龙山岩体的冷却史及其成矿关系的同位素研究. 地质科学, 32(4): 465~477.
- 朱金初, 李向东, 沈渭洲, 王银喜, 杨杰东. 1989. 广西花山复式花岗岩体成因的铷、钕和氧同位素研究. 地质学报, 63(3): 225~235.
- Chunfang. 2002. Early Yanshanian post-orogenic granitoids in the Nanling region; petrological constraints and geodynamic settings. Science in China, 45(8): 755~768.
- Chen Xiaoming, Wang Rucheng, Liu Changshi, Hu Huan, Zhang Wenlan, Gao Jianfeng. 2002. Isotopic dating and genesis for Fogang biotite granites of Conghua area, Guangdong province. Geol. J. China Uni., 8(3): 293~307 (in Chinese with English abstract).
- Compston W, Williams I S, Kirschvink J L, Zhang Z, Ma G. 1992. Zircon U-Pb ages for the Early Cambrian time-scale. J. Geol. Soc. London, 149: 171~184.
- Dodson M H, McClelland-Brown E. 1985. Isotopic and palaeomagnetic evidence for rates of cooling, uplift and erosion. In: The chronology of geological record. Geol. Soc. Mem. (London), 10: 315~325.
- Dostal J, Chatterjee A K. 2000. Contrasting behaviour of Nb/Ta and Zr/Hf ratios in a peraluminous granitic pluton (Nova Scotia, Canada). Chem. Geol., 163: 207~218.
- Eby G N. 1990. The A-type granitoids; A review of their occurrence and chemical characteristics and speculations on their petrogenesis. Lithos., 26: 115~134.
- Engelbreton D C, Cox A, Gordon R G. 1985. Relative motions between oceanic and continental plates in the Pacific basin. Geol. Soc. Am. Spec. Paper, 1~59.
- Fan Weiming, Wang Yunjun, Guo Feng, Peng Touping. 2003. Mesozoic mafic magmatism in Hunan-Jiangxi Provinces and the lithospheric extension. Earth Science Frontiers, 10(3): 159~169 (in Chinese with English abstract).
- Faure G. 1986. Principles of isotope geology. New York: Jones Willey & Sons, 141~249.
- Forster H J, Tischendorf G, Trumbull R B. 1997. An evaluation of the Rb vs. (Y+Nb) discrimination diagram to infer tectonic setting of silicic igneous rocks. Lithos., 40: 261~293.
- Fujimaki H. 1986. Partition coefficients of Hf, Zr and REE between zircon, apatite and liquid. Contrib. Mineral Petrol., 94: 42~45.
- Gilder S A, Gill J, Coe R S, Zhao X, Liu Z, Wang G, Yuan K, Liu W, Kuang G, Wu H. 1996. Isotopic and paleomagnetic constraints on the Mesozoic tectonic evolution of south China. J. Geophys Res., 101: 16137~16154.
- Green T H, Pearson N J. 1986. Rare earth element partitioning between titanite and coexisting silicate liquid at high pressure and temperature. Chem. Geol., 74: 201~216.
- Guo Lingzhi, Shi Yangshen, Ma Ruishi. 1983. On the formation and evolution of the Mesozoic-Cenozoic active continental margin and island arc tectonics of the western Pacific ocean. Acta Geologica Sinica, 57(1): 11~21 (in Chinese with English abstract).
- Hanson G N. 1978. The application of trace elements to the petrogenesis of igneous rocks of granitic compositions. Earth Planet Sci. Lett., 38: 26~43.
- Hong Dawei, Guo Wenqi, Li Gejing, Kang Wei, Xu Haiming. 1987. Petrology of the miarolitic granite belt in the southeast coast of Fujian province and their petrogenesis. Beijing: Science and Technology Press of Beijing, 1~132 (in Chinese with English abstract).
- Hong Dawei, Xie Xilin, Zhang Jisheng. 1998. Isotope geochemistry of granitoids in south China and their metallogeny. Resource Geology, 48: 251~263.

References

- Arth J G. 1976. Behaviour of trace elements during magmatic processes; a summary of theoretical models and their applications. J. Res. U. S. Geol. Survey, 4: 41~47.
- Bao Zhiwei, Zhao Zhenhua. 2003. Geochemistry and tectonic setting of the Fogang aluminous A-type granite, Guangdong province, China—a preliminary study. Geology-Geochemistry, 31(1): 52~61 (in Chinese with English abstract).
- Barbarin B. 1999. A review of the relationships between granitoid types, their origins and their geodynamic environments. Lithos., 46: 605~626.
- Boynton W V. 1984. Geochemistry of the rare earth elements: meteorite studies. In: Henderson P, ed. Rare Earth Elements Geochemistry. Amsterdam: Elsevier, 63~144.
- Bureau of Geology and Mineral Resources of Guangdong Province. 1988. Regional Geology of Guangdong Province. Beijing: Geology Press, 477~509 (in Chinese with English abstract).
- Chappell B W. 1999. Aluminum saturation in I- and S-type granite and the characterization of fractionated haplogranite. Lithos., 46: 535~551.
- Chappell B W, White A J R. 2001. Two contrasting granite types: 25 years later. Australia J. Earth Sci., 48: 489~499.
- Chen A. 2001. Mirror-image tectonics in the South China orogenic belt: tectonic evidence from western Fujian, Southeastern China. Tectonophysics, 305: 497~519.
- Chen J F, Jahn B M. 1998. Crustal evolution of southeastern China: Nd and Sr isotopic evidence. Tectonophysics, 284: 101~133.
- Chen Jiangfeng, Guo Xincheng, Tang Jiafu, Zhou Taixi. 1999. Nd isotopic model ages; implications of the growth of the continental crust of Southeastern China. J. Nanjing University, 35(6): 649~658 (in Chinese with English abstract).
- Chen Peirong, Kong Xinggong, Ni Qisheng, Zhang Bangtong, Liu Changshi. 1999. Ascertainment and implication of the Early Yanshanian bimodal volcanic associations from south Jiangxi province. Geological Review, 45 (Suppl): 734~741 (in Chinese with English abstract).
- Chen Peirong, Hua Renmin, Zhang Bangtong, Lu Jianjun, Fan

- Kelemen P B, Kingler R J, Johnson K T M. 1990. High field strength element depletions in arc basalt due to mantle-magma interaction. *Nature*, 345:521~524.
- Li X H, Chen Z, Liu D Y, Li W X. 2003. Jurassic gabbro-granite-syenite suites from Southern Jiangxi province, SE China: age, origin, and tectonic significance. *International Geology Review*, 45(10):898~921.
- Liegeois J P, Navez J, Hertogen J, Black R. 1998. Contrasting origin of post-collisional high-K calc-alkaline and shoshonitic versus alkaline and peralkaline granitoids, the use of sliding normalization. *Lithos.*, 45:1~28.
- Liu Changshi, Zhu Jinchu, Shen Weizhou, Xu Shijin. 1990. Classification and source materials of continental crust transformation series granitoids in south China. *Acta Geologica Sinica*, 64(1):43~52 (in Chinese with English abstract).
- Lu Songnian, Li Huimin. 1991. A precise U-Pb single zircon dating for the volcanics of Dahongyu Formation, Changcheng System in Jixian. *Bulletin of the Chinese Academy of Geological Sciences*, 22: 137~146 (in Chinese with English abstract).
- Mahood G, Hildreth W. 1983. Large partition coefficients for trace elements in high-silica rhyolites. *Geochim Cosmochim Acta*, 47: 11~30.
- Maniar P, Piccoli P. 1989. Tectonic discrimination of granitoids. *Geol. Soc. Amer Bull*, 101:635~643.
- Mao Jingwen, Li Hongyan, Pei Rongfu. 1995. Nd-Sr isotopic and petrogenetic studies of the Qianlishan granite stock, Hunan province. *Mineral Deposits*, 14(3):235~242 (in Chinese with English abstract).
- Martin H, Bonin B, Capdevila R, Jahn B M, Lameyre J, Wang Y. 1994. The Kuiqi peralkaline granitic complex (SE China): petrology and geochemistry. *Journal Petrology*, 35:983~1015.
- Maruyama S, Seno T. 1986. Orogeny and relative plate motions; example of the Japanese island. *Tectonophysics*, 127 (3~4): 305~329.
- Pearce J, Harris N B W, Tindle A G. 1984. Trace element discrimination diagrams for the tectonic interpretation of granitic rocks. *J. Petrol.*, 25:956~983.
- Qiu Jiansheng, Hu Jian, McInnes B I A, Jiang Shaoyong, Xu Xisheng, Allen C M. 2004a. Geochronology, geochemistry and petrogenesis of the Longwo granodioritic pluton in Guangdong province. *Acta Petrologica Sinica*, 20(6):1363~1374 (in Chinese with English abstract).
- Qiu Jiansheng, McInnes B I A, Xu Xisheng, Allen C M. 2004b. Zircon ELA-ICP-MS dating for Wuliting pluton at Dajishan, Southern Jiangxi and new recognition about its relation to Tungsten Mineralization. *Geological Review*, 50(2): 125~133 (in Chinese with English abstract).
- Raith J G. 1995. Petrogenesis of the Concordia granite gneiss and its relation to W-Mo mineralization in western Namaqualand, South Africa. *Precambrian Research*, 70:303~335.
- Ren Jishun, Chen Tingyu, Niu Baogui, Liu Zhigang, Liu Fengren. 1990. Tectonic evolution of the continental lithosphere and metallogeny in eastern China and adjacent areas. Beijing: Science Press, 1~205 (in Chinese).
- Ryerson F J, Watson E B. 1987. Rutile saturation in magmas; implications for Ti-Nb-Ta depletion in island-arc basalts. *Earth Planet Sci. Lett.*, 86: 225~239.
- Sewell R J, Chan L S, Fletcher C J N, Brewer T S, Zhu J C. 2000. Isotope zonation in basement crustal blocks of southeastern China; evidence for multiple terrane amalgamation. *Episodes*, 23(4):257~261.
- Shen Weizhou, Zhu Jinchu, Liu Changshi, Xu Shijin, Ling Hongfei. 1993. Sm-Nd isotopic study of basement metamorphic rocks in south China and its constraint on material sources of granitoids. *Acta Petrologica Sinica*, 9(2):115~124 (in Chinese with English abstract).
- Shen Weizhou, Ling Hongfei, Li Wuxian, Wang Dezi. 2000. Crust evolution in southeast China; evidence from Nd model ages of granitoids. *Science in China (Series D)*, 43: 36~49.
- Shu Liangshu, Zhou Xinmin. 2002. Late Mesozoic tectonism of southeast China. *Geological Review*, 48(3): 249~260 (in Chinese with English abstract).
- Shu Liangshu, Deng Ping, Wang Bin, Tan Zhenzhong, Yu Xingqi, Sun Yan. 2004. Lithology, kinematics and geochronology related to Late Mesozoic basin-mountain evolution in the Nanxiong-Zhuguang area, South China. *Science in China (Series D)*, 47(8): 673~688.
- Stolz A J, Jochum K P, Spettel B, Hofmann A W. 1996. Fluid and melt-related enrichment in the subarc mantle; evidence from Nb/Ta variations in island-arc basalts. *Geology*, 24:587~590.
- Sun S S, McDonough W F. 1989. Chemical and isotopic systematics of oceanic basalts; Implications for mantle composition and processes. In: Saunderson A D, Norry M J, eds. *Magmatism in the Ocean Basins*. *Geol. Soc. Spe. Publ.*, 42: 313~345.
- Sun Tao, Chen Peirong, Zhou Xinmin, Wang Rucheng, Wang Zhicheng. 2002. Strongly peraluminous granites in eastern Nanling mountains; China; study on muscovites. *Geological Review*, 48(5): 518~525 (in Chinese with English abstract).
- Wang Hongzhen, Yang Sennan, Li Sitian. 1983. Mesozoic and Cenozoic basin formation in east China and adjacent regions and development of the continental margin. *Acta Geologica Sinica*, 57(3): 213~223 (in Chinese with English abstract).
- Wang Yunjun, Zhang Y H, Fan Weimin. 2002. Numerical modeling for generation of Indo-Sinian peraluminous granitoids in Hunan Province; basaltic underplating vs. tectonic thickening. *Science in China (Series D)*; 45(11):1042~1056.
- Wang Y J, Fan W M, Guo F. 2003. Geochemistry of Early Mesozoic potassium-rich dioritic-granodioritic intrusions in southeastern Hunan province, south China; petrogenesis and tectonic implications. *Geochim. J.*, 37(4):427~448.
- Wang Yinxi, Yang Jiedong, Tao Xiancong, Li Huimin. 1988. A study of the Sm-Nd method for fossil, mineral and rock samples, and its application. *J. Nanjing Uni. (Natural Sciences Edition)*, 24(2):297~308 (in Chinese with English abstract).
- Whalen J B, Currie K L, Chappell B W. 1987. A-type granites: geochemical characteristics, discrimination and petrogenesis. *Contrib. Min. Petrol.*, 95: 407~419.
- Xu Xisheng, Deng Ping, O'Reilly S Y, Griffin W L, Zhou Xinmin, Tan Zhenzhong. 2003. Single zircon LAM-ICPMS U-Pb dating of Guidong complex (SE China) and its petrogenetic significance. *Chinese Science Bulletin*, 48(17): 1892~1899.
- Yuan Zhongxin, Zhang Zongqing. 1992. Sm-Nd isotopic characteristics of granitoids in the Nanling region and their petrogenetic analysis. *Geological Review*, 38(1): 1~15 (in Chinese with English abstract).
- Yurimoto H, Duck E F, Papike J J, Shearer C K. 1990. Are

- discontinuous chondrite-normalized REE patterns in pegmatitic granite systems the results of monazite fractionation? *Geochim Cosmochim Acta*, 54: 2141~2145.
- Zhang Wenlan, Hua Renmin, Wang Rucheng, Li Huimin, Chen Peirong. 2004. Single zircon U-Pb age of the Wuliting granite in Dajishan area of Jiangxi, and its geological implication. *Acta Geologica Sinica*, 78(3): 352~358 (in Chinese with English abstract).
- Zhao Zhenhua, Bao Zhiwei, Zhang Boyou. 1998. The geochemistry of Mesozoic basalts in south Hunan Province, South China. *Science in China (Series D)*, 28(Suppl): 7~14 (in Chinese).
- Zhao Zijie, Ma Daquan, Lin Huikun, Yuan Chunlin, Zhang Xiaohao. 1987. The geochemical characteristics and geological setting of the formation of Longwo and Fogang granitoid batholiths, Guangdong province. In: *Yichang Institute of Geology and Mineral Resource*, ed. *Research Reports of the Geology and Mineral Resources of Nanling (Series 1)*. Wuhan: Wuhan College of Geology Press, 28~68 (in Chinese with English abstract).
- Zheng Yongfei, Wei Chunsheng, Wang Zhengrong, Huang Yaosheng, Zhang Hong. 1997. An isotope study on the cooling history of the Dalongshan granitic massif and its bearing on mineralization process. *Scientia Geologica Sinica*, 32(4): 465~477 (in Chinese with English abstract).
- Zhou X M, Li W X. 2000. Origin of Late Mesozoic igneous rocks in Southeastern China: implications for lithosphere subduction and underplating of mafic magmas. *Tectonophysics*, 326: 259~277.
- Zhu Jinchu, Li Xiangdong, Shen Weizhou, Wang Yinxi, Yang Jiedong. 1989. Sr, Nd and O isotope studies on the genesis of the Huashan granite complex. *Acta Geologica Sinica*, 63(3): 225~235 (in Chinese with English abstract).

The Baishigang Pluton in Heyuan, Guangdong Province: A Highly Fractionated I-type Granite

QIU Jiansheng, HU Jian, WANG Xiaolei, JIANG Shaoyong, WANG Rucheng, XU Xisheng
State Key Laboratory for Mineral Deposit Research, Department of Earth Sciences, Nanjing University, 210093

Abstract

The Baishigang Pluton is located at the eastern end of the E-W-trending Fogang granite belt, and has an outcrop area of about 440 km². It intruded into the Lower Jurassic Lantang Group and is made up mainly of medium- to coarse-grained biotite granites. The major mineral phases of the granites are quartz (25%~35%), microperthite (45%~50%), plagioclase ($An=20\sim30$, 15%~20%) and biotite (5%~10%). Zircon U-Pb isotopic dating yields an age of 148.5 ± 1.6 Ma, indicating that it was formed in the Late Jurassic. Chemical analyses show that the granites have high SiO₂ (= 73.37%~76.05%) and moderate Al₂O₃ (12.00%~13.16%). They are slightly peraluminous with A/NKC values ranging mainly from 1.0 to 1.1 and normative corundum <1%. The rocks also have moderate to low alkaline contents ($K_2O+Na_2O=7.44\%\sim 8.48\%$, NK/A [molecular $(K_2O+Na_2O)/Al_2O_3$] = 0.75~0.88) and high potassium contents ($K_2O=4.22\%\sim 5.29\%$, $K_2O/Na_2O=1.31\sim 1.70$), thus can be ascribed to the rocks of the high-K calc-alkaline series. Trace and rare earth elements of the granites are characterized by rich Rb, Th, U, Pb and LREE, and poor Ba, Sr, P, Ti, and have high Rb/Sr (2.36~20.82) and low K/Rb ratios (<150 for most of the samples). Chondrite-normalized REE patterns are right-inclined and display significant negative europium anomalies ($\delta Eu=0.05\sim 0.28$). Their HFSE (e. g., Nb, Ta, Zr, Hf) concentrations and Ga/Al ratios are lower than that of the respective values of A-type granites. The granites have a more enriched Nd isotopic composition with initial ϵ_{Nd} values from -5.99 to -7.51, and show younger depleted-mantle model ages ($T_{DM}=1.42\sim 1.54$ Ga). Integrated geochemical data suggest that the rocks are highly fractionated I-type granites. By combining with comprehensively analyses of the tectonic settings, we suggest that the Baishigang pluton was formed in a post-collisional tensile environment, and the granites were generated by a two-stage process including mixing of underplating basaltic magma with the induced felsic melts in the source region and subsequent high-degree fractional crystallization.

Key words: highly fractionated I-type granite; crust-mantle interaction; geochemistry; petrogenesis; Baishigang pluton; Guangdong Province