

A RENEWAL THEOREM FOR STRONGLY ERGODIC MARKOV CHAINS IN DIMENSION $d \geq 3$ AND CENTERED CASE.

D. GUIBOURG and L. HERVÉ ¹

AMS subject classification : 60J10-60K05-47A55

Keywords : Markov chains, renewal theorem, spectral method.

Abstract. *In dimension $d \geq 3$, we present a general assumption under which the renewal theorem established by Spitzer [23] for i.i.d. sequences of centered nonlattice r.v. holds true. Next we appeal to an operator-type procedure to investigate the Markov case. Such a spectral approach has been already developed by Babillot [1], but the weak perturbation theorem of [18] enables us to greatly weaken the moment conditions of [1]. Our applications concern the v -geometrically ergodic Markov chains, the ρ -mixing Markov chains, and the iterative Lipschitz models, for which the renewal theorem of the i.i.d. case extends under the (almost) expected moment condition.*

1 Introduction

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let (E, \mathcal{E}) be a measurable space, and let $(X_n, S_n)_{n \geq 0}$ be a sequence of random variables (r.v.) defined on Ω and taking values in $E \times \mathbb{R}^d$. We denote by $\mathcal{L}_d(\cdot)$ the Lebesgue-measure on \mathbb{R}^d , by $\langle \cdot, \cdot \rangle$ the canonical scalar product on \mathbb{R}^d , by $\|\cdot\|$ the euclidean norm in \mathbb{R}^d , and by $B(\mathbb{R}^d)$ the Borel σ -algebra on \mathbb{R}^d , and we use the notation “*” to denote the matrix transposition.

Throughout this paper we assume that $d \geq 3$. Given some fixed nonnegative measurable function f on E , we are interested in the asymptotic behavior of the renewal-type measures $\sum_{n=1}^{+\infty} \mathbb{E}[f(X_n) 1_A(S_n - a)]$ ($A \in B(\mathbb{R}^d)$) when $a \in \mathbb{R}^d$ goes to infinity. Specifically we will consider the centered case in a sense that will be specified later. In the i.i.d. case, this corresponds to the well-known result established by Spitzer [23]: if $(X_n)_{n \geq 1}$ is a i.i.d. sequence of centered nonlattice r.v. such that $\mathbb{E}[|X_0|^{m_d}] < \infty$ with $m_d = \max\{d - 2, 2\}$, and if the covariance $d \times d$ -matrix $\Sigma = \mathbb{E}[X_0 X_0^*]$ is invertible, then the associated random walk $S_n = \sum_{k=1}^n X_k$ satisfies the following property when $\|a\| \rightarrow +\infty$:

$$\forall g \in \mathcal{C}_c(\mathbb{R}^d), \quad \sum_{n=1}^{+\infty} \mathbb{E}_\mu [g(S_n - a)] \sim \frac{C_d}{\langle \Sigma^{-1}a, a \rangle^{\frac{d-2}{2}}} \mathcal{L}_d(g) \quad (1)$$

with

$$C_d = 2^{-1} \pi^{-\frac{d}{2}} (\det \Sigma)^{-\frac{1}{2}} \Gamma\left(\frac{d-2}{2}\right), \quad (2)$$

where $\Gamma(\cdot)$ denotes the usual Gamma-function and $\mathcal{C}_c(\mathbb{R}^d)$ stands for the space of complex-valued compactly supported continuous functions on \mathbb{R}^d .

¹Université Européenne de Bretagne, I.R.M.A.R. (UMR-CNRS 6625), Institut National des Sciences Appliquées de Rennes. Denis.Guibourg@ens.insa-rennes.fr, Loic.Herve@insa-rennes.fr

In Section 2 we present some general assumptions on the characteristic-type functions $\mathbb{E}[f(X_n) e^{i\langle t, S_n \rangle}]$ under which the previous renewal theorem extends. In fact these assumptions are the “tailor-made” conditions for proving a multidimensional renewal theorem in the centered case with the help of the Fourier techniques. This part extends and specifies (and sometimes simplifies) some of the arguments and computations used in Babillot’s paper [1].

The purpose of Section 3 is to investigate the general assumptions of Section 2 in the case when $(X_n)_{n \geq 0}$ is a Markov chain. In this context we show in Subsection 3.1 that a natural way is to assume that, for each $t \in \mathbb{R}^d$, the linear operators $Q_n(t)$ defined by $(Q_n(t)f)(x) := \mathbb{E}_x[f(X_n) e^{i\langle t, S_n \rangle}]$ ($x \in E$) is a semi-group: $Q_{m+n}(t) = Q_n(t) \circ Q_m(t)$. The Markov random walks satisfy this assumption. We then have $\mathbb{E}_x[f(X_n) e^{i\langle t, S_n \rangle}] = (Q_1(t)^n f)(x)$, and if $Q_1(0)$ satisfies some “good” spectral properties on a Banach space \mathcal{B} , then the operator-perturbation method first introduced by Nagaev [21] can be used to obtain an accurate expansion of $Q_1(t)^n f$ for any $f \in \mathcal{B}$, thus of $\mathbb{E}_x[f(X_n) e^{i\langle t, S_n \rangle}]$. This spectral method has been already exploited to investigate the Markov renewal theorems, see e.g. [1, 12, 4, 14, 8]. In particular in [1], Property (1) is extended to centered Markov random walks in dimension $d \geq 3$. However, because of the use of the standard perturbation theory, the assumptions in [1] (and in all the previously cited works) need some operator-moment conditions which are restrictive, even not fulfilled in general when S_1 is unbounded.

In this work, we use the weak Nagaev method introduced in [15, 16, 2, 13], which allows to greatly weaken the moment conditions in limit theorems for strongly ergodic Markov chains and dynamical systems. This new approach, in which the Keller-Liverani perturbation theorem [18, 19, 10] plays a central role, is fully described in [17] and applied to prove some usual refinements of the central limit theorem (CLT). It is used in [11] to establish a one-dimensional (non-centered) Markov renewal theorem. Also mention that one of the more beautiful applications of this method concerns the convergence to stable laws, see [13, 5, 9].

This new approach is outlined in Subsection 3.2 and applied in Section 4 to the three following classical strongly ergodic Markov models:

- *The v -geometrically ergodic Markov chains [20],*
- *The ρ -mixing Markov chains [22],*
- *The iterative Lipschitz models [7].*

To have a good understanding of our improvements in term of moment conditions, let us consider an unbounded function $v : E \rightarrow [1, +\infty)$ and a v -geometrically ergodic Markov chain $(X_n)_{n \geq 0}$, that is we have: $\sup_{|f| \leq v} \sup_{x \in E} |Q^n f(x) - \pi(f)|/v(x) = O(\kappa^n)$ for some $\kappa \in (0, 1)$, where Q denotes the transition probability of the chain and π is the stationary distribution. Besides, let us consider the Markov random walk $S_n = \sum_{k=1}^n \xi(X_k)$ in \mathbb{R}^d associated to some measurable function $\xi : E \rightarrow \mathbb{R}^d$. The centered case corresponds to the assumption $\pi(\xi) = 0$, the (asymptotic) covariance matrix is given by $\Sigma := \lim_n \mathbb{E}[S_n S_n^*]/n$, provided that this limit exists, and finally the (Markov) nonlattice condition means that there exist no $b \in \mathbb{R}^d$, no closed subgroup H in \mathbb{R}^d , $H \neq \mathbb{R}^d$, no π -full Q -absorbing set $A \in \mathcal{E}$, and finally no bounded measurable function $\theta : E \rightarrow \mathbb{R}^d$ such that: $\forall x \in A, \xi(y) + \theta(y) - \theta(x) \in b + H$ $Q(x, dy)$ -p.s. We shall prove in Subsection 4.1 the next statement:

Let us assume that $(X_n)_{n \geq 0}$ is a v -geometrically ergodic Markov chain, that $\xi : E \rightarrow \mathbb{R}^d$

($d \geq 3$) is centered, nonlattice, and such that $\|\xi\|^{m_d+\delta_0} \leq C v$ with $m_d = \max\{d-2, 2\}$ and some constants $C, \delta_0 > 0$, and finally that the initial distribution μ of the chain is such that $\mu(v) < \infty$. Then the above asymptotic covariance matrix Σ is well-defined and positive definite, and $S_n = \sum_{k=1}^n \xi(X_k)$ satisfies (1).

In comparison with the optimal order m_d of the i.i.d. case, the condition $\|\xi\|^{m_d+\delta_0} \leq C v$ is the expected moment assumption (up to $\delta_0 > 0$) in the context of v -geometrically ergodic Markov chains. The corresponding result in [1] requires the following assumption: $\sup_{x \in E} \frac{1}{v(x)} \int_E |\xi(y)|^{m_d+\delta_0} v(y) Q(x, dy) < \infty$, see also [8]. This moment condition is not only clearly stronger than the previous one, but actually it is not fulfilled in general when ξ is unbounded. A typical example is presented in [17, § 3] for the usual linear autoregressive model and $\xi(x) = x$. Similar improvements concerning the moment conditions are obtained in Subsections 4.2 and 4.3 for the two others (above cited) Markov models.

Throughout this paper we shall use the following definitions. Let \mathcal{O} be an open subset of \mathbb{R}^d , let $(X, \|\cdot\|_X)$ be a normed vector space, and let $\tau \in [0, 1)$. We say that a function $V : \mathcal{O} \rightarrow X$ is uniformly τ -holderian if there exists a constant $C \geq 0$ such that we have:

$$\forall (t, t') \in \mathcal{O}^2, \|V(t) - V(t')\|_X \leq C \|t - t'\|^\tau.$$

If $m \in \mathbb{R}_+$, we shall say that $U \in \mathcal{C}_b^m(\mathcal{O}, X)$ if U is a function from \mathcal{O} to X satisfying the following properties :

U is $\lfloor m \rfloor$ -times continuously differentiable on \mathcal{O}

Each partial derivative of order $j = 0, \dots, \lfloor m \rfloor$ of U is bounded on \mathcal{O}

Each partial derivative of order $\lfloor m \rfloor$ of U is uniformly $(m - \lfloor m \rfloor)$ -holder on \mathcal{O} .

2 A general statement

In this section, we state a general result that will be applied later to the Markov context, but which has its own interest in view of other possible applications. Here $(X_n, S_n)_{n \geq 1}$ is a general sequence of random variables defined on Ω and taking values in $E \times \mathbb{R}^d$.

For any $R > 0$, we set $B(0, R) := \{t \in \mathbb{R}^d : \|t\| < R\}$, and for any $0 < r < r'$, we set $K_{r,r'} = \{t \in \mathbb{R}^d : r < \|t\| < r'\}$. We denote by ∇ and $Hess$ the gradient and the Hessian matrix, respectively.

Hypothesis $\mathcal{R}(m)$. *Given a real number $m > 0$ and a measurable function $f : E \rightarrow [0, +\infty)$ such that $\mathbb{E}[f(X_n)] < \infty$ for all $n \geq 1$, we will say that Hypothesis $\mathcal{R}(m)$ holds if:*

(i) *There exists $R > 0$ such that we have for all $t \in B(0, R)$ and all $n \geq 1$:*

$$\mathbb{E}[f(X_n) e^{i\langle t, S_n \rangle}] = \lambda(t)^n L(t) + R_n(t), \quad (3)$$

where $\lambda(0) = 1$, the functions $\lambda(\cdot)$ and $L(\cdot)$ are in $C_b^m(B(0, R), \mathbb{C})$, and the series $\sum_{n \geq 1} R_n(\cdot)$ uniformly converges on $B(0, R)$ and defines a function in $C_b^m(B(0, R), \mathbb{C})$.

(ii) *For all $0 < r < r'$, the series $\sum_{n \geq 1} \mathbb{E}[f(X_n) e^{i\langle \cdot, S_n \rangle}]$ uniformly converges on $K_{r,r'}$ and defines a function in $C_b^m(K_{r,r'}, \mathbb{C})$.*

If Hypothesis $\mathcal{R}(m)$ holds with $m \geq 2$, we define the following symmetric matrix:

$$\Sigma := -\text{Hess } \lambda(0).$$

In the sequel, $\vec{m} := \frac{\nabla \lambda(0)}{i}$ will be viewed as an asymptotic first moment vector in \mathbb{R}^d , and we shall assume that $\vec{m} = 0$. Under this centered assumption, $\Sigma = -\text{Hess } \lambda(0)$ may be viewed as an asymptotic covariance matrix. These facts are specified in the following proposition, in which we use the notation $F^{(\ell)}$ for the derivative of order ℓ of a complex-valued function F .

Proposition 1

(i) If Condition $\mathcal{R}(1)$ -(i) is fulfilled with $f = 1_E$, $L(0) = 1$, $\sup_{n \geq 1} |R_n^{(1)}(0)| < \infty$, and if $\forall n \geq 1$, $\mathbb{E}[|S_n|] < \infty$, then we have: $\frac{\nabla \lambda(0)}{i} = \lim_n \frac{1}{n} \mathbb{E}[S_n]$.

(ii) If Condition $\mathcal{R}(2)$ -(i) is fulfilled with $f = 1_E$, $L(0) = 1$, $\sup_{n \geq 1} |R_n^{(2)}(0)| < \infty$, and if in addition we have $\nabla \lambda(0) = 0$ and $\forall n \geq 1$, $\mathbb{E}[|S_n|^2] < \infty$, then we have: $\Sigma = \lim_n \frac{1}{n} \mathbb{E}[S_n S_n^*]$.

Proof. To simplify the computations, let us assume $d = 1$. The extension to $d \geq 2$ is obvious by using partial derivatives. By deriving $\mathbb{E}[e^{itS_n}] = \lambda(t)^n L(t) + R_n(t)$ at $t = 0$, we obtain: $i \mathbb{E}[S_n] = n \lambda^{(1)}(0) + L^{(1)}(0) + R_n^{(1)}(0)$. Hence Assertion (i). Next by deriving twice the same equality at $t = 0$, one gets: $-\mathbb{E}[S_n^2] = n \lambda^{(2)}(0) + L^{(2)}(0) + R_n^{(2)}(0)$. Hence Assertion (ii). \square

Notice that the assumption $\sup_{n \geq 1} |R_n^{(\ell)}(0)| < \infty$ used in Proposition 1 is natural in view of the desired regularity properties of $\sum_{n \geq 1} R_n(\cdot)$ in Hypothesis $\mathcal{R}(m)$ -(i).

For $k \in \mathbb{N}^*$, we denote by \mathcal{H}_k the space of all complex-valued continuous Lebesgue-integrable functions on \mathbb{R}^d , whose Fourier transform is compactly supported and k -times continuously differentiable on \mathbb{R}^d . The constant C_d is defined in (2).

Theorem 1 Assume that $d \geq 3$, that Hypothesis $\mathcal{R}(m)$ holds with $m = \max\{d - 2, 2 + \varepsilon\}$ for some $\varepsilon > 0$, that $\nabla \lambda(0) = 0$, and that Σ is positive definite. Then we have for all $g \in \mathcal{H}_{d-2}$:

$$\sum_{n=1}^{+\infty} \mathbb{E}[f(X_n) g(S_n - a)] \sim \frac{C_d L(0)}{\langle \Sigma^{-1} a, a \rangle^{\frac{d-2}{2}}} \mathcal{L}_d(g) \quad \text{when } \|a\| \rightarrow +\infty \quad (a \in \mathbb{R}^d). \quad (4)$$

It is well-known (see e.g. [3]) that the conclusion of Theorem 1 implies that, for all $a \in \mathbb{R}^d$, $U_a(B) := \sum_{n=1}^{+\infty} \mathbb{E}[f(X_n) 1_B(S_n - a)]$ ($B \in \mathcal{B}(\mathbb{R}^d)$) defines a positive Radon measure on \mathbb{R}^d , and that $\langle \Sigma^{-1} a, a \rangle^{d-2/2} U_a$ weakly converges to $C_d L(0) \mathcal{L}_d(\cdot)$ when $\|a\| \rightarrow +\infty$. Consequently:

Corollary 1 Under the assumptions of Theorem 1, the property (4) is fulfilled with any real-valued continuous compactly supported function g on \mathbb{R}^d , and also with $g = 1_B$ for any bounded Borel set B in \mathbb{R}^d whose boundary has a zero Lebesgue-measure.

The proof of Theorem 1 is based on some Fourier techniques partially derived from [1]. In the case $d = 3$ or 4, the optimal order 2 of the i.i.d. case (see Section 1) is here replaced with the order $2 + \varepsilon$. This will be needed in the proof of Lemma 2 below.

Proof of Theorem 1. Set $E_n(t) := \mathbb{E}[f(X_n) e^{i\langle t, S_n \rangle}]$ for each $t \in \mathbb{R}^d$. Let $h \in \mathcal{H}_{d-2}$, and let $b > 0$ be such that $\hat{h}(t) = 0$ if $\|t\| > b$. The inverse Fourier formula on h easily gives

$$\forall n \in \mathbb{N}^*, \quad (2\pi)^d \mathbb{E}[f(X_n) h(S_n)] = \int_{\|t\| \leq b} \hat{h}(t) E_n(t) dt.$$

Since $\lambda(t) = 1 - \frac{1}{2} \langle \Sigma t, t \rangle + o(\|t\|^2)$, one can choose $\alpha < R$ such that

$$\|t\| \leq \alpha \Rightarrow |\lambda(t)| \leq 1 - \frac{1}{4} \langle \Sigma t, t \rangle. \quad (5)$$

Now let $r \in (0, \alpha)$, and let χ be a real-valued compactly supported and indefinitely differentiable function on \mathbb{R}^d , such that its support is contained in $\{t \in \mathbb{R}^d : \|t\| \leq \alpha\}$ and $\chi(t) = 1$ for $\|t\| \leq r$. Let us write

$$(2\pi)^d \mathbb{E}[f(X_n) h(S_n)] = \int_{\|t\| \leq \alpha} \chi(t) \hat{h}(t) E_n(t) dt + \int_{r < \|t\| \leq b} (1 - \chi(t)) \hat{h}(t) E_n(t) dt.$$

Let $a \in \mathbb{R}^d$ and set $h_a(\cdot) := h(\cdot - a)$. We have $\widehat{h_a}(t) = \hat{h}(t) e^{-i\langle t, a \rangle}$. By applying the previous equalities to h_a and using Hypothesis $\mathcal{R}(m)$, one obtains

$$(2\pi)^d \sum_{n=1}^{+\infty} \mathbb{E}_\mu[f(X_n) h(S_n - a)] = I(a) + J(a) + K(a)$$

$$\begin{aligned} \text{with } I(a) &= \int_{\|t\| \leq \alpha} \chi(t) \hat{h}(t) \frac{\lambda(t)}{1 - \lambda(t)} L(t) e^{-i\langle t, a \rangle} dt \\ J(a) &= \int_{\|t\| \leq \alpha} \chi(t) \hat{h}(t) \left(\sum_{n \geq 1} R_n(t) \right) e^{-i\langle t, a \rangle} dt \\ K(a) &= \int_{r < \|t\| \leq b} (1 - \chi(t)) \hat{h}(t) \left(\sum_{n \geq 1} E_n(t) \right) e^{-i\langle t, a \rangle} dt. \end{aligned}$$

Indeed, it follows from (5) that $\forall t \in B(0, \alpha)$, $t \neq 0$, we have $|\lambda(t)| < 1$ and $\sum_{n \geq 1} |\lambda(t)|^n = \frac{|\lambda(t)|}{1 - |\lambda(t)|} \leq \frac{4}{\langle \Sigma t, t \rangle}$. Since $d \geq 3$ and Σ is invertible, the function $t \mapsto \frac{1}{\langle \Sigma t, t \rangle}$ is integrable near $t = 0$. So the term $I(a)$ derives from Lebesgue's theorem. The terms $J(a)$ and $K(a)$ follow from the uniform convergence of the series $\sum_{n \geq 1} R_n(\cdot)$ and $\sum_{n \geq 1} E_n(\cdot)$ on $B(0, \alpha)$ and $K_{r,b}$ respectively. Theorem 1 is then a consequence of the three next lemmas. \square

Remark 1 *The uniform convergence stated in Hypothesis $\mathcal{R}(m)$ for the series $\sum_{n \geq 1} R_n(\cdot)$ and $\sum_{n \geq 1} E_n(\cdot)$ is used for defining $J(a)$ and $K(a)$. Of course, alternative conditions may be used, as for instance: $\sum_{n \geq 1} \int_{B(0,R)} |R_n(t)| dt < \infty$ and $\sum_{n \geq 1} \int_{K_{r,r'}} |E_n(t)| dt < \infty$.*

Lemma 1 *We have $J(a) + K(a) = o(\|a\|^{-(d-2)})$ when $\|a\| \rightarrow +\infty$.*

Proof. By Hypothesis $\mathcal{R}(m)$, the integrands in $J(a)$ and $K(a)$ are respectively in $\mathcal{C}_b^m(B(0, R), \mathbb{C})$ (with $R > \alpha$) and in $\mathcal{C}_b^m(K_{r/2, 2b}, \mathbb{C})$. Since $m = \max\{d - 2, 2 + \varepsilon\} \geq d - 2$, this gives the desired result. \square

Now let us investigate $I(a)$. An easy computation yields $I(a) = I_1(a) + I_2(a) + I_3(a)$ with

$$\begin{aligned} I_1(a) &:= \int_{\|t\| \leq \alpha} \chi(t) \frac{\hat{h}(t)\lambda(t)L(t) - \hat{h}(0)L(0)}{1 - \lambda(t)} e^{-i\langle t, a \rangle} dt \\ I_2(a) &:= 2\hat{h}(0)L(0) \int_{\|t\| \leq \alpha} \frac{\chi(t)}{\langle \Sigma t, t \rangle} e^{-i\langle t, a \rangle} dt \\ I_3(a) &:= 2\hat{h}(0)L(0) \int_{\|t\| \leq \alpha} \chi(t) \frac{\lambda(t) - 1 + \frac{1}{2}\langle \Sigma t, t \rangle}{(1 - \lambda(t))\langle \Sigma t, t \rangle} e^{-i\langle t, a \rangle} dt. \end{aligned}$$

Lemma 2 *We have $I_1(a) + I_3(a) = o(\|a\|^{-(d-2)})$ when $\|a\| \rightarrow +\infty$.*

Proof. To study $I_1(a)$, let us define the following function on \mathbb{R}^d :

$$\theta_1(t) = \chi(t) \left(\hat{h}(t)\lambda(t)L(t) - \hat{h}(0)L(0) \right).$$

Then θ_1 has $d - 2$ continuous derivatives on \mathbb{R}^d and since $\theta_1(0) = 0$, we have $|\theta_1(t)| = O(\|t\|)$ on \mathbb{R}^d . Now set $u(t) = \frac{\theta_1(t)}{1 - \lambda(t)}$. Some standard derivative arguments give (see Remark (a) in Appendix A.0)

$$\forall i = 0, \dots, d - 2, \quad |u^{(i)}(t)| = O(\|t\|^{-(1+i)}), \quad (6)$$

where $u^{(i)}(\cdot)$ stands for any partial derivative of order i of u . In particular the partial derivatives of order $d - 2$ of u are Lebesgue-integrable on \mathbb{R}^d . Since $I_1(a) = \hat{u}(a)$, the claimed property for $I_1(a)$ follows from Proposition A.1 in Appendix A.1. The same property can be similarly established for $I_3(a)$ by defining the function

$$\theta_3(t) = \chi(t) \left(\lambda(t) - 1 + \frac{1}{2}\langle \Sigma t, t \rangle \right).$$

Indeed, if $d \geq 5$, then θ_3 has three continuous derivatives on \mathbb{R}^d and it can be easily seen that $|\theta_3(t)| = O(\|t\|^3)$, $|\theta_3^{(1)}(t)| = O(\|t\|^2)$ and $|\theta_3^{(2)}(t)| = O(\|t\|)$. Let $v(t) = \frac{\theta_3(t)}{(1 - \lambda(t))\langle \Sigma t, t \rangle}$. Remark (b) in Appendix A.0 then yields

$$\forall i = 0, \dots, d - 2, \quad |v^{(i)}(t)| = O(\|t\|^{-(1+i)}). \quad (7)$$

If $d = 3$ or 4 , then we only have $|\theta_3(t)| = O(\|t\|^{2+\varepsilon})$, $|\theta_3^{(1)}(t)| = O(\|t\|^{1+\varepsilon})$ and $|\theta_3^{(2)}(t)| = O(\|t\|^\varepsilon)$. Similarly, one gets

$$\forall i = 0, \dots, d - 2, \quad |v^{(i)}(t)| = O(\|t\|^{-(i+2-\varepsilon)}). \quad (8)$$

From either (7) or (8), the desired estimate on $I_3(a) = 2\hat{h}(0)L(0)\hat{v}(a)$ again follows from Proposition A.1. \square

Lemma 3 *Set $C'_d = (2\pi)^{\frac{d}{2}} 2^{\frac{d}{2}-1} \Gamma\left(\frac{d-2}{2}\right) (\det \Sigma)^{-\frac{1}{2}}$. We have $I_2(a) \sim \frac{C'_d \hat{h}(0)L(0)}{\langle \Sigma^{-1}a, a \rangle^{\frac{d-2}{2}}}$ when $\|a\| \rightarrow +\infty$.*

Proof. Let $S(\mathbb{R}^d)$ denote the so-called Schwartz space. Since $\chi \in S(\mathbb{R}^d)$ and the Fourier transform is a bijection on $S(\mathbb{R}^d)$, let us call $\psi \in S(\mathbb{R}^d)$ such that $\widehat{\psi} = \chi$. Then, setting $\psi_a(\cdot) := \psi(\cdot - a)$, we have

$$I_2(a) = 2\hat{h}(0) L(0) \int_{\mathbb{R}^d} \frac{\widehat{\psi_a}(t)}{\langle \Sigma t, t \rangle} dt.$$

Set $\Delta = \text{diag}(\sqrt{\lambda_1}, \dots, \sqrt{\lambda_d})$, where the λ_i 's are the eigenvalues of the covariance matrix Σ ($\lambda_i > 0$ because Σ is invertible), and let P be any orthogonal matrix of order d such that

$$P^{-1}\Sigma P = \Delta^2 = \text{diag}(\lambda_1, \dots, \lambda_d).$$

We have $\langle \Sigma t, t \rangle = \langle \Delta^2 P^{-1}t, P^{-1}t \rangle = \|\Delta P^{-1}t\|^2$. So, by setting $t = P\Delta^{-1}u$, one gets

$$I_2(a) = 2\hat{h}(0) L(0) \int (\det \Delta)^{-1} \widehat{\psi_a}(P\Delta^{-1}u) \frac{1}{\|u\|^2} du.$$

The Fourier transform of $g(v) = \psi(P\Delta v - a)$ is $\hat{g}(u) = (\det \Delta)^{-1} \widehat{\psi_a}(P\Delta^{-1}u)$. So we have

$$I_2(a) = 2\hat{h}(0) L(0) c \int \psi(P\Delta v - a) \frac{1}{\|v\|^{d-2}} dv, \quad (9)$$

where $c = (2\pi)^{\frac{d}{2}} 2^{\frac{d}{2}-2} \Gamma(\frac{d-2}{2})$. Indeed the equality (9) follows from the fact that we have $(\|\cdot\|^{-2})(v) = c\|v\|^{2-d}$ in the sense of tempered distribution on \mathbb{R}^d . (See also Appendix A.2 for an elementary proof of (9)). Now set $b = \Delta^{-1}P^{-1}a$, $F(x) = \psi(-P\Delta x)$. Furthermore, set $\beta = \|b\|$, $b = \beta\tilde{b}$, so $\|\tilde{b}\| = 1$, and define $F_\beta(x) = \beta^d F(\beta x)$ for any $\beta > 0$. Then

$$I_2(a) = 2\hat{h}(0) L(0) c \int F(b-v) \frac{1}{\|v\|^{d-2}} dv = 2\hat{h}(0) L(0) c \int F_\beta(\tilde{b}-w) \frac{1}{\|\beta w\|^{d-2}} dw,$$

so that $\beta^{d-2} I_2(a) = 2c\hat{h}(0) L(0) (F_\beta * f_{d-2})(\tilde{b})$, where $f_{d-2}(w) := \frac{1}{\|w\|^{d-2}}$, and $*$ denotes the convolution product on \mathbb{R}^d . By using some standard arguments of approximate identity, one can prove (see Appendix A.3.) that the following convergence holds uniformly in $\tilde{b} \in \mathbb{R}^d$ such that $\|\tilde{b}\| = 1$:

$$\lim_{\beta \rightarrow +\infty} (F_\beta * f_{d-2})(\tilde{b}) = \int_{\mathbb{R}^d} F(w) dw. \quad (10)$$

We have $\int F(w) dw = (\det \Delta)^{-1} \int \psi(y) dy = (\det \Sigma)^{-\frac{1}{2}}$, because $\int \psi(y) dy = \hat{\psi}(0) = \chi(0) = 1$, and $\beta = \|Pb\| = \|\Sigma^{-\frac{1}{2}}a\| = \langle \Sigma^{-1}a, a \rangle^{\frac{1}{2}}$. The above computations then yield Lemma 3. \square

3 Operator-type procedure in Markov models

In this section $(X_n, S_n)_{n \geq 0}$ still denotes a sequence of random variables taking values in $E \times \mathbb{R}^d$, but from now on $(X_n)_{n \geq 0}$ is assumed to be a Markov chain with state space E , transition probability $Q(x, dy)$, and initial distribution μ . We suppose that $S_0 = 0$. The functional action of $Q(x, dy)$ is given by $Qf(x) = \int_E f(y) Q(x, dy)$ provided that this integral is well defined. In this Markov context, we first present a general assumption providing an operator-type formula for the term $\mathbb{E}[f(X_n) e^{i\langle t, S_n \rangle}]$ of Hypothesis $\mathcal{R}(m)$. Next, we briefly compare the spectral method developed in [1] with that presented in [17].

3.1 A Markov context

For the moment we make the following abuse of notation: given $x \in E$, we denote by \mathbb{P}_x the underlying probability measure (and by \mathbb{E}_x the associated expectation) to refer to the case when the initial distribution of $(X_n)_{n \geq 0}$, is the Dirac distribution at x . This notation takes a precise (and usual) sense in the example below and in the setting of Section 4. For any bounded measurable function f on E , and for $n \in \mathbb{N}$, $t \in \mathbb{R}^d$, let us define:

$$(Q_n(t)f)(x) := \mathbb{E}_x[e^{i \langle t, S_n \rangle} f(X_n)] \quad (x \in E). \quad (11)$$

Let us observe that $Q_1(0) = Q$. Let us consider the following condition:

Condition (G). For all $t \in \mathbb{R}^d$, $(Q_n(t))_{n \in \mathbb{N}}$ is a semi-group, that is:

$$\forall (m, n) \in \mathbb{N}^2, \quad Q_{m+n}(t) = Q_m(t) \circ Q_n(t).$$

Under Condition (G), we have in particular $Q_n(t) = Q(t)^n$ where $Q(t) := Q_1(t)$ is defined by:

$$(Q(t)f)(x) := \mathbb{E}_x[e^{i \langle t, S_1 \rangle} f(X_1)] \quad (x \in E). \quad (12)$$

The $Q(t)$'s are called the Fourier operators, and in view of the study of Hypothesis $\mathcal{R}(m)$, one can see that (11) provides the following interesting formula :

$$\mathbb{E}_x[e^{i \langle t, S_n \rangle} f(X_n)] = (Q(t)^n f)(x) \quad (x \in E). \quad (13)$$

Example: the Markov random walks.

If $(X_n, S_n)_{n \in \mathbb{N}}$ is a Markov chain with state space $E \times \mathbb{R}^d$ and transition probability P satisfying the following property

$$\forall (x, a) \in E \times \mathbb{R}^d, \quad \forall A \in \mathcal{E}, \quad \forall B \in B(\mathbb{R}^d), \quad P((x, a), A \times B) = P((x, 0), A \times (B - a)),$$

the sequence $(X_n, S_n)_{n \in \mathbb{N}}$ is called a Markov random walk (MRW). Of course $(X_n)_{n \in \mathbb{N}}$ is then also a Markov chain, called the driving Markov chain of the MRW. We still assume that $S_0 = 0$. The previous translation property is equivalent to the following one: for any bounded measurable function F on $E \times \mathbb{R}^d$ and for all $a \in \mathbb{R}^d$, we have $(PF)_a = P(F_a)$, where we set for any function $G : E \times \mathbb{R}^d \rightarrow \mathbb{C}$: $G_a(x, b) := G(x, b + a)$. Let us check Condition (G). Let $t \in \mathbb{R}^d$ (fixed). Given $f : E \rightarrow \mathbb{R}$ bounded and measurable, we set: $F(x, b) := f(x) e^{i \langle t, b \rangle}$ for $x \in E$ and $b \in \mathbb{R}^d$. We have:

$$\begin{aligned} (Q_n(t)f)(x) &:= \mathbb{E}_{(x,0)}[e^{i \langle t, S_n \rangle} f(X_n)] &= \mathbb{E}_{(x,0)}[(PF)(X_{n-1}, S_{n-1})] \\ & &= \mathbb{E}_{(x,0)}[(PF)_{S_{n-1}}(X_{n-1}, 0)] \\ & &= \mathbb{E}_{(x,0)}[(PF_{S_{n-1}})(X_{n-1}, 0)] \\ & &= \mathbb{E}_{(x,0)}[e^{i \langle t, S_{n-1} \rangle} (PF)(X_{n-1}, 0)], \end{aligned}$$

(for the last equality, use: $F_a(y, b) = f(y) e^{i \langle t, b+a \rangle} = e^{i \langle t, a \rangle} F(y, b)$). Therefore we have: $(Q_n(t)f)(x) = \mathbb{E}_{(x,0)}[e^{i \langle t, S_{n-1} \rangle} g(X_{n-1})] = (Q_{n-1}(t)g)(x)$ with

$$g(\cdot) := (PF)(\cdot, 0) = \mathbb{E}_{(\cdot,0)}[f(X_1) e^{i \langle t, S_1 \rangle}] = (Q_1(t)f)(\cdot).$$

We have proved that: $(Q_n(t)f)(x) = (Q_{n-1}(t) \circ Q_1(t)f)(x)$, hence Condition (G) is fulfilled.

As a classical example of Markov random walk, recall that, if $(S_n)_{n \in \mathbb{N}}$ is a d -dimensional additive functional of a Markov chain $(X_n)_{n \in \mathbb{N}}$, then $(X_n, S_n)_{n \in \mathbb{N}}$ is a MRW.

3.2 Spectral methods

When $(X_n, S_n)_{n \geq 0}$ satisfies Condition (\mathcal{G}) , the spectral method consists in investigating the regularity conditions of Hypothesis $\mathcal{R}(m)$ via Formula (13). Let us outline the usual spectral method developed in [1] and the key idea of the weak spectral method presented in [17]. Given Banach spaces \mathcal{B} and X , we denote by $\mathcal{L}(\mathcal{B}, X)$ the space of the bounded linear operators from \mathcal{B} to X , with the simplifications $\mathcal{L}(\mathcal{B}) = \mathcal{L}(\mathcal{B}, \mathcal{B})$ and $\mathcal{B}' = \mathcal{L}(\mathcal{B}, \mathbb{C})$.

(I) *The spectral method via the standard perturbation theorem.*

Let $(\mathcal{B}, \|\cdot\|_{\mathcal{B}})$ be a Banach space of \mathbb{C} -valued measurable functions on E . The operator norm in $\mathcal{L}(\mathcal{B})$ is also denoted by $\|\cdot\|_{\mathcal{B}}$. Let us assume that $(X_n)_{n \geq 0}$ possesses a stationary distribution π defining a continuous linear form on \mathcal{B} . In [1], the assumptions are the following ones: for each $t \in \mathbb{R}^d$, the Fourier operator $Q(t)$ defined in (12) is in $\mathcal{L}(\mathcal{B})$, and

- (A) Q is strongly ergodic on \mathcal{B} , namely : $\lim_n Q^n = \pi$ in $\mathcal{L}(\mathcal{B})$,
- (B) For any compact set $K \subset \mathbb{R}^d \setminus \{0\}$: $\exists \rho < 1$, $\sup_{t \in K} \|Q(t)^n\|_{\mathcal{B}} = O(\rho^n)$,
- (C) $Q(\cdot) \in \mathcal{C}_b^m(\mathbb{R}^d, \mathcal{L}(\mathcal{B}))$ for some $m \in \mathbb{N}$.

Let $\rho_0 \in (\rho, 1)$. From (B), $(z - Q(t))^{-1}$ is defined for all $t \in K$ and $z \in \mathbb{C}$ such that $|z| = \rho_0$. By (C), we have: $(z - Q(\cdot))^{-1} \in \mathcal{C}_b^m(K, \mathcal{L}(\mathcal{B}))$, and from $Q(t)^n = \frac{1}{2i\pi} \oint_{|z|=\rho_0} z^n (z - Q(t))^{-1} dz$, Hypothesis $\mathcal{R}(m)$ -*(ii)* can be then deduced from (13) when $f \in \mathcal{B}$. Next, using (A) (C) and the standard perturbation theorem, one can prove that, for $t \in B := B(0, R)$ for some $R > 0$, $Q(t)^n = \lambda(t)^n \Pi(t) + N(t)^n$ where $\lambda(t)$ is the perturbed eigenvalue near $\lambda = 1$, $\Pi(t)$ is the corresponding rank-one eigenprojection, and $N(t) \in \mathcal{L}(\mathcal{B})$ satisfies $\sup_{t \in B} \|N(t)^n\|_{\mathcal{B}} = O(\kappa^n)$ for some $\kappa \in (0, 1)$. Since the previous eigen-elements inherit the regularity of $Q(\cdot)$, Hypothesis $\mathcal{R}(m)$ -*(i)* is fulfilled.

As already mentioned in Section 1, Condition (C) requires strong assumptions on Q and S_1 . In fact, by deriving formally (12) in the variable t (in case $d = 1$ and at $t = 0$ to simplify), one gets : $(Q^{(m)}(0)f)(x) := i^m \mathbb{E}_x[S_1^m f(X_1)]$. Therefore a necessary condition for (C) to be true is that $x \mapsto \mathbb{E}_x[S_1^m f(X_1)] \in \mathcal{B}$ for all $f \in \mathcal{B}$. This condition may be clearly quite restrictive or even not satisfied when S_1 is unbounded.

(II) *A weak spectral method via the Keller-Liverani perturbation theorem.*

The following alternative assumptions are proposed in [17]: (A) (B) hold true with respect to a whole family $\{\mathcal{B}_\theta, \theta \in I\}$ of Banach spaces (instead of a single one as in (I)), and (C) is replaced by the following condition:

(C') On each space \mathcal{B}_θ , $Q(\cdot)$ satisfies the Keller-Liverani perturbation theorem [18] near $t = 0$, and $Q(\cdot) \in \mathcal{C}_b^k(\mathbb{R}^d, \mathcal{L}(\mathcal{B}_\theta, \mathcal{B}_{\theta'}))$ for suitable $\mathcal{B}_\theta \subset \mathcal{B}_{\theta'}$ ($k = 0, \dots, m$).

The above regularity hypothesis will involve (in substance) that we have, for some suitable $b, c \in I$, the following condition: $\forall f \in \mathcal{B}_b, x \mapsto \mathbb{E}_x[S_1^m f(X_1)] \in \mathcal{B}_c$ (with $\mathcal{B}_b \subset \mathcal{B}_c$). Thanks to the "gap" between \mathcal{B}_b and \mathcal{B}_c , this condition is less restricting than in (I). Of course, the passage from (C') to the regularity properties of the maps $(z - Q(\cdot))^{-1}$ is here more difficult than in the method (I) because $(z - Q(t))^{-1}$ must be seen as elements of $\mathcal{L}(\mathcal{B}_\theta, \mathcal{B}_{\theta'})$ according to a procedure involving several suitable values $(\theta, \theta') \in I^2$. Such results have been presented in [10, 17, 9].

4 Application to three classical Markov models

In this section, the weak spectral method is applied to the three classical models cited in Section 1: the v -geometrically ergodic Markov chains, the ρ -mixing Markov chains, and the iterative Lipschitz models. In order to use directly some technical results of [17], we consider Markov random walks of the form $S_n = \sum_{k=1}^n \xi(X_k)$, where $\xi : E \rightarrow \mathbb{R}^d$ is a centered functional. We establish below that, in the three above models, the property (1) holds under the nonlattice condition and under some moment conditions on ξ of order $m_d + \varepsilon$, where m_d has been defined in Section 1 by $m_d = \max\{d - 2, 2\}$. First, let us give some common definitions and preliminary properties that will be used in the three models.

Let $(X_n)_{n \geq 0}$ be a Markov chain with state space (E, \mathcal{E}) , transition probability $Q(x, dy)$, stationary distribution π , and initial distribution μ . Let $\xi = (\xi_1, \dots, \xi_d)$ be a \mathbb{R}^d -valued measurable functional on E , and let us define the following associated random walk:

$$S_n = \sum_{k=1}^n \xi(X_k).$$

Throughout this section, we assume that $d \geq 3$, and that ξ is π -centered (i.e. ξ is π -integrable and $\pi(\xi_i) = 0$ for $i = 1, \dots, d$).

Obviously $(X_n, S_n)_{n \in \mathbb{N}}$ is a special case of Markov random walk (with $(X_n)_{n \geq 0}$ as driving chain), and the Fourier operators $Q(t)$ in (12) are here given by the kernels

$$Q(t)(x, dy) = e^{i\langle t, \xi(y) \rangle} Q(x, dy). \quad (14)$$

All the Banach spaces (say $(\mathcal{B}, \|\cdot\|_{\mathcal{B}})$) used in this section are either composed of complex-valued π -integrable functions on E , or composed of classes (modulo π) of such functions. In addition \mathcal{B} contains 1_E (or $Cl(1_E)$), and π always defines a continuous linear form on \mathcal{B} (i.e. $\pi \in \mathcal{B}'$), so that the following rank-one projection can be defined on \mathcal{B} :

$$\Pi f = \pi(f)1_E \quad (f \in \mathcal{B}). \quad (15)$$

We will say that $(X_n)_{n \geq 0}$ is strongly ergodic on \mathcal{B} if Q continuously acts on \mathcal{B} (i.e. $Q \in \mathcal{L}(\mathcal{B})$) and satisfies the following property:

$$\exists \kappa_0 < 1, \exists C > 0, \forall n \geq 1, \quad \|Q^n - \Pi\|_{\mathcal{B}} \leq C \kappa_0^n, \quad (16)$$

where $\|\cdot\|_{\mathcal{B}}$ denotes the operator norm on \mathcal{B} . Condition (16) is equivalent to that already mentioned in Subsection 3.2 : $\lim_n \|Q^n - \Pi\|_{\mathcal{B}} = 0$. We will repeatedly use the following consequence of Formula (13):

$$f \in \mathcal{B}, \mu \in \mathcal{B}' \Rightarrow \forall n \geq 1, \forall t \in \mathbb{R}^d, \quad \mathbb{E}_{\mu}[f(X_n) e^{i\langle t, S_n \rangle}] = \mu(Q(t)^n f). \quad (17)$$

All the next conditions on the initial distribution μ are satisfied with $\mu = \pi$, and in Subsections 4.1 and 4.3 when μ is the Dirac distribution at any $x \in E$.

It will be seen that the assumptions of the next corollaries always imply that the following asymptotic covariance matrix is well-defined:

$$\Sigma = \lim_n \frac{1}{n} \mathbb{E}_{\pi}[S_n S_n^*]. \quad (18)$$

Now let us introduce the following classical (Markov) nonlattice condition:

Nonlattice Condition : (Q, ξ) , or merely ξ , is said nonlattice if there exist no $b \in \mathbb{R}^d$, no closed subgroup H in \mathbb{R}^d , $H \neq \mathbb{R}^d$, no π -full Q -absorbing set $A \in \mathcal{E}$, and finally no bounded measurable function $\theta : E \rightarrow \mathbb{R}^d$ such that

$$\forall x \in A, \quad \xi(y) + \theta(y) - \theta(x) \in b + H \quad Q(x, dy) - a.s.$$

As usual, $A \in \mathcal{E}$ is said to be π -full if $\pi(A) = 1$, and Q -absorbing if $Q(a, A) = 1$ for all $a \in A$. Finally each partial derivative $\frac{\partial^k Q}{\partial t_{p_1} \dots \partial t_{p_k}}(t)$ of the Fourier kernels $Q(t)$ (see (14)) is defined by means of the kernel

$$Q_{(p_1, \dots, p_k)}(t)(x, dy) = i^k \left(\prod_{s=1}^k \xi_{p_s}(y) \right) e^{i\langle t, \xi(y) \rangle} Q(x, dy). \quad (19)$$

4.1 Applications to the v -geometrically ergodic Markov chains.

We suppose that \mathcal{E} is countably generated. Let $v : E \rightarrow [1, +\infty)$ be some fixed unbounded function, and let us assume that $(X_n)_{n \geq 0}$ is v -geometrically ergodic, namely : we have (16) on the weighted supremum-normed space $(\mathcal{B}_v, \|\cdot\|_v)$ of all measurable functions $f : E \rightarrow \mathbb{C}$ satisfying the condition :

$$\|f\|_v = \sup_{x \in E} \frac{|f(x)|}{v(x)} < \infty.$$

Notice that the previous assumption involves $\pi(v) < \infty$. The v -geometrical ergodicity condition can be investigated with the help of the so-called drift conditions. For this fact, and for the classical examples of such models, we refer to [20].

The method (II) outlined in Subsection 3.2 is applied here with the following spaces : for $0 < \theta \leq 1$, we denote by $(\mathcal{B}_\theta, \|\cdot\|_\theta)$ the weighted supremum-normed space associated to v^θ , where we set

$$\|f\|_\theta = \sup_{x \in E} \frac{|f(x)|}{v(x)^\theta}.$$

Corollary 2 *Assume that $\mu(v) < \infty$ and that $\xi : E \rightarrow \mathbb{R}^d$ ($d \geq 3$) is a π -centered nonlattice functional such that we have with $m_d = \max\{d - 2, 2\}$:*

$$\exists C > 0, \exists \delta_0 > 0, \quad \|\xi\|^{m_d + \delta_0} \leq C v. \quad (20)$$

Let $f \in \mathcal{B}_\theta$, $f \geq 0$, where ϑ is such that $0 < \vartheta < 1 - \frac{m}{m_d + \delta_0}$ with $m \in (2, 2 + \delta_0)$ if $d = 3, 4$, and $m = d - 2$ if $d \geq 5$.

Then we have (4) with $L(0) = \pi(f)$ and Σ defined by (18).

In order to present a general study of Hypothesis $\mathcal{R}(m)$ in the present context, we consider below any real numbers m, η such that $0 < m < \eta$, we set $\tau = m - [m]$, where $[m]$ denotes the integer part of m , and we assume that $\xi : E \rightarrow \mathbb{R}^d$ is a measurable function such that:

$$\exists C > 0, \quad \|\xi\|^\eta \leq C v. \quad (21)$$

Thanks to Theorem 1, Corollary 2 follows from the next Propositions 2-3 (applied with $\eta = m_d + \delta_0$) and from Remark 2.

Proposition 2 *Let ϑ_m be such that $0 < \vartheta_m < \vartheta_m + \frac{m}{\eta} < 1$. Assume that ξ satisfies Condition (21), that $\mu(v) < \infty$, and that $f \in \mathcal{B}_{\vartheta_m}$. Then Hypothesis $\mathcal{R}(m)$ -(i) holds true with $L(0) = \pi(f)$.*

Proof. For any $\kappa \in (0, 1)$, let us define

$$\mathcal{D}_\kappa = \{z : z \in \mathbb{C}, |z| \geq \kappa, |z - 1| \geq (1 - \kappa)/2\}. \quad (22)$$

Lemma 4 *Let $\theta \in (0, 1]$. Then Property (16) is satisfied on \mathcal{B}_θ . Moreover, there exists $\kappa_\theta \in (0, 1)$ such that, for all $\kappa \in (\kappa_\theta, 1)$, there exists $R_\kappa > 0$ such that we have:*

$$\begin{aligned} \forall z \in \mathcal{D}_\kappa, \forall t \in B(0, R_\kappa), \quad (z - Q(t))^{-1} \in \mathcal{L}(\mathcal{B}_\theta), \\ \sup \{ \|(z - Q(t))^{-1}\|_\theta : z \in \mathcal{D}_\kappa, t \in B(0, R_\kappa) \} < \infty. \end{aligned}$$

Proof. By hypothesis (v -geometrical ergodicity), we have (16) on \mathcal{B}_1 . The fact that (16) extends to \mathcal{B}_θ is a well-known consequence of the so-called drift conditions and Jensen's inequality, see e.g. [17, Lem. 10.1]. The second assertion follows from the Keller-liverani perturbation theorem [18], see [17, Lem. 10.1 and § 4]. \square

Now let $\kappa \in (\kappa_{\vartheta_m}, 1)$. Let $\Gamma_{0,\kappa}$ be the oriented circle centered at $z = 0$, with radius κ , and let $\Gamma_{1,\kappa}$ be the oriented circle centered at $z = 1$, with radius $\frac{1-\kappa}{2}$. From (16) and the Keller-liverani theorem (both applied on $\mathcal{B}_{\vartheta_m}$), we can write for all $t \in B(0, R_\kappa)$ the following equality in $\mathcal{L}(\mathcal{B}_{\vartheta_m})$ (e.g. see [17, Sec. 7]):

$$Q(t)^n = \lambda(t)^n \Pi(t) + N(t)^n \quad (23)$$

where $\lambda(t)$ is the perturbed eigenvalue near $\lambda(0) = 1$, and $\Pi(t), N(t)^n$ can be defined by the following line integrals:

$$\Pi(t) = \frac{1}{2i\pi} \oint_{\Gamma_{1,\kappa}} (z - Q(t))^{-1} dz \quad \text{and} \quad N(t)^n = \frac{1}{2i\pi} \oint_{\Gamma_{0,\kappa}} z^n (z - Q(t))^{-1} dz. \quad (24)$$

For any probability measure ν on E such that $\nu(v) < \infty$ and for any $g \in \mathcal{B}_{\vartheta_m}$, we define

$$L_{\nu,g}(t) := \nu(\Pi(t)g) = \frac{1}{2i\pi} \oint_{\Gamma_{1,\kappa}} \nu((z - Q(t))^{-1}g) dz, \quad (25)$$

$$R_{n,\nu,g}(t) := \nu(N(t)^n g) = \frac{1}{2i\pi} \oint_{\Gamma_{0,\kappa}} z^n \nu((z - Q(t))^{-1}g) dz. \quad (26)$$

The previous quantities are well defined since $g \in \mathcal{B}_{\vartheta_m}$ and $\nu \in \mathcal{B}'_1 \subset \mathcal{B}'_{\vartheta_m}$. From (17) we have (3) with $\forall t \in B(0, R_\kappa)$:

$$L(t) = L_{\mu,f}(t) \quad \text{and} \quad R_n(t) = R_{n,\mu,f}(t). \quad (27)$$

In particular we have $L(0) = \mu(\Pi f) = \pi(f)$. Besides it can be easily proved (see [17, Sect.7] for details) that we have with possibly R_κ reduced:

$$\forall t \in B(0, R_\kappa), \lambda(t) = \frac{\pi(Q(t)1_E) - R_{1,\pi,1_E}(t)}{L_{\pi,1_E}(t)}. \quad (28)$$

Finally, from Lemma 4 and (24), there exists $C > 0$ such that: $\forall t \in B(0, R_\kappa)$, $R_n(t) \leq C \kappa^n$. Therefore, the series $\sum_{n \geq 1} R_n(t)$ converge, and we have

$$\sum_{n \geq 1} R_n(t) = \frac{1}{2i\pi} \oint_{\Gamma_{0,\kappa}} \frac{z}{1-z} \mu((z - Q(t))^{-1} f) dz. \quad (29)$$

From (27) (28) (29), the desired regularity properties in Hypothesis $\mathcal{R}(m)$ -(i) will be fulfilled if we prove that $\pi(Q(\cdot)1_E)$, $\pi((z - Q(\cdot))^{-1}1_E)$, and $\mu((z - Q(\cdot))^{-1}f)$ are in $\mathcal{C}_b^m(B(0, R), \mathbb{C})$ for some $R > 0$ and have uniform bounded derivatives in $z \in \mathcal{D}_\kappa$ (thus in $z \in \Gamma_{0,\kappa} \cup \Gamma_{1,\kappa}$). Let Q_k denote any kernel of the form $Q_{(p_1, \dots, p_k)}$ defined in (19).

Lemma 5 *Let $0 < \theta < \theta' \leq 1$, let $k \in \mathbb{N}$ such that $0 \leq k \leq \lfloor m \rfloor$, let $R > 0$, and set $B = B(0, R)$. Then we have the following properties :*

- (i) *If $\tau' \in (0, 1)$ and $\theta + \frac{k+\tau'}{\eta} \leq \theta'$, then $Q_k \in \mathcal{C}_b^{\tau'}(B, \mathcal{L}(\mathcal{B}_\theta, \mathcal{B}_{\theta'}))$,*
- (ii) *If $k \leq \lfloor m \rfloor - 1$ and $\theta + \frac{k+1}{\eta} < \theta'$, then $Q_k \in \mathcal{C}_b^1(B, \mathcal{L}(\mathcal{B}_\theta, \mathcal{B}_{\theta'}))$.*

Proof. Lemma 5 follows from (21): this is established in [17, Lem. 10.4-5]. \square

To make easier the use of Lemma 5, let us define for any fixed $u > 0$: $T_u(\theta) := \theta + u/\eta$. Since $m = \lfloor m \rfloor + \tau$ and $\vartheta_m + m/\eta < 1$, one can choose $\tau' \in (\tau, 1)$ and $\delta > 0$ such that

$$T_{\tau'} T_{1+\delta}^{\lfloor m \rfloor}(\vartheta_m) := \vartheta_m + \frac{\lfloor m \rfloor}{\eta} + \frac{\tau'}{\eta} + \frac{\lfloor m \rfloor \delta}{\eta} = 1.$$

Lemma 5 shows that $Q(\cdot) \in \mathcal{C}_b^m(B, \mathcal{L}(\mathcal{B}_{\vartheta_m}, \mathcal{B}_1))$. Since $1_E \in \mathcal{B}_{\vartheta_m}$ and $\pi \in \mathcal{B}'_1$, one gets: $\pi(Q(\cdot)1_E) \in \mathcal{C}_b^m(B, \mathbb{C})$. Lemma 6 below provides the above claimed regularity properties involving the resolvent $(z - Q(t))^{-1}$, and thus completes the proof of Proposition 2. \square

Lemma 6 *There exists $\kappa \in (\kappa_{\vartheta_m}, 1)$ and $R > 0$ such that, for any probability measure ν on E satisfying $\nu(v) < \infty$ and for any $g \in \mathcal{B}_{\vartheta_m}$, the map $t \mapsto r_z(t) := \nu((z - Q(t))^{-1}g)$ is in $\mathcal{C}_b^m(B(0, R), \mathbb{C})$ for all $z \in \mathcal{D}_\kappa$, and we have*

$$\forall \ell = 0, \dots, \lfloor m \rfloor, \sup \{ |r_z^{(\ell)}(t)|, z \in \mathcal{D}_\kappa, t \in B(0, R) \} < \infty,$$

where $r_z^{(\ell)}(\cdot)$ denotes any partial derivative of order ℓ of $r_z(\cdot)$. Moreover, if $m \notin \mathbb{N}$, then the τ -Hölder coefficient of $r_z^{(\lfloor m \rfloor)}$ on $B(0, R)$ is uniformly bounded in $z \in \mathcal{D}_\kappa$.

Proof. On the basis of Lemmas 4-5, Lemma 6 is established in [17, Prop. 10.3] in the case $m \in \mathbb{N}$. The case $m \notin \mathbb{N}$ can be obtained by slightly extending the method of [17]. It can be also deduced from [9, Th. 3.3] which specifies and generalizes the Taylor expansions obtained

in [15, 10]. Let us verify that the assumptions of [9, Th. 3.3] are fulfilled. Define the spaces $\mathcal{B}^{[m]+1} := \mathcal{B}_{\vartheta_m}$, $\mathcal{B}^0 = \mathcal{B}_{T_r, T_{1+\delta}^{[m]}(\vartheta_m)} = \mathcal{B}_1$ and

$$\forall j = 1, \dots, [m], \quad \mathcal{B}^j := \mathcal{B}_{T_{1+\delta}^{[m]+1-j}(\vartheta_m)}.$$

Note that $\mathcal{B}^{[m]+1} \subset \mathcal{B}^{[m]} \subset \dots \subset \mathcal{B}^1 \subset \mathcal{B}^0$. Then Lemmas 4-5 and [9, Th. 3.3] show that there exist $\kappa \in (\kappa_{\vartheta_m}, 1)$ and $R > 0$ such that $r_z(t)$ admits a Taylor expansion of order $[m]$ at any point $t_0 \in B(0, R)$, with a remainder $O(|t - t_0|^m)$ which is uniform in $z \in \mathcal{D}_\kappa$ and $t_0 \in B(0, R)$. As already observed in [9, Rk. 3.6], the passage to the m -differentiability properties of $r_z(\cdot)$ can be deduced from a general lemma in [6]. \square

Proposition 3 *Assume that ξ is nonlattice and satisfies Condition (21), that $\mu(v) < \infty$, and that $f \in \mathcal{B}_{\vartheta_m}$, where ϑ_m is defined as in Proposition 2. Then Hypothesis $\mathcal{R}(m)$ -(ii) is fulfilled.*

Proof. Let $0 < r < r'$. The nonlattice assumption yields the following result (see [17, Lem. 10.1, Prop. 5.4]):

Lemma 7 *For any $\theta \in (0, 1]$, there exists $\rho_\theta < 1$ such that: $\sup_{t \in K_{r, r'}} \|Q(t)^n\|_\theta = O(\rho_\theta^n)$, where $\|\cdot\|_\theta$ stands here for the operator norm on \mathcal{B}_θ .*

Lemma 7 and (17) imply that the series $\mathcal{E}(t) := \sum_{n \geq 1} \mathbb{E}_\mu[f(X_n) e^{i\langle t, S_n \rangle}]$ uniformly converge on $K_{r, r'}$ (use $f \in \mathcal{B}_{\vartheta_m}$ and $\mu \in \mathcal{B}'_1 \subset \mathcal{B}'_{\vartheta_m}$). Setting $\theta_j := T_{1+\delta}^{[m]-j}(\vartheta_m)$ for $j = 0, \dots, [m]$, we define $\rho = \max\{\rho_1, \rho_{\theta_0}, \dots, \rho_{\theta_{[m]}}\}$. Let $\rho_0 \in (\rho, 1)$, and let Γ denote the oriented circle centered at $z = 0$, with radius ρ_0 . Since $Q(t)^n = \frac{1}{2i\pi} \oint_\Gamma z^n (z - Q(t))^{-1} dz$ in $\mathcal{L}(\mathcal{B}_{\vartheta_m})$ for all $t \in K_{r, r'}$, we have

$$\mathcal{E}(t) = \sum_{n \geq 1} \mu(Q(t)^n f) = \frac{1}{2i\pi} \oint_\Gamma \frac{z}{1-z} \mu((z - Q(t))^{-1} f) dz.$$

The desired regularity of $\mathcal{E}(\cdot)$ can be then deduced as in the proof of Proposition 2. \square

Remark 2 *If ξ is π -centered and satisfies (21) with $\eta > 2$, then we have $\nabla \lambda(0) = 0$. If $\mu(v) < \infty$, then we have: $-\text{Hess } \lambda(0) = \lim_n \frac{1}{n} \mathbb{E}_\mu[S_n S_n^*]$. Finally, if ξ is nonlattice, then $\Sigma := -\text{Hess } \lambda(0)$ is positive definite.*

Proof of Remark 2. Assume that $\mu(v) < \infty$. We have $\mathbb{E}_\mu[\|S_n\|^2] < \infty$ (use (21)). From (26) and Lemma 6, it follows that $\sup_{n \geq 1} |R_{n, \mu, 1_E}^{(\ell)}(0)| < \infty$ for $\ell = 1, 2$. By Proposition 1(i), we get: $\nabla \lambda(0)/i = \lim_n \frac{1}{n} \mathbb{E}_\mu[S_n]$. When applied with $\mu = \pi$, this gives $\nabla \lambda(0) = 0$ since ξ is π -centered. The second point of Remark 2 follows from Proposition 1(ii), and the last assertion is established in [17, Sect. 5.2]. \square

The following remarks also apply to the examples of the two next subsections.

Remark 3 Even in case $d \geq 5$ for which Hypothesis $\mathcal{R}(m_d)$ is needed in Theorem 1 with $m_d = d - 2$, the above use of [9, Th. 3.3] (or [17, Sect. 7]) does not allow to prove (4) under the moment assumption $\|\xi\|^{m_d} \leq C v$. This is due to the fact that, when ξ is unbounded, the property $Q \in \mathcal{C}^\varepsilon(\mathbb{R}^d, \mathcal{L}(\mathcal{B}_\theta))$ is not fulfilled in general and must be replaced in the derivation procedure (of both [17, 9]) by $Q \in \mathcal{C}^\varepsilon(\mathbb{R}^d, \mathcal{L}(\mathcal{B}_\theta, \mathcal{B}_{\theta'}))$ with $\theta < \theta'$. This yields a "gap" between the spaces $\mathcal{B}_{\vartheta_m}$ and \mathcal{B}_v (used in Lemma 6) which is slightly bigger than the expected one. This is the reason why the order in (20) is $m_d + \delta_0$ with some arbitrary small $\delta_0 > 0$.

Remark 4 Property (4) could be investigated in the general setting of the sequences $(X_n, S_n)_{n \geq 0}$ satisfying Condition (\mathcal{G}) of Subsection 3.1, provided that the driving Markov chain $(X_n)_{n \geq 0}$ is v -geometrically ergodic. The moment conditions then focus on (S_1, X_1) .

4.2 Applications to the ρ -mixing Markov chains

Let us assume that the σ -field \mathcal{E} is countably generated, and that $(X_n)_{n \geq 0}$ is a ρ -mixing Markov chain. Equivalently (see [22]), $(X_n)_{n \geq 0}$ possesses a stationary distribution π and satisfies the strong ergodicity condition (16) on the usual Lebesgue space $\mathbb{L}^2(\pi)$. For instance, this condition is fulfilled when $(X_n)_{n \geq 0}$ is uniformly ergodic (i.e. satisfies (16) on the space \mathcal{B}^∞ of bounded measurable complex-valued functions on E , or equivalently $(X_n)_{n \geq 0}$ is aperiodic, ergodic, and satisfies the so-called Doeblin condition). Let us first present a statement in the stationary case.

Corollary 3 If $\xi : E \rightarrow \mathbb{R}^d$ ($d \geq 3$) is a π -centered nonlattice functional such that

$$\exists \delta_0 > 0, \quad \pi(\|\xi\|^{m_d + \delta_0}) < \infty, \quad (30)$$

then we have (1) in the stationary case ($\mu = \pi$) with Σ defined by (18).

In comparison with the i.i.d. case, the moment condition (30) is the expected one (up to $\delta_0 > 0$). Corollary 3 is a special case of the next one. We denote by $\mathbb{L}^p(\pi)$, $1 \leq p \leq \infty$, the usual Lebesgue space associated to π .

Corollary 4 Assume that ξ satisfies the assumptions of Corollary 3 and that the initial distribution μ is of the form $\mu = \phi d\pi$, where $\phi \in \mathbb{L}^{r'}(\pi)$ for some $r' > \frac{\eta}{\eta - m}$, where $\eta = m_d + \delta_0$, $m \in (2, 2 + \delta_0)$ if $d = 3, 4$, and $m = d - 2$ if $d \geq 5$. If $s > \frac{\eta}{\eta - m}$ and $\frac{\eta s}{\eta + m s} > \frac{r'}{r' - 1}$, then, for any $f \in \mathbb{L}^s(\pi)$, $f \geq 0$, we have (4) with $L(0) = \pi(f)$ and Σ defined by (18).

Given $r' > \eta/(\eta - m)$, the two last conditions on s are fulfilled if s is sufficiently large. Indeed, let us set $r = r'/(r' - 1)$ (ie. $1/r + 1/r' = 1$). Since $1/r' < 1 - m/\eta$, we have $1 < r < \eta/m$, and since $\frac{\eta s}{\eta + m s} \nearrow \frac{\eta}{m}$ when $s \rightarrow +\infty$, we have $\eta s/(\eta + m s) > r$ when s is large enough.

Thanks to Theorem 1, Corollary 4 is a consequence of the next Proposition 4. Let us consider any real numbers m, η such that $0 < m < \eta$, set $\tau = m - \lfloor m \rfloor$, and assume that $\xi : E \rightarrow \mathbb{R}^d$ is a measurable function such that:

$$\pi(\|\xi\|^\eta) < \infty. \quad (31)$$

Let $r' > \frac{\eta}{\eta-m}$, $r := \frac{r'}{r'-1}$, and let s be such that $s > \frac{\eta}{\eta-m}$ and $\frac{\eta s}{\eta+ms} > r$. Note that $r < s$.

Proposition 4 *Assume that we have (31), that $\mu = \phi d\pi$ with $\phi \in \mathbb{L}^{r'}(\pi)$, and that $f \in \mathbb{L}^s(\pi)$, with r' and s above defined. Then Hypothesis $\mathcal{R}(m)$ -(i) holds true with $L(0) = \pi(f)$. If in addition ξ is nonlattice, then Hypothesis $\mathcal{R}(m)$ -(ii) is fulfilled.*

Proof. We are going to apply the procedure of the previous subsection to some suitable spaces chosen in the family $\{\mathcal{B}_\theta := \mathbb{L}^\theta(\pi), \theta \in (1, +\infty)\}$. First the conclusions of Lemma 4 are true w.r.t. these spaces, see [22] and [17, Prop. 4.1]. Second, from the nonlattice condition, Lemma 7 is valid, see [17, Prop. 5.4, Sec. 5.3]. Now let us define for any fixed $u > 0$: $T_u(\theta) := \eta\theta/(\eta + u\theta)$. Then, by using (31) and an easy extension of [17, Lem. 7.4], we can prove that we have for any $\theta \in I := [r, s]$ and $\tau' \in (0, 1)$:

$$\begin{aligned} \forall j = 1, \dots, [m]: \quad T_1^j(\theta) \in I &\Rightarrow Q(\cdot) \in \mathcal{C}_b^j(B, \mathcal{L}(\mathbb{L}^\theta, \mathbb{L}^{T_1^j(\theta)})) \\ \forall j = 0, \dots, [m]: \quad T_{\tau'} T_1^j(\theta) \in I &\Rightarrow Q(\cdot) \in \mathcal{C}_b^{j+\tau'}(B, \mathcal{L}(\mathbb{L}^\theta, \mathbb{L}^{T_{\tau'} T_1^j(\theta)})), \end{aligned}$$

where $B = B(0, R)$ (for any $R > 0$). Next, from $r < \frac{\eta s}{\eta+ms}$, one can fix $\tau' \in (\tau, 1)$ such that

$$r \leq T_{\tau'} T_1^{[m]}(s) = \frac{\eta s}{\eta + ([m] + \tau')s}.$$

By using the spaces $\mathcal{B}^{[m]+1} := \mathbb{L}^s(\pi)$, $\mathcal{B}^0 = \mathbb{L}^r(\pi)$ and

$$\forall j = 1, \dots, [m], \quad \mathcal{B}^j := \mathbb{L}^{T_1^{[m]+1-j}(s)}(\pi),$$

one can prove as in Subsection 4.1 that the conclusions of Lemma 6 are fulfilled for any probability measure ν on E defining a continuous linear form on $\mathbb{L}^r(\pi)$ and for any $g \in \mathbb{L}^s(\pi)$. Then the two conclusions of Proposition 4 can be proved as in Subsection 4.1. \square

To complete the proof of Corollary 4, notice that, if (31) holds with $\eta > 2$, then the conclusions of Remark 2 are fulfilled (here the condition on μ is that of Proposition 4).

4.3 Applications to iterative Lipschitz models

Here (E, d) is a non-compact metric space in which every closed ball is compact, and it is endowed with its Borel σ -field \mathcal{E} . Let (G, \mathcal{G}) be a measurable space, let $(\varepsilon_n)_{n \geq 1}$ be a i.i.d. sequence of random variables taking values in G , and let $F : E \times G \rightarrow E$ be a measurable function. Given an initial E -valued r.v. X_0 independent of $(\varepsilon_n)_{n \geq 1}$, the random iterative model associated to $(\varepsilon_n)_{n \geq 1}$, F and X_0 is defined by (see [7])

$$X_n = F(X_{n-1}, \varepsilon_n), \quad n \geq 1.$$

Let us consider the two following random variables which are classical in these models [7] :

$$\mathcal{C} := \sup \left\{ \frac{d(F(x, \varepsilon_1), F(y, \varepsilon_1))}{d(x, y)}, x, y \in E, x \neq y \right\} \quad \text{and} \quad \mathcal{M} = 1 + \mathcal{C} + d(F(x_0, \varepsilon_1), x_0)$$

where x_0 is some fixed point in E . It is well-known that, if $\mathcal{C} < 1$ almost surely and if

$$\exists s \geq 0, \exists \delta_0 > 0, \quad \mathbb{E}[\mathcal{M}^{(s+1)m_d + \delta_0}] < \infty, \quad (32)$$

then $(X_n)_{n \geq 0}$ possesses a stationary distribution π such that $\pi(d(\cdot, x_0)^{(s+1)m_d + \delta_0}) < \infty$.

Corollary 5 *Let us assume that $\mathcal{C} < 1$ almost surely, that (32) holds, that the initial distribution μ is such that $\mu(d(\cdot, x_0)^{(s+1)m_d + \delta_0}) < \infty$, and finally that $\xi : E \rightarrow \mathbb{R}^d$ ($d \geq 3$) is a π -centered nonlattice function satisfying the following condition:*

$$\exists S \geq 0, \forall (x, y) \in E \times E, \quad \|\xi(x) - \xi(y)\| \leq S d(x, y) [1 + d(x, x_0) + d(y, x_0)]^s, \quad (33)$$

where s is the real number in (32). Then we have (1) with Σ defined by (18).

Notice that (33) corresponds to the general weighted-Lipschitz condition introduced in [7]. Some weaker assumptions on \mathcal{C} are presented at the end of this subsection, as well as some (a priori) less restrictive condition than the nonlattice assumption.

Example: the linear autoregressive models in \mathbb{R}^3 . As an illustration, let us consider in $E = \mathbb{R}^3$, equipped with some norm $\|\cdot\|$, the autoregressive model $X_n = AX_{n-1} + \varepsilon_n$, where $X_0, \varepsilon_1, \varepsilon_2, \dots$ are independent \mathbb{R}^3 -valued random variables, and A is a contractive matrix of order 3 ($\sup_{\|x\| \leq 1} \|Ax\| < 1$). Then, taking the distance $d(x, y) = \|x - y\|$, $(X_n)_n$ is an iterative Lipschitz model and we have $\mathcal{C} < 1$. Let us consider the centered random walk $S_n = X_1 + \dots + X_n - n\mathbb{E}_\pi[X_0]$ associated to the functional $\xi(x) := x - \mathbb{E}_\pi[X_0]$. Note that ξ satisfies (33) with $s = 0$. Assume that ξ is nonlattice and that we have

$$\exists \delta_0 > 0, \quad \mathbb{E}[\|X_0\|^{2+\delta_0} + \|\varepsilon_1\|^{2+\delta_0}] < \infty. \quad (34)$$

Then, from Corollary 5, the above random walk S_n satisfies (1). This statement extends the 3-dimensional renewal theorem of the i.i.d. case (obtained here in the special case $A = 0$) under the same moment condition up to $\delta_0 > 0$. Also notice that, except the moment condition on ε_1 in (34), the previous result does not require any special assumption on the law of ε_1 . Anyway, it remains true if $X_n = A_n X_{n-1} + \varepsilon_n$ where $(A_n)_{n \geq 1}$ is a sequence of r.v. taking values in the set of matrices of order 3, such that $\|A_1\| < 1$ a.s. and $(A_n, \varepsilon_n)_{n \geq 1}$ is i.i.d. and independent of X_0 .

Corollary 5 follows from the more general Corollary 6 below, that will be proved by applying the operator-type method to the weighted Hölder-type spaces $\mathcal{B}_{\alpha, \beta, \gamma}$ defined as follows. For $x \in E$, set $p(x) = 1 + d(x, x_0)$, and given any $0 < \alpha \leq 1$ and $0 < \beta \leq \gamma$, define for $(x, y) \in E^2$:

$$\Delta_{\alpha, \beta, \gamma}(x, y) = p(x)^{\alpha\gamma} p(y)^{\alpha\beta} + p(x)^{\alpha\beta} p(y)^{\alpha\gamma}.$$

The space $\mathcal{B}_{\alpha, \beta, \gamma}$ is by definition composed of the \mathbb{C} -valued functions f on E such that

$$m_{\alpha, \beta, \gamma}(f) = \sup \left\{ \frac{|f(x) - f(y)|}{d(x, y)^\alpha \Delta_{\alpha, \beta, \gamma}(x, y)}, x, y \in E, x \neq y \right\} < \infty.$$

Set $|f|_{\alpha, \gamma} = \sup_{x \in E} \frac{|f(x)|}{p(x)^{\alpha(\gamma+1)}}$ and $\|f\|_{\alpha, \beta, \gamma} = m_{\alpha, \beta, \gamma}(f) + |f|_{\alpha, \gamma}$. Then $(\mathcal{B}_{\alpha, \beta, \gamma}, \|\cdot\|_{\alpha, \beta, \gamma})$ is a Banach space.

Corollary 6 *Let us assume that the assumptions of Corollary 5 hold, and let us fix any real number α such that $0 < \alpha \leq \min\{1, \frac{\delta_0}{4(s+1)}\}$. Then we have (4), with $L(0) = \pi(f)$ and Σ defined by (18), for each nonnegative function $f \in \mathcal{B}_{\alpha, \vartheta, \vartheta}$, where ϑ is some real number such that $\vartheta > s + 1$ (the condition on ϑ is specified below in function of α and δ_0).*

Corollary 6 is a consequence of Theorem 1 and of the next Proposition 5 which, given $m \in \mathbb{N}$, gives Hypothesis $\mathcal{R}(m + \delta)$ for some $\delta > 0$ (this is convenient for our purpose).

We assume below that $\mathcal{C} < 1$ a.s. and that we have:

$$\exists m \in \mathbb{N}, \exists s \geq 0, \exists \delta_0 > 0, \quad \mathbb{E}[\mathcal{M}^{(s+1)m + \delta_0}] < \infty, \quad (35)$$

and that $\xi : E \rightarrow \mathbb{R}^d$ is a measurable function satisfying (33) with s given in (35).

Let $k \in \mathbb{N}$, $\alpha \in (0, 1]$, $\delta' \in (0, \alpha)$, and let β be such that $s + 1 \leq \beta \leq \gamma$. Set $B = B(0, R)$ (for any $R > 0$).

Lemma 8 *Let us assume that $\gamma' \geq \gamma + \frac{(s+1)(k+\delta')}{\alpha}$ and $\mathbb{E}[\mathcal{M}^{\alpha(\gamma'+\beta)}] < \infty$. Then we have $Q(\cdot) \in \mathcal{C}_b^{k+\delta'}(B, \mathcal{L}(\mathcal{B}_{\alpha, \beta, \gamma}, \mathcal{B}_{\alpha, \beta, \gamma'}))$.*

Lemma 8 can be proved by using the arguments of [17, Lem. B.4-4']. Now, let us define for any $u > 0$: $T_u(\gamma) := \gamma + (s+1)u/\alpha$. Let α be fixed such that $0 < \alpha \leq \min\{1, \frac{\delta_0}{4(s+1)}\}$. Set $\vartheta := s + 1 + \delta'$ for some $\delta' \in (0, \alpha)$ (specified below), and define:

$$\gamma_m := T_{\delta'} T_{1+\delta'}^m(\vartheta) = s + 1 + \delta' + (s+1)(m(1+\delta') + \delta')/\alpha.$$

We have $\alpha(\gamma_m + \vartheta) = (s+1)m + 2\alpha(s+1) + 2\alpha\delta' + \delta'(s+1)(m+1)$. Now, using $\alpha \leq \frac{\delta_0}{4(s+1)}$, we can choose $\delta' \in (0, \alpha)$ sufficiently small such that $\alpha(\gamma_m + \vartheta) \leq (s+1)m + \delta_0$.

Proposition 5 *Let us assume that we have $\mathcal{C} < 1$ a.s. and (35), that ξ satisfies (33) with s given in (35), that $\mu(d(\cdot, x_0)^{(s+1)m + \delta_0}) < \infty$, and finally that $f \in \mathcal{B}_{\alpha, \vartheta, \vartheta}$. Then, for any $\delta \in (0, \delta')$, Hypothesis $\mathcal{R}(m + \delta)$ -(i) holds with $L(0) = \pi(f)$. If in addition ξ is nonlattice, then Hypothesis $\mathcal{R}(m + \delta)$ -(ii) is fulfilled.*

Proof. We apply the procedure of Subsection 4.1 with the spaces $\mathcal{B}_\gamma := \mathcal{B}_{\alpha, \vartheta, \gamma}$, $\gamma \in [\vartheta, \gamma_m]$, with $\alpha, \vartheta, \gamma_m$ fixed above. First, Lemma 4 is valid on each \mathcal{B}_γ , see [17, Prop. 11.2-11.4]. Second, from the nonlattice condition, Lemma 7 applies on each \mathcal{B}_γ , see [17, Prop. 11.8]. Next, observe that $\gamma_m > \vartheta + (s+1)(m + \delta')/\alpha$. Then, by using Lemma 8 and the spaces $\mathcal{B}^{[m]+1} := \mathcal{B}_{\alpha, \vartheta, \vartheta}$, $\mathcal{B}^0 = \mathcal{B}_{\alpha, \vartheta, \gamma_m}$ and

$$\forall j = 1, \dots, [m], \quad \mathcal{B}^j := \mathcal{B}_{\alpha, \vartheta, T_{1+\delta'}^{m+1-j}(\vartheta)},$$

one can prove as in Subsection 4.1 that Lemma 6 holds with here any $g \in \mathcal{B}_{\alpha, \vartheta, \vartheta}$ and any probability measure ν on E defining a continuous linear form on $\mathcal{B}_{\alpha, \vartheta, \gamma_m}$. Since $\alpha(\gamma_m + 1) \leq (s+1)m + \delta_0$, the last condition holds if $\mu(d(\cdot, x_0)^{(s+1)m + \delta_0}) < \infty$. Then the conclusions of Proposition 5 can be established as in Subsection 4.1. \square

To complete the proof of Corollary 6, notice that, under the assumptions of Proposition 5 with $m \geq 2$ in (35), the conclusions of Remark 2 are fulfilled.

Remark 5 *By repeating the previous proof with a more precise use of the results of [17], one can establish that under the following assumption (weaker than (32) and $C < 1$)*

$$\exists s \geq 0, \exists \delta_0 > 0, \mathbb{E}[(1 + C^\alpha) \mathcal{M}^{(s+1)m_d + \delta_0}] < \infty, \mathbb{E}[C^\alpha \max\{C, 1\}^{(s+1)m_d + \delta_0}] < 1,$$

the conclusion of Corollary 6 remains true if $\mu(d(\cdot, x_0)^{(s+1)m_d + \delta_0}) < \infty$, and if the π -centered function ξ satisfies (33) and the following non-arithmeticity condition (a priori weaker for iterative models than the nonlattice assumption):

There exist no $t \in \mathbb{R}^d$, $t \neq 0$, no $\lambda \in \mathbb{C}$, $|\lambda| = 1$, no π -full Q -absorbing set $A \in \mathcal{E}$, and finally no bounded function $w \in \mathcal{B}_{\alpha, \vartheta, \gamma_{m_d}}$ whose modulus is nonzero constant on A , such that we have : $\forall x \in A, e^{i\langle t, \xi(y) \rangle} w(y) = \lambda w(x) \quad Q(x, dy) - a.s.$

5 Conclusion

In this paper we extend the well-known renewal theorem established by Spitzer [23] in dimension $d \geq 3$ for centered nonlattice i.i.d. sequences. In a first step, given a measurable space E and a sequence $(X_n, S_n)_{n \geq 0}$ of $E \times \mathbb{R}^d$ -valued random variables, we present the “tailor-made” assumptions on $\mathbb{E}[f(X_n) e^{i\langle t, S_n \rangle}]$ under which such a renewal theorem can be proved by Fourier techniques. In a second step, considering the case when $(X_n)_{n \geq 0}$ is a Markov chain, we give a general setting containing the Markov random walks, for which the spectral method may provide an efficient way to study the term $\mathbb{E}[f(X_n) e^{i\langle t, S_n \rangle}]$. This approach, already used in [1], is improved here by appealing to the weak spectral method developed in [17]. Our main improvements concern the operator-type moment assumptions: when applied to the ν -geometrically ergodic Markov chains, the ρ -mixing Markov chains and the iterative Lipschitz models, these assumptions are fulfilled under the (almost) expected moment condition in comparison with the i.i.d. case. In particular, our results apply to unbounded r.v. S_1 , while the moment condition derived from [1] is not satisfied in general when S_1 is not bounded.

The present weak spectral procedure should also enable to improve some of the results of [1, 8] concerning the non-centered Markov renewal theorem in dimension $d \geq 2$. Work in these directions by the first author is in progress. Finally this method could be used to study the renewal theorem for Birkhoff sums in dynamical systems by using the so-called Perron-Frobenius operator, as already developed for other purposes in [9].

APPENDIX A. Complements in the proof of Theorem 1

A.0. Inequalities for the derivatives of $1/w$.

Let $\gamma = (\gamma_1, \dots, \gamma_d) \in \mathbb{N}^d$, and set $|\gamma| = \gamma_1 + \dots + \gamma_d$ and $\gamma! = \gamma_1! \dots \gamma_d!$. If $\beta \in \mathbb{N}^d$ is such that $\beta \leq \gamma$, namely $\beta_i \leq \gamma_i$ for each $i \in \{1, \dots, d\}$, we set $\binom{\gamma}{\beta} = \frac{\gamma!}{\beta!(\gamma-\beta)!}$.

Let Ω be an open subset of \mathbb{R}^d , let $m \in \mathbb{N}^*$. We denote by $\mathcal{C}^m(\Omega, \mathbb{C})$ the space of m -times continuously differentiable complex-valued functions on Ω . If $\gamma \in \mathbb{N}^d$ is such that $|\gamma| \leq m$, we denote by ∂^γ the derivative operator defined on $\mathcal{C}^m(\Omega, \mathbb{C})$ by :

$$\partial^\gamma := \frac{\partial^{|\gamma|}}{\partial x_1^{\gamma_1} \dots \partial x_d^{\gamma_d}} = \partial_1^{\gamma_1} \dots \partial_d^{\gamma_d} \quad \text{where} \quad \partial_j := \frac{\partial}{\partial x_j}.$$

Let us recall (Leibniz's formula) that, if f and g are in $\mathcal{C}^m(\Omega, \mathbb{C})$, then we have for all $\gamma \in \mathbb{N}^d$ such that $|\gamma| \leq m$,

$$\partial^\gamma(f \cdot g) = \sum_{\beta \leq \gamma} \binom{\gamma}{\beta} \partial^\beta f \cdot \partial^{\gamma-\beta} g.$$

Let V be a bounded neighborhood of 0 in \mathbb{R}^d .

Proposition A.0. *Assume that $w : \bar{V} \rightarrow \mathbb{C}$ is m -times continuously differentiable, and that there exist some constants $c > 0$ and $d \geq 0$ such that we have for all $x \in V$: $|w(x)| \geq c \|x\|^2$ and $\sum_{j=1}^d |(\partial_j w)(x)| \leq d \|x\|$. Then, for each $\gamma \in \mathbb{N}^d$ such that $|\gamma| \leq m$, there exists a constant $C_\gamma \geq 0$ such that*

$$\forall x \in V \setminus \{0\}, \quad \left| \partial^\gamma \left(\frac{1}{w} \right) (x) \right| \leq \frac{C_\gamma}{\|x\|^{2+|\gamma|}}. \quad (36)$$

Proof. Let $\gamma = (\gamma_1, \dots, \gamma_d) \in \mathbb{N}^d$ be such that $|\gamma| \leq m$, set $M_\gamma = \max_{x \in \bar{V}} |\partial^\gamma w(x)|$, and let $K > 0$ be such that $\|x\| \leq K$ for all $x \in V$. To prove (36), we are going to use an induction on $l = |\gamma| \in \{0, \dots, m\}$. First, (36) is obvious if $l = 0$, and if $l = 1$, then (36) follows from the following equalities

$$\forall j \in \{1, \dots, d\}, \quad \forall x \in V \setminus \{0\}, \quad \partial_j \left(\frac{1}{w} \right) (x) = - \frac{\partial_j w(x)}{w^2(x)}.$$

Now let $l \in \{1, \dots, m-1\}$, suppose that (36) is fulfilled for each $\gamma \in \mathbb{N}^d$ such that $|\gamma| \leq l$, and consider any $\gamma \in \mathbb{N}^d$ such that $|\gamma| = l+1$. Since $\partial^\gamma(w^{-1} \cdot w) = 0$, the Leibniz formula gives on $V \setminus \{0\}$:

$$\partial^\gamma \left(\frac{1}{w} \right) = - \frac{1}{w} \sum_{\beta \leq \gamma, \beta \neq \gamma} \binom{\gamma}{\beta} \partial^\beta \left(\frac{1}{w} \right) \cdot \partial^{\gamma-\beta} w.$$

Let $\beta \leq \gamma$ be such that $|\gamma| - |\beta| \geq 2$ and let $x \in V \setminus \{0\}$. We have :

$$\left| \frac{1}{w(x)} \partial^\beta \left(\frac{1}{w} \right) (x) \partial^{\gamma-\beta} w(x) \right| \leq \frac{C_\beta M_{\gamma-\beta}}{c \|x\|^{4+|\beta|}} \leq \frac{K^{|\gamma|-|\beta|-2} C_\beta M_{\gamma-\beta}}{c \|x\|^{2+|\gamma|}}.$$

Let $\beta \leq \gamma$ be such that $|\gamma| - |\beta| = 1$, and let $x \in V \setminus \{0\}$. Then we have :

$$\left| \frac{1}{w(x)} \partial^\beta \left(\frac{1}{w} \right) (x) \partial^{\gamma-\beta} w(x) \right| \leq \frac{d C_\beta}{c \|x\|^{3+|\beta|}} = \frac{d C_\beta}{c \|x\|^{2+|\gamma|}}.$$

This yields (36) for $|\partial^\gamma(\frac{1}{w})|$. □

Remarks.

(a) The derivative estimates (6) needed for the function $u(t) = \frac{\theta_1(t)}{1-\lambda(t)}$ in the proof of Lemma 2 follows from this proposition. Indeed, let us set $w(t) = 1 - \lambda(t)$ for $t \in V := [-\alpha, \alpha]^d$. Since $\nabla\lambda(0) = 0$ and Σ is positive definite, w satisfies the hypotheses of Proposition A.0. Besides, one may assume that $\sqrt{d}\alpha < 1$, so that each $t \in V$ satisfies $\|t\| < 1$. Now let $\gamma \in \mathbb{N}^d$ be such that $|\gamma| \leq d-2$, and let $C > 0$ and $D_\gamma \geq 0$ be such that we have for all $t \in V : |\theta_1(t)| \leq C\|t\|$ and $|\partial^\gamma \theta_1(t)| \leq D_\gamma$. On the one hand, we have

$$\forall t \in V \setminus \{0\}, \quad |\theta_1(t) \partial^\gamma \left(\frac{1}{w}\right)(t)| \leq \frac{C C_\gamma}{\|t\|^{1+|\gamma|}}.$$

On the other hand, for $\beta \in \mathbb{N}^d$ such that $\beta \neq \gamma$, $\beta \leq \gamma$, we have since $|\beta| \leq |\gamma| - 1$ and $\|t\| < 1$,

$$\forall t \in V \setminus \{0\}, \quad |\partial^{\gamma-\beta} \theta_1(t) \partial^\beta \left(\frac{1}{w}\right)(t)| \leq \frac{D_{\gamma-\beta} C_\beta}{\|t\|^{2+|\beta|}} \leq \frac{D_{\gamma-\beta} C_\beta}{\|t\|^{1+|\gamma|}}.$$

So (6) follows from Leibniz's formula.

(b) The derivative estimates (7) (8) on the function $v(t) = \frac{\theta_3(t)}{(1-\lambda(t))\langle \Sigma t, t \rangle}$ in the proof of Lemma 2 can be derived similarly. Indeed, set $s(t) = \langle \Sigma t, t \rangle$ for $t \in [-\alpha, \alpha]^d$. Then the function $s(\cdot)$ satisfies the assumptions of Proposition A.0 on $V = [-\alpha, \alpha]^d$. Let $\gamma \in \mathbb{N}^d$ be such that $|\gamma| \leq d-2$, and let $t \in V \setminus \{0\}$. From Leibniz's formula, the partial derivative $\partial^\gamma v(t)$ is a sum of terms of the form (up to some binomial coefficients)

$$\partial^\beta \theta_3(t) \partial^\delta \left(\frac{1}{w}\right)(t) \partial^\eta \left(\frac{1}{s}\right)(t) \quad \text{with } |\beta| + |\delta| + |\eta| = |\gamma|,$$

where $w(t) = 1 - \lambda(t)$. By proceeding as above (consider here the three cases $|\beta| = 0, 1, 2$, and the case $|\beta| \geq 3$), one can then establish (7) and (8).

A.1. Some inequalities for Fourier transforms.

Let $u : \mathbb{R}^d \setminus \{0\} \rightarrow \mathbb{C}$. We assume that there exists $\alpha > 0$ such that we have $u(x) = 0$ for all $x \in \mathbb{R}^d \setminus [-\alpha, \alpha]^d$.

Proposition A.1. *Let $k \in \mathbb{N}^*$. If u is k -times continuously differentiable on $\mathbb{R}^d \setminus \{0\}$ and satisfies the following conditions:*

- (i)_k each partial derivative of order $\leq k-1$ of u is $O(\|x\|^{-s})$ on $\mathbb{R}^d \setminus \{0\}$ for some $s < d-1$.
- (ii)_k each partial derivative of order k of u is Lebesgue-integrable on \mathbb{R}^d ,

then we have $\hat{u}(a) = o\left(\frac{1}{\|a\|^k}\right)$ when $\|a\| \rightarrow +\infty$.

This proposition is a consequence of the following lemma.

Lemma A.1. *If u is continuously differentiable on $\mathbb{R}^d \setminus \{0\}$ and satisfies (i)₁-(ii)₁, then we have $\widehat{\frac{\partial u}{\partial x_j}}(a) = ia_j \hat{u}(a)$ for $j = 1, \dots, d$, and $\hat{u}(a) = o\left(\frac{1}{\|a\|}\right)$ when $\|a\| \rightarrow +\infty$.*

In fact, the second conclusion in Lemma A.1 proves Proposition A.1 in the case $k = 1$, while the first conclusion of Lemma A.1 clearly allows to establish Proposition A.1 by induction.

Proof of Lemma A.1. Below, $\langle \cdot, \cdot \rangle$ and $\|\cdot\|$ denote the scalar product and the euclidean norm in both \mathbb{R}^d and \mathbb{R}^{d-1} . Let us write $x = (x_1, x') \in \mathbb{R} \times \mathbb{R}^{d-1}$ and $a = (a_1, a') \in \mathbb{R} \times \mathbb{R}^{d-1}$, and define

$$u_1(x_1) = \int_{|x'| \leq \alpha} u(x_1, x') e^{-i\langle a', x' \rangle} dx',$$

so that we have

$$\forall a \in \mathbb{R}^d, \quad \hat{u}(a) = \int_{-\alpha}^{\alpha} u_1(x_1) e^{-ia_1 x_1} dx_1.$$

Since u is continuous on $\mathbb{R}^d \setminus \{0\}$ and $x' \mapsto 1_{\{|x'| \leq \alpha\}}(x') \|x'\|^{-s}$ is Lebesgue-integrable on \mathbb{R}^{d-1} (because $s < d-1$), we deduce from Condition $(i)_1$ and Lebesgue's theorem that u_1 is continuous on \mathbb{R} . Moreover, given any segment K in \mathbb{R}^* , $\frac{\partial u}{\partial x_1}$ is bounded on the compact set $K \times [-\alpha, \alpha]^{d-1}$, and it then follows from Lebesgue's theorem that u_1 is continuously differentiable on K , with :

$$\forall x_1 \in K, \quad u_1'(x_1) = \int_{|x'| \leq \alpha} \frac{\partial u}{\partial x_1}(x_1, x') e^{-i\langle a', x' \rangle} dx'.$$

Now write

$$\hat{u}(a) = \lim_{\varepsilon \rightarrow 0} \left(\int_{\varepsilon}^{\alpha} u_1(x_1) e^{-ia_1 x_1} dx_1 + \int_{-\alpha}^{-\varepsilon} u_1(x_1) e^{-ia_1 x_1} dx_1 \right).$$

For $a_1 \neq 0$, an easy integration by parts gives (notice that $u_1(\alpha) = 0$ because $u(\alpha, \cdot) = 0$)

$$\int_{\varepsilon}^{\alpha} u_1(x_1) e^{-ia_1 x_1} dx_1 = \frac{e^{-ia_1 \varepsilon} u_1(\varepsilon)}{ia_1} + \frac{1}{ia_1} \int_{\varepsilon}^{\alpha} u_1'(x_1) e^{-ia_1 x_1} dx_1.$$

Using the continuity of $u_1(\cdot)$ at 0, the above expression of $u_1'(x_1)$, and finally the fact that $\frac{\partial u}{\partial x_1}$ is Lebesgue-integrable on \mathbb{R}^d by hypothesis, one gets

$$\lim_{\varepsilon \rightarrow 0} \left(\int_{\varepsilon}^{\alpha} u_1(x_1) e^{-ia_1 x_1} dx_1 \right) = \frac{u_1(0)}{ia_1} + \frac{1}{ia_1} \int_{[0, \alpha] \times [-\alpha, \alpha]^{d-1}} \frac{\partial u}{\partial x_1}(x) e^{-i\langle a, x \rangle} dx.$$

Similarly one can prove that

$$\lim_{\varepsilon \rightarrow 0} \left(\int_{-\alpha}^{-\varepsilon} u_1(x_1) e^{-ia_1 x_1} dx_1 \right) = \frac{-u_1(0)}{ia_1} + \frac{1}{ia_1} \int_{[-\alpha, 0] \times [-\alpha, \alpha]^{d-1}} \frac{\partial u}{\partial x_1}(x) e^{-i\langle a, x \rangle} dx.$$

So $\hat{u}(a) = \frac{1}{ia_1} \widehat{\frac{\partial u}{\partial x_1}}(a)$. On the same way, we have $\hat{u}(a) = \frac{1}{ia_j} \widehat{\frac{\partial u}{\partial x_j}}(a)$ ($j = 2, \dots, d$). This proves the first assertion of Lemma A.1. This also gives $\|a\| |\hat{u}(a)| \leq \sum_{j=1}^d |a_j| |\hat{u}(a)| = \sum_{j=1}^d |\widehat{\frac{\partial u}{\partial x_j}}(a)|$, and since $\widehat{\frac{\partial u}{\partial x_j}}(a) \rightarrow 0$ when $\|a\| \rightarrow +\infty$ (by $(ii)_1$), this yields the second assertion. \square

A.2. An elementary proof of (9).

Let us recall that we set in (9) : $c = (2\pi)^{\frac{d}{2}} 2^{\frac{d}{2}-2} \Gamma(\frac{d-2}{2})$. Equality (9) follows from the following proposition. Let $S(\mathbb{R}^d)$ denote the Schwartz space.

Proposition A.2. *Let $g \in S(\mathbb{R}^d)$. Then we have : $\int_{\mathbb{R}^d} \hat{g}(u) \frac{1}{\|u\|^2} du = c \int_{\mathbb{R}^d} g(v) \frac{1}{\|v\|^{d-2}} dv$.*

Proof. First observe that $\frac{1}{\|u\|^2} = \int_0^{+\infty} e^{-x\|u\|^2} dx$, so that we obtain by Fubini's theorem, Parseval's formula, and by setting $\gamma_x(u) = e^{-x\|u\|^2}$

$$\int_{\mathbb{R}^d} \hat{g}(u) \frac{1}{\|u\|^2} du = \int_0^{+\infty} \int_{\mathbb{R}^d} \left(\hat{g}(u) e^{-x\|u\|^2} du \right) dx = \int_0^{+\infty} \left(\int_{\mathbb{R}^d} g(v) \widehat{\gamma_x}(v) dv \right) dx,$$

with $\widehat{\gamma_x}(v) = \left(\frac{\pi}{x}\right)^{\frac{d}{2}} e^{-\frac{\|v\|^2}{4x}}$ from a usual computation. Now Fubini's theorem again yields

$$\int_0^{+\infty} \int_{\mathbb{R}^d} g(v) \widehat{\gamma_x}(v) dv dx = \pi^{\frac{d}{2}} \int_{\mathbb{R}^d} g(v) \left(\int_0^{+\infty} x^{-\frac{d}{2}} e^{-\frac{\|v\|^2}{4x}} dx \right) dv.$$

Finally, we have $\int_0^{+\infty} x^{-\frac{d}{2}} e^{-\frac{\|v\|^2}{4x}} dx = 2^{d-2} \Gamma\left(\frac{d-2}{2}\right) \|v\|^{-(d-2)}$ (by setting $y = \frac{\|v\|^2}{4x}$). \square

A.3. Proof of (10).

Recall that $*$ denotes the convolution product on \mathbb{R}^d , and that we set $f_{d-2}(w) := \|w\|^{2-d}$ in (10). Let us consider any $F \in S(\mathbb{R}^d)$, and set $F_\beta(x) = \beta^d F(\beta x)$ for any $\beta > 0$. Then :

Proposition A.3. $\lim_{\beta \rightarrow +\infty} (F_\beta * f_{d-2})(\tilde{b}) = \int F(w) dw$ uniformly in \tilde{b} such that $\|\tilde{b}\| = 1$.

Proof. Let $\varepsilon > 0$. Since $\|\tilde{b}\| = 1$, there clearly exists $0 < \eta = \eta(\varepsilon) < 1$ independent of \tilde{b} such that we have $|f_{d-2}(\tilde{b} - w) - 1| \leq \varepsilon$ if $\|w\| < \eta$. From $\int F_\beta(w) dw = \int F(w) dw$, one gets

$$\begin{aligned} \left| \int F(w) dw - (F_\beta * f_{d-2})(\tilde{b}) \right| &\leq \int |F_\beta(w)| |1 - f_{d-2}(\tilde{b} - w)| dw \\ &= \int_{\|w\| < \eta} + \int_{\|w\| \geq \eta} := A_{\eta, \tilde{b}} + B_{\eta, \tilde{b}}. \end{aligned}$$

We have $A_{\eta, \tilde{b}} \leq \varepsilon \int |F(w)| dw$. Besides we have $|F(\cdot)| \leq \frac{D}{(1+\|\cdot\|)^{d+1}}$ for some $D > 0$ (because $F \in S(\mathbb{R}^d)$), thus $B_{\eta, \tilde{b}} \leq B'_{\eta, \tilde{b}} + B''_{\eta, \tilde{b}}$ with $B'_{\eta, \tilde{b}} = \int_{\|w\| \geq \eta} |F_\beta(w)| dw$ and

$$B''_{\eta, \tilde{b}} = \int_{\|w\| \geq \eta} \left(\frac{D \beta^d}{\beta^{d+1} \|w\|^{d+1}} \right) \left(\frac{1}{\|\tilde{b} - w\|^{d-2}} \right) dw = \frac{D}{\beta} \int_{\|w\| \geq \eta} \frac{dw}{\|w\|^{d+1} \|\tilde{b} - w\|^{d-2}}.$$

For β large enough, we have $B'_{\eta, \tilde{b}} = \int_{\|y\| \geq \eta\beta} |F(y)| dy \leq \varepsilon$, and by decomposing $\int_{\|w\| \geq \eta}$ according that $\|\tilde{b} - w\| \leq 2$ or $\|\tilde{b} - w\| > 2$, and observing that $\|\tilde{b} - w\| > 2 \Rightarrow \|w\| > 1$, we obtain

$$B''_{\eta, \tilde{b}} \leq \frac{D}{\beta} \left(\frac{1}{\eta^{d+1}} \int_{\{\|\tilde{b}-w\| \leq 2\}} \frac{dw}{\|\tilde{b} - w\|^{d-2}} + \frac{1}{2^{d-2}} \int_{\{\|w\| > 1\}} \frac{dw}{\|w\|^{d+1}} \right).$$

Since the two previous integrals are finite, we have $B''_{\eta, \tilde{b}} \leq \varepsilon$ for β large enough. \square

References

- [1] BABILLOT M. *Théorie du renouvellement pour des chaînes semi-markoviennes transientes*. Ann. I. H. Poincaré, sect. B, Tome 24, No 4, 507-569 (1988).
- [2] BÁLINT P., GOUËZEL S. *Limit theorems in the stadium billiard*. Communications in Mathematical Physics **263**, 451-512 (2006).
- [3] BREIMAN L. *Probability* Classic in Applied Mathematics, SIAM, 1993.
- [4] BROISE A., DAL'BO F., PEIGNÉ M. *Études spectrales d'opérateurs de transfert et applications*. Astérisque 238 (1996).
- [5] D. BURACZEWSKI, E. DAMEK, Y. GUIVARC'H. *Convergence to stable laws for a class of multidimensional stochastic recursions*. Accepted for the publication in Probability Theory and Related Fields (2009).
- [6] CAMPANATO S. *Proprieta di una famiglia di spazi funzionali*. Ann. Scuola Norm. Sup. Pisa (3) 18 (1964), 137-160.
- [7] DUFLO M. *Random Iterative Models*. Applications of Mathematics, Springer-Verlag Berlin Heidelberg (1997).
- [8] FUH C.D, LAI T.L. *Asymptotic expansions in multidimensional Markov renewal theory and first passage times for Markov random walks*. Adv. in Appl. Probab. **33**, 652-673 (2001).
- [9] GOUËZEL S. *Necessary and sufficient conditions for limit theorems in Gibbs-Markov maps*. Preprint (2008).
- [10] GOUËZEL S., LIVERANI C. *Banach spaces adapted to Anosov systems*. Ergodic Theory Dyn. Syst. **26**, 189-217 (2006).
- [11] GUIBOURG D. *Théorème de renouvellement pour chaînes de Markov fortement ergodiques. Applications aux modèles itératifs Lipschitziens*. C. R. Acad. Sci. Paris, Ser. I 346 (2008) 435-438.
- [12] GUIVARC'H Y., HARDY J. *Théorèmes limites pour une classe de chaînes de Markov et applications aux difféomorphismes d'Anosov*. Ann. Inst. Henri Poincaré, Vol. 24, No 1, p. 73-98 (1988).
- [13] GUIVARC'H Y., LE PAGE E. *On spectral properties of a family of transfer operators and convergence to stable laws for affine random walks*. Ergodic Theory Dynam. Systems, 28 (2008) no. 2, pp. 423-446.
- [14] HENNION H., HERVÉ L. *Limit theorems for Markov chains and stochastic properties of dynamical systems by quasi-compactness*. Lecture Notes in Mathematics No 1766, Springer (2001).
- [15] HENNION H., HERVÉ L. *Central limit theorems for iterated random lipschitz mappings*. Annals of Proba. Vol. 32 No. 3A (2004) 1934-1984.

- [16] HERVÉ L. *Théorème local pour chaînes de Markov de probabilité de transition quasi-compacte. Applications aux chaînes V -géométriquement ergodiques et aux modèles itératifs.* Ann. I. H. Poincaré - PR 41 (2005) 179-196.
- [17] HERVÉ L, PÈNE F. *The Nagaev method via the Keller-Liverani theorem.* To appear in Bull. Soc. Math. France. See arXiv:0901.4617.
- [18] KELLER G., LIVERANI C. *Stability of the Spectrum for Transfer Operators.* Ann. Scuola Norm. Sup. Pisa. Cl. Sci. (4) Vol. XXVIII (1999) 141-152.
- [19] LIVERANI C. *Invariant measures and their properties. A functional analytic point of view.* Dynamical Systems. Part II: Topological Geometrical and Ergodic Properties of Dynamics. Pubblicazioni della Classe di Scienze, Scuola Normale Superiore, Pisa. Centro di Ricerca Matematica "Ennio De Giorgi": Proceedings. Published by the Scuola Normale Superiore in Pisa (2004).
- [20] S.P. MEYN AND R.L. TWEEDIE. *Markov chains and stochastic stability.* Springer Verlag, New York, Heidelberg, Berlin (1993).
- [21] NAGAEV S.V. *Some limit theorems for stationary Markov chains.* Theory of probability and its applications 11 4 (1957) 378-406.
- [22] ROSENBLATT M. *Markov processes. Structure and asymptotic behavior.* Springer-Verlag, New York (1971).
- [23] SPITZER F. *Principles of random walks.* Van Nostrand, Princeton, 1964.