

DISCRETIZED CONFIGURATIONS AND PARTIAL PARTITIONS

AARON ABRAMS, DAVID GAY, AND VALERIE HOWER

ABSTRACT. We show that the discretized configuration space of k points in the n -simplex is homotopy equivalent to a wedge of spheres of dimension $n - k + 1$. This space is homeomorphic to the order complex of the poset of ordered partial partitions of $\{1, \dots, n + 1\}$ with exactly k parts. We also compute the Euler characteristic in two different ways, thereby obtaining a topological proof of a combinatorial recurrence satisfied by the Stirling numbers of the second kind.

1. INTRODUCTION

1.1. Configurations. The discretized configuration space $D_k(X)$ was introduced in [1] as a combinatorial model of the classical configuration space of k -tuples of distinct points in a space X . (When $k = 2$ this has classically been called the “deleted product.”) To define $D_k(X)$ it is required that X have the structure of a cell complex. In [1], and in the works of several subsequent authors, the space X is a graph, i.e. a finite 1-complex.

To study $D_k(X)$ for higher dimensional X , it is natural to begin with some basic building blocks. In this paper we consider the discretized configuration spaces $D_k(\Delta^n)$ where Δ^n is the n -dimensional simplex. We prove

Theorem 1. *The space $D_k(\Delta^n)$ is homotopy equivalent to a wedge of spheres of dimension $n - k + 1$.*

Theorem 2. *The number of spheres in the wedge is $\sum_{i=1}^{k-1} (-1)^{i+k+1} \binom{k}{i+1} i^{n+1}$.*

Note that if X is any simplicial complex then $D_k(X)$ is built out of (products of) spaces of the form $D_i(\Delta^n)$.

1.2. Partitions. The partition lattice Π_n is a classical combinatorial object studied since antiquity. It is a poset whose elements are set partitions of $[n] = \{1, \dots, n\}$ and whose ordering is given by refinement. The order complex of Π_n has the homotopy type of a wedge of spheres (see [4]). The (larger) lattice $\Pi_{\leq n}$ of *partial partitions* has recently also been shown to have the homotopy type of a wedge of spheres [2].

These results are proved using the machinery of algebraic combinatorics. The standard technique is to find a shelling of the order complex, often via an EL- or CL-labelling of the poset [4]. One nice feature of these arguments is that they often produce an explicit basis for the (unique) nonzero homology group, and one can usually compute its Betti number.

1.3. Connection. The face poset of the space $D_k(\Delta^{n-1})$ can be identified with a combinatorial object closely related to $\Pi_{\leq n}$, namely the poset $\tilde{\Pi}_{\leq n}^k$ of *ordered partial partitions of $[n]$ with exactly k parts*. Thus Theorems 1 and 2 imply the following.

Theorem 3. *The poset $\tilde{\Pi}_{\leq n}^k$ has the homotopy type of a wedge of spheres of dimension $n - k$. The number of spheres is the same as the expression given in Theorem 2, with n replaced by $n - 1$.*

Note that the symmetric group S_k acts on the space $D_k(X)$ (and on the poset $\tilde{\Pi}_{\leq n}^k$) by permuting coordinates. The quotient poset $\Pi_{\leq n}^k$ is naturally a sub-poset of $\Pi_{\leq n}$; its elements are the *partial partitions of $[n]$ with exactly k parts*. However, $\Pi_{\leq n}^k$ does not have the homotopy type of a wedge of spheres, as can be seen already in the case $k = 2$ (where one obtains a real projective plane $\mathbb{R}P^{n-2}$).

1.4. Paper contents. We start with definitions, examples and results (Section 2). The proof of Theorem 1 is a direct computation using algebraic topology. It takes two steps: in Section 3, we give an inductive argument that the spaces are simply connected (when the dimension at least 2), and then in Section 4 a spectral sequence computation shows that the spaces have the same homology as a wedge of spheres. Together, these allow us to use the Whitehead theorem to deduce the result.

In particular, we do not give a shelling of the spaces $D_k(\Delta^n)$, although we suspect that one may be possible.

Problem. *Find a combinatorial proof of Theorem 1.*

In Section 5 we describe two combinatorial proofs of Theorem 2. In the first, we use the interpretation as a partition lattice to count the i -dimensional cells in $D_k(\Delta^n)$. This count involves the Stirling numbers of the second kind, denoted

$$\left\{ \begin{matrix} N \\ K \end{matrix} \right\}$$

which (by definition) means the number of partitions of a set of size N into exactly K nonempty subsets.

We also obtain a recurrence for the top Betti number by following through the spectral sequence. One can prove that the expression in Theorem 2

agrees with the actual Betti numbers for small k , and satisfies the same recurrence; this establishes Theorem 2 in a different way.

2. DEFINITIONS, THEOREM, EXAMPLES

2.1. Configurations. Let $k, n \geq 1$ be fixed integers. The n -simplex Δ^n is the largest simplicial complex on the vertex set $[n + 1]$. The space $D_k(\Delta^n)$ is the largest cell complex that is contained in the product $(\Delta^n)^k$ minus its *diagonal* $\{(x_1, \dots, x_k) \in (\Delta^n)^k \mid x_i = x_j \text{ for some } i \neq j\}$. One can also describe this as the union of those open cells of $(\Delta^n)^k$ whose closure misses the diagonal; explicitly,

$$D_k(\Delta^n) = \bigcup_{\substack{\bar{\sigma}_i \text{ pairwise disjoint} \\ \text{closed cells in } \Delta^n}} \sigma_1 \times \cdots \times \sigma_k$$

We visualize this space by imagining k “robots” in the space Δ^n ; then $D_k(\Delta^n)$ is the space of allowable configurations if a robot is said to “occupy” the entire closure of the cell in whose interior it is contained, and a configuration is “allowable” if no two robots occupy the same point of Δ^n . The maximum dimension of a cell of $D_k(\Delta^n)$ is $n - k + 1$.

Theorem 1 (again). *For all $n \geq 1$ and $2 \leq k \leq n + 1$, the space $D_k(\Delta^n)$ is homotopy equivalent to a wedge of spheres of dimension $n - k + 1$.*

Proof. If $n - k + 1 = 0$ then $D_k(\Delta^n)$ is a discrete set of $k! \geq 2$ points, hence a wedge of 0-spheres. If $n - k + 1 = 1$ then $D_k(\Delta^n)$ is a connected 1-complex, hence up to homotopy, a wedge of circles. If $n - k + 1 > 1$ then $2 \leq k < n$, so Proposition 1 (Section 3) shows that $D_k(\Delta^n)$ is simply connected and Proposition 2 (Section 4) shows that $D_k(\Delta^n)$ has the homology of a wedge of spheres of dimension $n - k + 1$. As there is clearly a map from a wedge of spheres inducing isomorphisms on homology, the Whitehead theorem implies the result. \square

2.2. Examples. Let K_d denote the complete graph on the vertex set $[d]$ (i.e., the 1-skeleton of Δ^{d-1}). Let S^d denote the d -sphere. Let e_i denote the i th standard basis vector of \mathbb{R}^d , for $1 \leq i \leq d$.

Example 1 ($k = 2$). We view Δ^n as the convex hull of the $n + 1$ standard basis vectors e_i in \mathbb{R}^{n+1} . Then the *Gauss map* from $D_2(\Delta^n)$ to \mathbb{R}^{n+1} given by $(x, y) \mapsto x - y$ is a homeomorphism onto S , the boundary of the polytope with vertex set $\{e_i - e_j \mid i, j \in [n + 1], i \neq j\}$ in \mathbb{R}^{n+1} . To see this, note that the map is obviously continuous and surjective, and given a point in S one can determine the coordinates x and y of the preimage as the “positive” and “negative” parts of S . Thus the map is a bijection.

The complex S is contained in the hyperplane $\{\sum x_i = 0\} \cong \mathbb{R}^n$ and is homeomorphic to S^{n-1} . This agrees with Theorems 1 and 2. Note also that $D_2(\Delta^{n-1})$ is contained in $D_2(\Delta^n)$ as an equator.

We next describe some small cases.

$n = 2$. The space $D_2(\Delta^2)$ is a connected 1-complex with six vertices, each of valence two, and six edges. It is a hexagon. It is the same as $D_2(K_3)$, since neither robot may venture into the interior of the 2-cell of Δ^2 .

$n = 3$. The space $D_2(\Delta^3)$ is (the surface of) a cuboctahedron (Figure 1). Deleting the triangular 2-cells leaves the planar 2-complex $D_2(K_4)$.




FIGURE 1. $D_2(\Delta^3)$.

$n = 4$. The space $D_2(\Delta^4)$ is homeomorphic to S^3 . The space $D_2(K_5)$ is a subcomplex which is itself homeomorphic to a closed orientable surface of genus six. This surface is a Heegaard splitting of $D_2(\Delta^4)$. The complementary handlebodies are made of prisms ($\Delta^2 \times I$ and $I \times \Delta^2$) and 3-simplices ($v \times \Delta^3$ and $\Delta^3 \times v$).

Example 2 ($k = n$). In this case $D_k(\Delta^n)$ is a connected 1-complex; a simple count reveals that the rank of H_1 is $\frac{1}{2}(n-2)(n+1)! + 1$. It is true but not obvious that this agrees with Theorem 2. If one quotients by the action of the symmetric group S_k that permutes coordinates, the result is a graph whose first Betti number is $\binom{n}{2}$.

2.3. Partitions. An *ordered partition* π of the set $[n] = \{1, \dots, n\}$ is an r -tuple (for some r) of disjoint non-empty sets $\pi = (S_1, \dots, S_r)$ whose union is $[n]$. (The ordering is on the set of parts, but each part is an unordered set.) An *ordered partial partition* of $[n]$ is the same, except the union is only required to be a subset of $[n]$. An *ordered partial partition with exactly k parts* is an ordered partial partition with $r = k$.

Let $\tilde{\Pi}_{\leq n}^k$ be the poset whose elements are the ordered partial partitions of $[n]$ with exactly k parts. The partial order is given by $(S_1, \dots, S_k) \leq (T_1, \dots, T_k)$ iff $S_i \subseteq T_i$ for each i .

Note that $\tilde{\Pi}_{\leq n}^k$ is not a lattice: the meet $(S_i) \wedge (T_i)$ is $(S_i \cap T_i)$ if all these sets are nonempty, but it is otherwise nonexistent; and similarly the join $(S_i) \vee (T_i)$ is $(S_i \cup T_i)$ if all these sets are disjoint, but otherwise it does not exist. Of course we may add a top and bottom element if we like. For comparison, the poset of partial partitions $\Pi_{\leq n}$ (see [2]) has a top element consisting of the 1-part partition $[n]$, and it is a lattice provided one includes an empty partition at the bottom.

Proof of Theorem 3. The face poset of $D_k(\Delta^{n-1})$ is isomorphic to the poset $\tilde{\Pi}_{\leq n}^k$, by mapping the face $\sigma_1 \times \cdots \times \sigma_k$ to the element (S_i) where S_i is the set of vertices of the cell σ_i . Thus the order complex of $\tilde{\Pi}_{\leq n}^k$ is homeomorphic to $D_k(\Delta^{n-1})$, so this is equivalent to Theorems 1 and 2. \square

We remark once again that the quotient $\Pi_{\leq n}^k$ of $\tilde{\Pi}_{\leq n}^k$ by the symmetric group S_k does not have the homotopy type of a wedge of spheres. When $k = 2$, for example, the action is antipodal and the quotient is a projective plane. Nevertheless $\Pi_{\leq n}^k$ is a sub-poset of $\Pi_{\leq n}$, which is a wedge of spheres up to homotopy (see the introduction).

3. THE FUNDAMENTAL GROUP

This is the first of the propositions referred to in the proof of Theorem 1.

Proposition 1. *If $1 \leq k < n$ then $D_k(\Delta^n)$ is simply connected.*

Proof. If $k = 1$ then $D_k(\Delta^n) = \Delta^n$, which is simply connected. If $k = 2$ we have already seen that $D_2(\Delta^n)$ is homeomorphic to S^{n-1} , which is simply connected if $n > 2$. We proceed by induction on k ; let $k > 2$ be fixed. Note that the hypothesis means that if all robots are at vertices, then there are at least two unoccupied vertices.

To prove the theorem we will construct a set of generators of $\pi_1(D_k(\Delta^n))$ and then we will show that each is null-homotopic.

Since $D_k(\Delta^n) \subset (\Delta^n)^k$, projection onto the first factor induces a map $\rho : D_k(\Delta^n) \rightarrow \Delta^n$. The inverse image of a point in the interior of an i -cell of Δ^n is isomorphic to $D_{k-1}(\Delta^{n-i})$. In particular, if v is a vertex of Δ^n then $\rho^{-1}(v)$ is simply connected, by induction.

Let v be the vertex $n + 1$ of Δ^n and let T be the spanning tree of the 1-skeleton of Δ^n (that is, K_{n+1}) consisting of all edges incident with v . The space $\rho^{-1}(T)$ is the union of $n + 1$ *vertex spaces* (i.e. the preimages of the vertices), each of which is a copy of the simply connected space $D_{k-1}(\Delta^{n-1})$, and n *edge spaces* (preimages of edges), each of which is a copy of the connected (but not necessarily simply connected) space $I \times D_{k-1}(\Delta^{n-2})$. The edge spaces are attached to the vertex spaces by embeddings at the ends

$\{0, 1\} \times D_{k-1}(\Delta^{n-2})$. Thus by the Seifert-van Kampen theorem, $\rho^{-1}(T)$ is simply connected.

Now consider $Y = \rho^{-1}(K_{n+1})$. The space Y is obtained from $\rho^{-1}(T)$ by attaching $\binom{n}{2}$ edge spaces $I \times D_{k-1}(\Delta^{n-2})$ indexed by the pairs $i, j \in [n]$ with $i < j$. As there are no new vertex spaces, each such edge space results in an HNN extension of the fundamental group; thus $\pi_1(Y)$ is free of rank $\binom{n}{2}$.

Note that the entire 1-skeleton of $D_k(\Delta^n)$ is contained in Y . Thus a generating set for $\pi_1(Y)$ will also generate $\pi_1(D_k(\Delta^n))$. We now describe such a generating set.

Fix a basepoint $\star \in \rho^{-1}(v)$. For each $i, j \in [n]$ choose a path α_{ij} in $\rho^{-1}(v)$ from \star to a configuration x with i and j unoccupied and each robot at a vertex of Δ^n . Let γ_{ij} be the loop starting at x that leaves all robots fixed except the first, and moves the first robot around the triangle $v \rightarrow i \rightarrow j \rightarrow v$. The loop $\alpha_{ij}\gamma_{ij}\alpha_{ij}^{-1}$ represents the generator of $\pi_1(Y)$ arising from attaching the edge space $\rho^{-1}([i, j])$. Letting i, j vary, these $\binom{n}{2}$ loops form a free basis for $\pi_1(Y)$.

But clearly each of these generators of $\pi_1(Y)$ is null-homotopic in $D_k(\Delta^n)$, as the loop γ_{ij} bounds a 2-simplex in $D_k(\Delta^n)$. We conclude that $D_k(\Delta^n)$ is simply connected, as desired. \square

4. HOMOLOGY

Here we prove the second of the propositions referred to in the proof of Theorem 1, using a spectral sequence to compute the homology of $D_k(\Delta^n)$. All homology groups will have integer coefficients.

Recall or observe:

- (1) $D_k(\Delta^n)$ has dimension $n - k + 1$.
- (2) If $n - k + 1 > 0$ then $D_k(\Delta^n)$ is connected.
- (3) $D_2(\Delta^n)$ is homeomorphic to a sphere of dimension $n - 1$.
- (4) $D_k(\Delta^{k-1})$ is $k!$ points.

Let $n \geq 1$ and $1 \leq k \leq n$ be fixed.

Again we consider the projection $\rho : D_k(\Delta^n) \longrightarrow \Delta^n$ onto the first coordinate. Note that ρ satisfies

$$\begin{aligned} \rho^{-1}([i_1]) &\cong D_{k-1}(\Delta^{n-1}) \\ \rho^{-1}([i_1, i_2]) &\cong \mathbb{R} \times D_{k-1}(\Delta^{n-2}) \\ \rho^{-1}([i_1, i_2, i_3]) &\cong \mathbb{R}^2 \times D_{k-1}(\Delta^{n-3}) \\ &\vdots \\ \rho^{-1}([i_1, i_2, i_3, \dots, i_{n-k+1}]) &\cong \mathbb{R}^{n-k} \times D_{k-1}(\Delta^{k-1}) \\ \rho^{-1}([i_1, i_2, i_3, \dots, i_{n-k+2}]) &\cong \mathbb{R}^{n-k+1} \times D_{k-1}(\Delta^{k-2}) \end{aligned}$$

where $i_1, i_2, i_3, \dots, i_{n-k+2}$ are distinct vertices of Δ^n and the face $[\cdot]$ of Δ^n is the interior of the convex hull of the given vertices. We use $\Delta^n(k)$ to denote the k -dimensional faces of Δ^n and $\Delta_{\leq k}^n = \bigcup_{i \leq k} \Delta^n(i)$ for the k -skeleton of Δ^n .

The map ρ gives a filtration of $D_k(\Delta^n)$ as follows

$$\emptyset = X_{-1} \subset X_0 \subset X_1 \subset X_2 \subset \dots \subset X_{n-k} \subset X_{n-k+1} = D_k(\Delta^n)$$

where $X_p = \rho^{-1}(\Delta_{\leq p}^n)$. Moreover

$$\begin{aligned} X_p \setminus X_{p-1} &= \bigsqcup_{f \in \Delta^n(p)} \rho^{-1}(f) \\ &\cong \bigsqcup_{f \in \Delta^n(p)} \mathbb{R}^p \times D_{k-1}(\Delta^{n-p-1}) \end{aligned}$$

We can hence construct a spectral sequence [3, p 327] (E^r, d^r) with

$$E_{p,q}^1 = H_{p+q}(X_p \setminus X_{p-1}) \implies H_{p+q}(D_k(\Delta^n))$$

converging to homology with closed supports. Since $H_p(\mathbb{R}^p) = \mathbb{Z}$ is the only nonzero homology group of \mathbb{R}^p we have

$$\begin{aligned} E_{p,q}^1 &= H_{p+q}(X_p \setminus X_{p-1}) \\ &= \bigoplus_{f \in \Delta^n(p)} H_{p+q}(\mathbb{R}^p \times D_{k-1}(\Delta^{n-p-1})) \\ &= \bigoplus_{f \in \Delta^n(p)} H_q(D_{k-1}(\Delta^{n-p-1})), \end{aligned}$$

where we have used the Künneth formula.

Proposition 2. *Let $n \geq 1$ and $1 \leq k \leq n$. Then*

$$H_r(D_k(\Delta^n)) = \begin{cases} \mathbb{Z}, & \text{if } r = 0, \\ 0, & \text{if } 0 < r < n - k + 1, \end{cases}$$

and $H_{n-k+1}(D_k(\Delta^n))$ is free abelian and nontrivial. Thus $D_k(\Delta^n)$ has the same homology as a wedge of spheres of dimension $n - k + 1$.

Proof. We induct on k ; the cases $k = 1, 2$ are observations (2), (3) above. Assume the theorem holds for configurations of $k - 1$ robots. Let (E^r, d^r) be the spectral sequence from above. Using our inductive hypothesis, the E^1 term has nonzero entries only along the diagonal line $p + q = n - k + 1$ and along row $q = 0$. The entries in the E^1 term are as follows

$$\begin{array}{c|cccccccc}
 & E_{0,n-k+1}^1 & & & & & & & \\
 q & 0 & E_{1,n-k}^1 & & & & & & \\
 & 0 & 0 & E_{2,n-k-1}^1 & & & & & \\
 & 0 & 0 & 0 & E_{3,n-k-2}^1 & & & & \\
 & 0 & 0 & 0 & 0 & \ddots & & & \\
 & 0 & 0 & 0 & 0 & \cdots & \ddots & & \\
 & 0 & 0 & 0 & 0 & \cdots & \cdots & E_{n-k,1}^1 & \\
 & E_{0,0}^1 & E_{1,0}^1 & E_{2,0}^1 & E_{3,0}^1 & \cdots & \cdots & E_{n-k,0}^1 & E_{n-k+1,0}^1 \\
 \hline
 & p=0 & 1 & 2 & 3 & \cdots & \cdots & n-k & n-k+1
 \end{array}$$

where

$$\begin{aligned}
 \text{rank} E_{p,n-k+1-p}^1 &= \binom{n+1}{p+1} b_{n-k+1-p}(D_{k-1}(\Delta^{n-p-1})) \text{ for } 0 \leq p \leq n-k, \\
 \text{rank} E_{p,0}^1 &= \begin{cases} \binom{n+1}{p+1} & \text{for } 0 \leq p \leq n-k \\ (k-1)! \binom{n+1}{p+1} & \text{for } p = n-k+1, \end{cases}
 \end{aligned}$$

and all other entries are zero.

The only possible nonzero higher differentials are the horizontal maps $d_{p,0}^1 : E_{p,0}^1 \rightarrow E_{p-1,0}^1$ for $1 \leq p \leq n - k + 1$, where

$$d_{p,0}^1 : \bigoplus_{f \in \Delta^n(p)} H_0(D_{k-1}(\Delta^{n-p-1})) \longrightarrow \bigoplus_{g \in \Delta^n(p-1)} H_0(D_{k-1}(\Delta^{n-p}))$$

is a direct sum of maps

$$H_0(D_{k-1}(\Delta^{n-p-1})) \longrightarrow \bigoplus_{g \in f(p-1)} H_0(D_{k-1}(\Delta^{n-p})) \quad \text{for } f \in \Delta^n(p)$$

Note that if $\dim f \leq n - k$ and $g \in f(p - 1)$ then the map

$$\mathbb{Z} \cong H_0(D_{k-1}(\Delta^{n-p-1})) \longrightarrow H_0(D_{k-1}(\Delta^{n-p})) \cong \mathbb{Z}$$

is injective (hence an isomorphism) as it is induced by inclusion. Thus, computing $E_{p,0}^2 = \ker d_{p,0}^1 / \text{im} d_{p+1,0}^1$ is equivalent to computing the p th homology group of Δ^n for $p < n - k$, and we obtain $E_{p,0}^2 = 0$ for $p < n - k$.

For $p = n - k$, we have $d_{n-k,0}^1$ is equivalent to ∂_{n-k} , where ∂ is the boundary map for the n -simplex. For each $(n - k + 1)$ -face f of Δ^n , let γ_f be a generator for the factor of $H_0(D_{k-1}(\Delta^{k-2}))$ corresponding to the fiber over f . Then

$$d_{n-k+1,0}^1 \Big|_{\bigoplus_f \mathbb{Z} \cdot \gamma_f} \cong \partial_{n-k+1}$$

and hence $\text{im} d_{n-k+1,0}^1 \cong \text{im} \partial_{n-k+1} = \ker \partial_{n-k} \cong \ker d_{n-k,0}^1$, which yields $E_{n-k,0}^2 = 0$. Thus the only nonzero entries of the E^2 term are $E_{0,0}^2 \cong \mathbb{Z}$ and also along the line $p + q = n - k + 1$ where we have free abelian groups. We hence obtain the integer homology groups of $D_k(\Delta^n)$ from $E^2 = E^\infty$ by adding along the lines $p + q = r$, which proves the inductive step. \square

5. THE EULER CHARACTERISTIC

In this section we discuss the Euler characteristic of $D_k(\Delta^n)$ from two points of view: first via the interpretation of $D_k(\Delta^n)$ as the order complex of the poset $\tilde{\Pi}_{\leq n+1}^k$, and second via the spectral sequence computation from Section 4.

5.1. Stirling numbers. Recall that the symmetric group S_k acts on $D_k(\Delta^n)$ by permuting coordinates. This is a free cellular action, i.e. the quotient $UD_k(\Delta^n)$ inherits a natural cell structure. An i -dimensional cell of $D_k(\Delta^n)$ corresponds to an ordered partial partition of $[n + 1]$ which uses exactly $k + i$ of the elements from $[n + 1]$, and an i -dimensional cell of $UD_k(\Delta^n)$ corresponds to an *unordered* partial partition of $[n + 1]$ which uses exactly $k + i$ of the elements from $[n + 1]$.

The Stirling number of the second kind, denoted $S(n, k)$ or $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$, is by definition the number of ways to partition a set of size n into exactly k nonempty subsets. (These partitions are unordered.) Thus the number of i -cells of $UD_k(\Delta^n)$ is

$$\binom{n+1}{k+i} \left\{ \begin{matrix} k+i \\ k \end{matrix} \right\},$$

and the Euler characteristic of $UD_k(\Delta^n)$ is

$$\sum_{i=0}^{n-k+1} (-1)^i \binom{n+1}{k+i} \left\{ \begin{matrix} k+i \\ k \end{matrix} \right\}.$$

As the Euler characteristic is multiplicative under covers, it follows that the Euler characteristic of $D_k(\Delta^n)$ is

$$(1) \quad k! \sum_{i=0}^{n-k+1} (-1)^i \binom{n+1}{k+i} \left\{ \begin{matrix} k+i \\ k \end{matrix} \right\}.$$

There are many well-known formulae and recurrences for the Stirling numbers; for instance

$$\left\{ \begin{matrix} n \\ k \end{matrix} \right\} = \frac{1}{k!} \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} j^n.$$

This provides one avenue to a proof of Theorem 2. By Equation (1), that would follow from the combinatorial identity

$$\sum_{j=0}^k (-1)^{n+j-1} \binom{k}{j} (j-1)^{n+1} = k! \sum_{i=0}^{n-k+1} (-1)^i \binom{n+1}{k+i} \left\{ \begin{matrix} k+i \\ k \end{matrix} \right\},$$

which can indeed be proven by elementary combinatorial techniques.

Example 3 ($k = 3$). The preceding identity shows that $D_3(\Delta^n)$ is a wedge of $(2^{n+1} - 3)$ 2-spheres.

Example 4 ($k = 4$). Similarly, $D_4(\Delta^n)$ is a wedge of $(3^{n+1} - 4 \cdot 2^{n+1} + 6)$ 3-spheres.

5.2. Betti numbers from the spectral sequence. Another approach to computing the Euler characteristic is to notice that the spectral sequence yields a recursion satisfied by the top Betti number. One can then establish Theorem 2 by showing that the formula in the statement of the theorem satisfies the same recursion. Again, this can be carried out by elementary (but non-trivial) combinatorial arguments.

Let $\beta_{k,n}$ denote the rank of $H_{n-k+1}(D_k(\Delta^n))$. We establish the following recurrence for $\beta_{k,n}$ in terms of the values of $\beta_{k-1,i}$ for $i < n$.

Theorem 4. *For $1 \leq k \leq n + 1$, we have*

$$\beta_{k,n} = -\binom{n}{k-1} + \sum_{p=0}^{n-k+1} \binom{n+1}{k+p-1} \beta_{k-1,k+p-2}.$$

It would be nice to have a geometric interpretation of this result. For instance, it may be possible to describe bases for the top homology groups in such a way that geometric relationships between the bases for different values of k and n shed light on this recurrence.

In any case, the theorem itself follows immediately from the next two lemmas.

Lemma. *For $3 \leq k \leq n$, define $Y_{k,n} := \text{rank} E_{n-k+1,0}^2$ where (E^r, d^r) is the spectral sequence for $D_k(\Delta^n)$. Then, we have*

$$Y_{k,n} = (k-1)! \binom{n+1}{k-1} - \binom{n}{k-1}.$$

Proof. As in Section 4, for each $(n-k+1)$ -face f of Δ^n , let γ_f be a generator for the factor of $H_0(D_{k-1}(\Delta^{k-2}))$ corresponding to the fiber over f . Then we have the following

$$\begin{aligned} \operatorname{im} d_{n-k+1,0}^1 &= \operatorname{im} d_{n-k+1,0}^1 \Big|_{\bigoplus_f \mathbb{Z} \cdot \gamma_f} \\ \ker d_{n-k+1,0}^1 &= \underbrace{\mathbb{Z} \binom{n+1}{n-k+2} \oplus \dots \oplus \mathbb{Z} \binom{n+1}{n-k+2}}_{(k-1)!-1 \text{ factors}} \oplus \ker d_{n-k+1,0}^1 \Big|_{\bigoplus_f \mathbb{Z} \cdot \gamma_f} \end{aligned}$$

where

$$\ker d_{n-k+1,0}^1 \Big|_{\bigoplus_f \mathbb{Z} \cdot \gamma_f} \cong \ker \partial_{n-k+1} = \operatorname{im} \partial_{n-k+2}$$

and ∂ is the boundary map for the n -simplex. Next, we note that

$$\begin{aligned} \operatorname{rank}(\operatorname{im} \partial_{n-k+2}) &= \binom{n+1}{n-k+3} - \operatorname{rank}(\ker \partial_{n-k+2}) \\ &= \binom{n+1}{n-k+3} - \operatorname{rank}(\operatorname{im} \partial_{n-k+3}). \end{aligned}$$

Thus, if we define $f_k := \operatorname{rank}(\operatorname{im} \partial_{n-k+2})$ then we have $f_2 = 1$ and $f_k = \binom{n+1}{n-k+3} - f_{k-1}$ for $3 \leq k \leq n$. Iterating, we find $f_k = \sum_{j=0}^{k-2} (-1)^{k+j} \binom{n+1}{j}$.

Thus

$$\begin{aligned} Y_{k,n} &= [(k-1)! - 1] \binom{n+1}{k-1} + f_k \\ &= (k-1)! \binom{n+1}{k-1} + \sum_{j=0}^{k-1} (-1)^{k+j} \binom{n+1}{j} \\ &= (k-1)! \binom{n+1}{k-1} - \binom{n}{k-1} \end{aligned}$$

where in the last line we have used the elementary combinatorial identity $\sum_{j=0}^K (-1)^j \binom{N}{j} = (-1)^K \binom{N-1}{K}$. \square

Lemma. For $3 \leq k \leq n$, let $\beta_{k,n}$ be the rank of the homology group $H_{n-k+1}(D_k(\Delta^n))$. Then $\beta_{k,n}$ satisfies the following recursion:

$$\beta_{k,n} = Y_{k,n} + \sum_{p=0}^{n-k} \binom{n+1}{p+1} \beta_{k-1,n-p-1},$$

where $Y_{k,n}$ is as in the above lemma.

Proof. Since the spectral sequence (E^r, d^r) collapses at E^2 and there is no torsion in the E^2 -term, we have

$$\begin{aligned}\beta_{k,n} &= \text{rank}E_{n-k+1,0}^2 + \sum_{p=0}^{n-k} \text{rank}E_{p,n+1-k-p}^2 \\ &= Y_{k,n} + \sum_{p=0}^{n-k} \text{rank}E_{p,n+1-k-p}^1.\end{aligned}$$

Since $\text{rank}E_{p,n+1-k-p}^1 = \binom{n+1}{p+1}\beta_{k-1,n-p-1}$, the lemma holds. \square

REFERENCES

- [1] A. Abrams, “Configuration spaces of colored graphs,” *Geometriae Dedicata* 92 (2002), pp. 185–194.
- [2] P. Hanlon, P. Hersh, J. Shareshian, “A $GL_n(q)$ -analogue of the partition lattice,” preprint available at <http://www4.ncsu.edu/~plhersh/hhs09.pdf>.
- [3] S. MacLane, *Homology*, Springer-Verlag, Berlin, 1963.
- [4] M. Wachs, “Poset topology: tools and applications,” *IAS/Park City Mathematics Institute*, 2004.