

The homotopy type of toric arrangements

Luca MOCI* Simona SETTEPANELLA†

October 29, 2010

Abstract

A toric arrangement is a finite set of hypersurfaces in a complex torus, every hypersurface being the kernel of a character. In the present paper we build a CW-complex homotopy equivalent to the arrangement complement, with a combinatorial description similar to that of the well-known Salvetti complex. If the toric arrangement is defined by a Weyl group, we also provide an algebraic description, very handy for cohomology computations. In the last part we give a description in terms of tableaux for a toric arrangement appearing in robotics.

Keywords:

Hyperplane arrangements, toric arrangements, CW-complex, Salvetti complex, Weyl groups, Young tableaux

MSC (2010):

52C35, 20F55, 05E45, 55P15

Introduction

A toric arrangement is a finite set of hypersurfaces in a complex torus $T = (\mathbb{C}^*)^n$, in which every hypersurface is the kernel of a character $\chi \in X \subset \text{Hom}(T, \mathbb{C}^*)$ of T . Let \mathcal{R}_X be the complement of the arrangement: its geometry and topology have been studied by many authors, see for instance [9], [4], [3], [13]. In particular, in [10] and [3] the De Rham cohomology of \mathcal{R}_X has been computed, and recently in [15] a *wonderful model* has been built.

In the present paper we build a topological model \mathcal{S} for \mathcal{R}_X . This model is a regular CW-complex, similar to the one introduced by Salvetti ([17]) for the complement of hyperplane arrangements. Indeed for a wide class of arrangements, which we call *thick*, the cells of \mathcal{S} are in bijection with pairs $[C \prec F]$, where C is a *chamber* of the *real* toric arrangement and F is a *facet* adjacent to it (according to the definitions given in Section 1.3).

The model \mathcal{S} is well suited for homology and homotopy computations, which will be developed in future papers (see for instance [19]). Furthermore, the

*Institut fuer Mathematik, Technische Universitaet Berlin, Strasse des 17. Juni, 136, Berlin. moci@math.tu-berlin.de (This work was partially supported by a Sofia Kovalevskaya Research Prize of Alexander von Humboldt Foundation awarded to Olga Holtz.)

†LEM, Scuola Superiore Sant'Anna, Pisa, Italy. s.settepanella@sssup.it (Thanks to financial support from the European Commission 6th FP (Contract CIT3-CT-2005-513396), Project: DIME - Dynamics of Institutions and Markets in Europe)

jumping loci in the local system cohomology of a CW-complex are affine algebraic varieties. In the theory of hyperplane arrangements such objects, called *characteristic varieties*, proved to be of fundamental importance. It is then a remarkable fact that the characteristic varieties can be defined also in the toric case.

In Section 2 we focus on a toric arrangement associated to an affine Weyl group \widetilde{W} . In this case the chambers are in bijection with the elements of the corresponding finite Weyl group W , and the cells of \mathcal{S} are given by the pairs (w, Γ) , where $w \in W$ and Γ is a proper subset of the set S of generators of \widetilde{W} . This generalizes a construction introduced in [18] and [6].

In the last Section we give a description of the facets of the real toric arrangement defined by the Weyl group A_n in the torus corresponding to the root lattice. This description in terms of Young tableaux turns out to be interesting since it coincides with the complex describing the space of all periodic legged gaits of a robot body (see [2]).

Acknowledgements We are grateful to the organizers of the research program "Configuration Spaces: Geometry, Combinatorics and Topology" at Centro De Giorgi (Pisa), which provided us a significant occasion to work together. In particular we wish to thank Fred Cohen and Mario Salvetti for several valuable suggestions. We also thank Priyavrat Deshpande for many stimulating conversations we had while we were completing the present paper.

1 The CW-complex

1.1 Main definitions

Let $T = (\mathbb{C}^*)^n$ be a complex torus and $X \subset \text{Hom}(T, \mathbb{C}^*)$ be a finite set of characters of T . The kernel of every $\chi \in X$ is a hypersurface of T :

$$H_\chi := \{t \in T \mid \chi(t) = 1\}.$$

Then X defines on T the *toric arrangement*:

$$\mathcal{T}_X := \{H_\chi, \chi \in X\}.$$

Let \mathcal{R}_X be the *complement* of the arrangement:

$$\mathcal{R}_X := T \setminus \bigcup_{\chi \in X} H_\chi.$$

Let $\pi : V \rightarrow T$ be the universal covering of T . Then V is a complex vector space of rank n , and π is the quotient map $\pi : V \rightarrow V/\Lambda$, where Λ is a lattice in V . Then the preimage $\pi^{-1}(H_\chi)$ of a hypersurface $H_\chi \in \mathcal{T}_X$ is the union of an infinite family of parallel hyperplanes. Thus

$$\mathcal{A}_X := \{H \text{ hyperplane of } V \mid \exists \chi \in X \text{ s.t. } \pi(H) = H_\chi\}$$

is a periodic affine hyperplane arrangement in V . Let \mathcal{M}_X be its complement:

$$\mathcal{M}_X := V \setminus \bigcup_{\chi \in X} \pi^{-1}(H_\chi).$$

By definition, π maps \mathcal{M}_X on \mathcal{R}_X . Moreover the equations defining the hyperplanes in \mathcal{A}_X can always be assumed to have integral (hence real) coefficients since they are given by elements of Λ . Thus by [17] there is an (infinite) CW-complex $\tilde{\mathcal{S}} \subset \mathcal{M}_X$ and a map $\varphi : \mathcal{M}_X \rightarrow \tilde{\mathcal{S}}$ giving a homotopy equivalence. Furthermore, we can build $\tilde{\mathcal{S}}$ in such a way that it is invariant under the action of translation in Λ : for instance by building the cells relative to a fundamental domain and then inductively, defining for each cell above the other cells of its Λ -orbit by translation. Thus $\pi(\tilde{\mathcal{S}})$ is a finite CW-complex, which will be denoted by \mathcal{S} , and the image of every cell of $\tilde{\mathcal{S}}$ is a cell of \mathcal{S} . Moreover, since φ is Λ -equivariant, it is well defined the map

$$\varphi_\pi(t) := (\pi \circ \varphi)(\pi^{-1}(t))$$

which makes the following diagram commutative:

$$\begin{array}{ccc} \mathcal{M}_X & \xrightarrow{\varphi} & \tilde{\mathcal{S}} \\ \pi \downarrow & & \pi \downarrow \\ \mathcal{R}_X & \xrightarrow{\varphi_\pi} & \mathcal{S} \end{array} \quad (1)$$

Lemma 1.1 *The map φ_π is a homotopy equivalence between \mathcal{R}_X and \mathcal{S} .*

Proof. The map φ is a homotopy equivalence hence, by definition, there is a continuous map $\psi : \tilde{\mathcal{S}} \rightarrow \mathcal{M}_X$ such that $\psi\varphi$ is homotopic to the identity map $id_{\mathcal{M}_X}$ and $\varphi\psi$ is homotopic to $id_{\tilde{\mathcal{S}}}$. Namely, since $\tilde{\mathcal{S}}$ is a deformation retract, the homotopy inverse ψ is simply the inclusion map, which is clearly Λ -equivariant. Hence the map

$$\psi_\pi(t) := (\pi \circ \psi)(\pi^{-1}(t))$$

is well defined and makes the following diagram commutative:

$$\begin{array}{ccc} \tilde{\mathcal{S}} & \xrightarrow{\psi} & \mathcal{M}_X \\ \pi \downarrow & & \pi \downarrow \\ \mathcal{S} & \xrightarrow{\psi_\pi} & \mathcal{R}_X. \end{array} \quad (2)$$

Let $I = [0, 1]$ be the unit interval and $F : \mathcal{M}_X \times I \rightarrow \mathcal{M}_X$ be the continuous map such that $F(x, 0) = \psi(\varphi(x))$ and $F(x, 1) = id_{\mathcal{M}_X}(x)$. Again, since F is Λ -equivariant, we can define the map:

$$F_\pi(t) := (\pi \circ F)(\pi^{-1}(t))$$

In this way we get the commutative diagram:

$$\begin{array}{ccc} \mathcal{M}_X \times I & \xrightarrow{F} & \mathcal{M}_X \\ \pi \downarrow & & \pi \downarrow \\ \mathcal{R}_X \times I & \xrightarrow{F_\pi} & \mathcal{R}_X. \end{array} \quad (3)$$

By construction map F_π is a continuous map such that $F_\pi(x, 1) = id_{\mathcal{R}_X}$ and

$$F_\pi(x, 0) = (\psi\varphi)_\pi(x) = \pi\psi\varphi\pi^{-1}(x) = \pi\psi\pi^{-1}\pi\varphi\pi^{-1}(x) = \psi_\pi \circ \varphi_\pi(x).$$

Hence F_π gives the required homotopy equivalence. \square

1.2 Salvetti complex for affine arrangements

In order to describe the structure of \mathcal{S} , we now have to focus on the real counterparts of the complex arrangements above.

Let $V_{\mathbb{R}}$ be the real part of V . In other words, let $V_{\mathbb{R}} \doteq \mathbb{R}^n$ be a real vector space, and let $V \doteq V_{\mathbb{R}} \otimes_{\mathbb{R}} \mathbb{C}$ be its complexification. Then we identify $V_{\mathbb{R}}$ with a subspace of V via the map $v \mapsto v \otimes 1$.

Let $\mathcal{A}_{X,\mathbb{R}}$ be the corresponding hyperplane arrangement on $V_{\mathbb{R}}$ and $\mathcal{M}_{X,\mathbb{R}} = \mathcal{M}_X \cap V_{\mathbb{R}}$ its complement. Since the image of \mathbb{R} under the map $\mathbb{C} \rightarrow \mathbb{C}/\mathbb{Z} \xrightarrow{\sim} \mathbb{C}^*$ is the circle

$$S^1 := \{z \in \mathbb{C} \mid |z| = 1\}$$

we have that the image of $V_{\mathbb{R}}$ under the map $\pi : V \rightarrow V/\Lambda \xrightarrow{\sim} T$ is a compact torus $T_{\mathbb{R}} \subset T$. A *real toric arrangement* $\mathcal{T}_{X,\mathbb{R}}$ is naturally defined on $T_{\mathbb{R}}$ with hypersurfaces $H_{X,\mathbb{R}} := H_X \cap T_{\mathbb{R}}$ and complement $\mathcal{R}_{X,\mathbb{R}} = \mathcal{R}_X \cap T_{\mathbb{R}}$. Furthermore π restricts to universal covering map $\pi : V_{\mathbb{R}} \rightarrow T_{\mathbb{R}}$ and $\pi(\mathcal{M}_{X,\mathbb{R}}) = \mathcal{R}_{X,\mathbb{R}}$.

We recall the following definitions:

1. a *chamber* of $\mathcal{A}_{X,\mathbb{R}}$ is a connected component of $\mathcal{M}_{X,\mathbb{R}}$;
2. a *space* of $\mathcal{A}_{X,\mathbb{R}}$ is an intersection of elements in $\mathcal{A}_{X,\mathbb{R}}$;
3. a *facet* of $\mathcal{A}_{X,\mathbb{R}}$ is the intersection of a space and the closure of a chamber.

Let $\mathbf{S} := \{\tilde{F}^k\}$ be the stratification of $V_{\mathbb{R}}$ into facets \tilde{F}^k induced by the arrangement $\mathcal{A}_{X,\mathbb{R}}$ (see [1]), where superscript k stands for codimension.

Then the k -cells of the complex $\tilde{\mathcal{S}}$ described in [17] bijectively correspond to pairs

$$[\tilde{C} \prec \tilde{F}^k]$$

where $\tilde{C} = \tilde{F}^0$ is a chamber of \mathbf{S} and $\tilde{F}^i \prec \tilde{F}^j \Leftrightarrow \text{clos}(\tilde{F}^i) \supset \tilde{F}^j$ is the standard partial ordering in \mathbf{S} (see also [16]).

Let $|\tilde{F}|$ be the affine subspace spanned by \tilde{F} , and let us consider the subarrangement

$$\mathcal{A}_{\tilde{F}} = \{H \in \mathcal{A}_{X,\mathbb{R}} : \tilde{F} \subset H\}.$$

A cell $[\tilde{C} \prec \tilde{F}^k]$ is in the boundary of $[\tilde{D} \prec \tilde{G}^j]$ ($k < j$) if and only if

- i) $\tilde{F}^k \prec \tilde{G}^j$
 - ii) the chambers \tilde{C} and \tilde{D} are contained in the same chamber of $\mathcal{A}_{\tilde{F}^k}$.
- (4)

Previous conditions are equivalent to say that \tilde{C} is the chamber of $\mathcal{A}_{X,\mathbb{R}}$ which is the "closest" to \tilde{D} among those which contain \tilde{F}^k in their closure. The standard notation $[\tilde{C} \prec \tilde{F}^k] \in \partial_{\tilde{\mathcal{S}}}[\tilde{D} \prec \tilde{G}^j]$ will be used.

It is a simple remark that the above description of the Salvetti complex $\tilde{\mathcal{S}}$ is Λ -invariant. Indeed each translation $t \in \Lambda$ acts on the stratification $\mathbf{S} := \{\tilde{F}^k\}$ sending a k -facet F^k into the k -facet $t.F^k$. Then the translation t acts on $\tilde{\mathcal{S}}$ sending a k -cell $[C \prec F^k]$ in the k -cell $[t.C \prec t.F^k]$.

1.3 Salvetti Complex for toric arrangements

In order to give a similar description for \mathcal{S} , we introduce the following definitions:

1. a *chamber* of $\mathcal{T}_{X,\mathbb{R}}$ is a connected component of $R_{X,\mathbb{R}}$;
2. a *layer* of $\mathcal{T}_{X,\mathbb{R}}$ is a connected component of an intersection of elements of $\mathcal{T}_{X,\mathbb{R}}$;
3. a *facet* of $\mathcal{T}_{X,\mathbb{R}}$ is an intersection of a layer and the closure of a chamber.

Lemma 1.2

1. If \tilde{C} is a chamber of $\mathcal{A}_{X,\mathbb{R}}$, $\pi(\tilde{C})$ is a chamber of $\mathcal{T}_{X,\mathbb{R}}$;
2. If \tilde{L} is a space of $\mathcal{A}_{X,\mathbb{R}}$, $\pi(\tilde{L})$ is a layer of $\mathcal{T}_{X,\mathbb{R}}$;
3. If \tilde{F} is a facet of $\mathcal{A}_{X,\mathbb{R}}$, $\pi(\tilde{F})$ is a facet of $\mathcal{T}_{X,\mathbb{R}}$;

Proof. The first statement is clear, as well as the second one since $\pi(\tilde{L})$ must be connected. The third claim is a direct consequence of the previous two. \square

Now, let us consider pairs

$$[C \prec F^k]$$

where $C = F^0$ is a chamber of $\mathcal{T}_{X,\mathbb{R}}$, F^k a k -codimensional facet of $\mathcal{T}_{X,\mathbb{R}}$ and $F^i \prec F^j \Leftrightarrow \text{clos}(F^i) \supset F^j$.


By Lemma 1.2 the quotient map $\pi(\tilde{F})$ of a facet is still a facet in the real torus and, because of the surjectivity of π , we get that any facet F in $\mathcal{T}_{X,\mathbb{R}}$ is the image $F = \pi(\tilde{F})$ of an affine one.

We notice that

$$\pi([\tilde{C} \prec \tilde{F}]) = \pi([\tilde{D} \prec \tilde{G}]) \implies [\pi(\tilde{C}) \prec \pi(\tilde{F})] = [\pi(\tilde{D}) \prec \pi(\tilde{G})].$$

Indeed if $\pi([\tilde{C} \prec \tilde{F}]) = \pi([\tilde{D} \prec \tilde{G}])$ there is a translation $t \in \Lambda$ which sends $[\tilde{C} \prec \tilde{F}]$ in $[\tilde{D} \prec \tilde{G}]$. As a simple consequence $\tilde{D} = t.\tilde{C}$ and $\tilde{F} = t.\tilde{G}$, i.e. $\pi(\tilde{C}) = \pi(\tilde{D})$ and $\pi(\tilde{F}) = \pi(\tilde{G})$.

Then there is a natural surjective map from the cells of \mathcal{S} to the set of pairs $[C \prec F]$, but this map in general is not injective. Let us consider the simple example defined by $\mathcal{A} = \{x \in \mathbb{R} \mid x \in \mathbb{Z}\}$.



The chambers \tilde{C}_i for $i \in \mathbb{Z}$ are the open intervals $(i, i + 1)$ and the 1-codimensional facets are the points. The toric arrangement depends on the chosen lattice. For example we can quotient in two different way as in the above figure.

Namely, the picture on the left corresponds to the choice $\Lambda = \mathbb{Z}$, i.e. $\pi : x \mapsto e^{2\pi ix}$, whereas the picture on the right is given by $\Lambda = 2\mathbb{Z}$ and $\pi : x \mapsto e^{\pi ix}$. As shown in the pictures the complex in the former example cannot be described by the two pairs $[C_{-1} \prec C_{-1}]$, $[C_{-1} \prec e_0]$ since it has 3 cells. Furthermore, this CW-complex is not regular (the closure of its cells is not contractible). On the other hand, in the latter example we have a regular CW-complex with two 0-dimensional cells and four 1-dimensional cells.

Now we will focus on the case in which \mathcal{S} maps bijectively on the set of pairs $[C \prec F]$, since then the description of the complex \mathcal{S} is particularly striking. Since $\mathcal{S} = \pi(\tilde{\mathcal{S}})$ is a complex homotopic to the complement \mathcal{R}_X , \mathcal{S} is described by pairs of the form $[C \prec F]$ if and only if the map

$$\pi([\tilde{C} \prec \tilde{F}]) \longrightarrow [\pi(\tilde{C}) \prec \pi(\tilde{F})] \quad (5)$$

is injective.

Moreover, if the definition (5) holds then we can define the boundary of a pair $[C \prec F]$. We need first to introduce new notations.

Notations. Let $P_0 \subset V$ be a fundamental parallelogram for $\pi : V \rightarrow T$ containing the origin of V . Let $\mathcal{A}_{0,X}$ be the subarrangement of \mathcal{A}_X made by all the hyperplanes that intersect P_0 (see, for instance, figure (8) in the next Section).

We will say that a maximal dimensional cell $[\tilde{C} \prec \tilde{F}^n]$ is in $\mathcal{A}_{0,X}$ if its support $|\tilde{F}^n|$ is the intersection of some of the hyperplanes in $\mathcal{A}_{0,X}$. While a

k -cell $[\tilde{C} \prec \tilde{F}^k]$ is in $\mathcal{A}_{0,X}$ if it is in the boundary of a n -cell in $\mathcal{A}_{0,X}$. Let $\tilde{\mathcal{S}}_0$ be the set of all such cells.

With previous notations if (5) is injective (i.e. it is a bijection) we define the boundary as follow:

$[C \prec F^k]$ is in the boundary of $[D \prec G^j]$ ($k < j$) if and only if there are cells $[\tilde{C} \prec \tilde{F}^k] \in \pi^{-1}([C \prec F^k]) \cap \tilde{\mathcal{S}}_0$ and $[\tilde{D} \prec \tilde{G}^j] \in \pi^{-1}([D \prec G^j]) \cap \tilde{\mathcal{S}}_0$ such that $[\tilde{C} \prec \tilde{F}^k] \in \partial_{\tilde{\mathcal{S}}}[\tilde{D} \prec \tilde{G}^j]$.

Obviously this boundary map commutes with the one in $\tilde{\mathcal{S}}$ and we get that the map in (5) is a bijection of CW-complexes.

Toric arrangement for which \mathcal{S} is in bijection with pairs $[C \prec F]$ are easily characterized as follows.

Definition 1.3 A toric arrangement \mathcal{T}_X is thick if the quotient map

$$\pi : V \longrightarrow T$$

is injective on the closure $\text{clos}(\tilde{C})$ of every chamber \tilde{C} of the associated affine arrangement $\mathcal{A}_{X,\mathbb{R}}$.

We notice that every toric arrangement is covered by a thick one and the fiber of the covering map is finite; hence our assumption is not very restrictive.

We have the following

Lemma 1.4 A toric arrangement \mathcal{T}_X is thick if and only if

$$[\pi(\tilde{C}) \prec \pi(\tilde{F})] = [\pi(\tilde{D}) \prec \pi(\tilde{G})] \iff \pi([\tilde{C} \prec \tilde{F}]) = \pi([\tilde{D} \prec \tilde{G}])$$

for any two cells $[\tilde{C} \prec \tilde{F}], [\tilde{D} \prec \tilde{G}] \in \tilde{\mathcal{S}}$

Proof. By previous considerations, it is enough to prove that the thick condition is equivalent to

$$[\pi(\tilde{C}) \prec \pi(\tilde{F})] = [\pi(\tilde{D}) \prec \pi(\tilde{G})] \implies \pi([\tilde{C} \prec \tilde{F}]) = \pi([\tilde{D} \prec \tilde{G}])$$

\implies : Let \mathcal{T}_X be thick and $[\pi(\tilde{C}) \prec \pi(\tilde{F})] = [\pi(\tilde{D}) \prec \pi(\tilde{G})]$ for two given k -cells in $\tilde{\mathcal{S}}$. This implies that $\pi(\tilde{C}) = \pi(\tilde{D})$ and $\pi(\tilde{F}) = \pi(\tilde{G})$, i.e. there are translations $t, t' \in \Lambda$ such that $\tilde{D} = t.\tilde{C}$ and $\tilde{G} = t'.\tilde{F}$.

By construction $t.\tilde{F}$ is a facet in the closure $\text{clos}(D)$. We get two facets $t.\tilde{F}$ and \tilde{G} both in $\text{clos}(D)$ and with the same image $\pi(t.\tilde{F}) = \pi(\tilde{F}) = \pi(\tilde{G})$. By hypothesis π is injective on $\text{clos}(D)$ then $t.\tilde{F} = \tilde{G}$, i.e. $t = t'$ which implies that $\pi([\tilde{C} \prec \tilde{F}]) = \pi([\tilde{D} \prec \tilde{G}])$.

\Leftarrow Let \tilde{F} and \tilde{G} two facets in $\text{clos}(\tilde{C})$ such that $\pi(\tilde{F}) = \pi(\tilde{G})$ then

$$\pi([\tilde{C} \prec \tilde{F}]) = [\pi(\tilde{C}) \prec \pi(\tilde{F})] = [\pi(\tilde{C}) \prec \pi(\tilde{G})] = \pi([\tilde{C} \prec \tilde{G}]).$$

As a consequence if $t \in \Lambda$ is the translation such that $\tilde{F} = t.\tilde{G}$ then $t.\tilde{C} = \tilde{C}$. It follows that t is the identity and we get $\tilde{F} = \tilde{G}$, i.e. π is injective on $\text{clos}(\tilde{C})$
 \square

By Lemma 1.4 the map defined in (5) is a bijection if and only if \mathcal{T}_X is a thick toric arrangement. Hence the set of pairs $[C \prec F]$ is a CW-complex $\overline{\mathcal{S}}$ and we get the following theorem

Theorem 1 *Let \mathcal{T}_X be a thick toric arrangement. Then its complement \mathcal{R}_X has the same homotopy type of the CW-complex $\overline{\mathcal{S}}$.*

Then in this case the complex \mathcal{S} has a nice combinatorial description, totally analogue to that of the classical Salvetti complex [17].

Moreover if a toric arrangement is thick then the maximal dimensional cells $[\tilde{C} \prec \tilde{F}^n]$ in $\mathcal{A}_{0,X}$ are in one to one correspondence with the n -dimensional facets of $\overline{\mathcal{S}}$. Then the boundary in a thick toric arrangement \mathcal{T}_X can be completely described knowing the boundary in the associated finite complex $\mathcal{A}_{0,X}$.

This allows to better understand the fundamental group of the complement and to perform computations on integer cohomology.

Furthermore, in this case \mathcal{S} is a *regular* CW-complex.

Remark 1.5 *The number of chambers of $\mathcal{T}_{X,\mathbb{R}}$ can be computed by formulae given in [8] and [13]. However, the combinatorics of the layers in $\mathcal{T}_{X,\mathbb{R}}$ is more complicated than the one of spaces of $\mathcal{A}_{X,\mathbb{R}}$. Hence, an enumeration of the facets is not easy to provide in the general case. Thus from now on we focus on the arrangements defined by roots systems. In this case the chambers are parametrized by the elements of the Weyl group, and the poset of layers has been described in [12].*

2 Weyl toric arrangements

In this section we give a simpler description of the above complex for the case of toric arrangements associated to affine Weyl groups, by taking as Λ the coroot lattice (for the theory of Weyl groups see, for instance, [1]). Indeed in this case the toric arrangement is thick. Using this description, we give an example of how the integer cohomology of these arrangements can be computed.

2.1 Notations and Recalls.

Toric arrangement associated to a Weyl group. Let Φ be a root system, $\Lambda = \langle \Phi^\vee \rangle$ be the lattice spanned by the coroots, and Δ be its dual lattice (which is called the *cocharacters* lattice). Then we define a torus $T = T_\Delta$ having Δ as group of characters. Namely, if \mathfrak{g} is the semisimple complex Lie algebra associated to Φ and \mathfrak{h} is a Cartan subalgebra, T is defined as the quotient $T \doteq \mathfrak{h}/\Lambda$.

Each root α takes integer values on Δ , so it induces a map

$$e^\alpha : T \rightarrow \mathbb{C}/\mathbb{Z} \simeq \mathbb{C}^*$$

which is a character of the torus. Let X be the set of these characters; more precisely, since α and $-\alpha$ define the same hypersurface, we set

$$X \doteq \{e^\alpha, \alpha \in \Phi^+\}.$$

In this way to every affine Weyl group \widetilde{W} we associate a toric arrangement $\mathcal{T}_{\widetilde{W}}$, with complement $\mathcal{R}_{\widetilde{W}}$.

We will call these arrangements Weyl toric arrangements. They have been studied in [11] and [12].

Remark 2.1

1. Let G be the semisimple, simply connected linear algebraic group associated to \mathfrak{g} . Then T is the maximal torus of G corresponding to \mathfrak{h} , and \mathcal{R}_X is known as the set of regular points of T .
2. One may take as Δ the root lattice (or equivalently, take as Λ the character lattice). But in this way one obtains as T a maximal torus of the semisimple adjoint group G^a , which is the quotient of G by its center.

Let (\widetilde{W}, S) be the Coxeter system associated to \widetilde{W} and

$$\mathcal{A}_{\widetilde{W}} = \{H_{\widetilde{w}s_i\widetilde{w}^{-1}} \mid \widetilde{w} \in \widetilde{W} \text{ and } s_i \in S\}$$

the arrangement in \mathbb{C}^n obtained by complexifying the reflection hyperplanes of \widetilde{W} , where, in a standard way, the hyperplane $H_{\widetilde{w}s_i\widetilde{w}^{-1}}$ is the hyperplane fixed by the reflection $\widetilde{w}s_i\widetilde{w}^{-1}$.

We can view Λ as a subgroup of \widetilde{W} , acting by translations. Then it is well known that $\widetilde{W}/\Lambda \simeq W$, where W is the finite reflection group associated to \widetilde{W} (see for instance [21]). As a consequence, the toric arrangement can be described as:

$$\mathcal{T}_{\widetilde{W}} = \{H_{[w]s_i[w^{-1}]} \mid w \in W \text{ and } s_i \in S\}$$

where two hypersurfaces $H_{[w]s_i[w^{-1}]}$ and $H_{[\overline{w}]s_i[\overline{w}^{-1}]}$ are equal if and only if there is a translation $t \in \Lambda$ such that $tw s_i (tw)^{-1} = \overline{w} s_i \overline{w}^{-1}$, i.e. $\overline{w} = tw$.

By [12], these hypersurfaces intersect in

$$\frac{|W|}{|W_{S \setminus \{s_i\}}|}$$

local copies of the finite hyperplane arrangement $A_{W_{S \setminus \{s_i\}}}$ associated to the group generated by $S \setminus \{s_i\}$, $s_i \in S$.

For example in the affine Weyl group \widetilde{A}_n generated by $\{s_0, \dots, s_n\}$ for any generator s_i the finite reflection group associated to $S \setminus \{s_i\}$ is a copy of the finite Coxeter group A_n .

Then we have

Proposition 2.2 *The toric arrangement $\mathcal{T}_{\widetilde{W}}$ is thick.*

Proof. Since Λ is the coroot lattice, if $t \in \Lambda$ is a translation such that there is a n -codimensional facet $\widetilde{F}^n \in \text{clos}(\widetilde{C}) \cap \text{clos}(t\widetilde{C})$ for an affine chamber \widetilde{C} , then t is the identity (see [1]).

If $\mathcal{T}_{\widetilde{W}}$ is not thick then there are two facets \widetilde{F}_1 and \widetilde{F}_2 in the closure $\text{clos}(\widetilde{C})$ of a chamber \widetilde{C} such that $\pi(\widetilde{F}_1) = \pi(\widetilde{F}_2)$, i.e. there is a translation $t \in \Lambda$ such that $\widetilde{F}_2 = t\widetilde{F}_1$. Hence \widetilde{F}_2 is a facet in $\text{clos}(\widetilde{C}) \cap \text{clos}(t\widetilde{C})$. In particular all the n -codimensional facets \widetilde{F}^n in the closure of \widetilde{F}_2 are in the closure of both \widetilde{C} and $t\widetilde{C}$. This is a contradiction and it concludes the proof. \square

Then we can construct the Salvetti complex for these arrangements in a way which is very similar to the one known for affine Coxeter arrangements.

Salvetti Complex for affine Artin groups It is well known (see, for instance, [6], [18]) that the cells of Salvetti complex $\widetilde{\mathcal{S}}_W$ for arrangements $\mathcal{A}_{\widetilde{W}}$ are of the form $E(\widetilde{w}, \Gamma)$ with $\Gamma \subset S$ and $\widetilde{w} \in \widetilde{W}$. Indeed if $\widetilde{\alpha} \in \{\widetilde{w}s\widetilde{w}^{-1} | s \in S, \widetilde{w} \in \widetilde{W}\}$ is a reflection, the chambers are in one to one correspondence with the elements of the group \widetilde{W} as follows. Fixed a base chamber C_0 , it corresponds to $1 \in \widetilde{W}$. Now if C corresponds to \widetilde{w} , the chamber D separated from C by the reflection hyperplane $H_{\widetilde{\alpha}}$ corresponds to the element $\widetilde{\alpha}\widetilde{w} \in \widetilde{W}$. The notation $D \simeq \widetilde{\alpha}\widetilde{w}$ will be used.

If \widetilde{F}^k is a k -codimensional facet then the k -cell $[\widetilde{C} \prec \widetilde{F}^k]$ corresponds to the pair $E(\widetilde{w}, \Gamma)$ where $\widetilde{w} \simeq \widetilde{C}$ and $\Gamma = \{s_{i_1}, \dots, s_{i_k}\}$ is the unique subset of cardinality k in S such that

$$|F^k| = \bigcap_{j=1}^k H_{\widetilde{w}s_{i_j}\widetilde{w}^{-1}}.$$

If \widetilde{W}_Γ is the finite subgroup generated by $s \in \Gamma$, by [6] the integer boundary map can be expressed as follows:

$$\partial_k(E(\widetilde{w}, \Gamma)) = \sum_{s_j \in \Gamma} \sum_{\beta \in \widetilde{W}_\Gamma^{\Gamma \setminus \{s_j\}}} (-1)^{l(\beta) + \mu(\Gamma, s_j)} E(\widetilde{w}\beta, \Gamma \setminus \{s_j\}). \quad (6)$$

where $\widetilde{W}_\Gamma^{\Gamma \setminus \{\sigma\}} = \{w \in \widetilde{W}_\Gamma : l(ws) > l(w) \forall s \in \Gamma \setminus \{\sigma\}\}$ and $\mu(\Gamma, s_j) = \#\{s_i \in \Gamma | i \leq j\}$.

Remark 2.3 *Instead of the co-boundary operator we prefer to describe its dual, i.e. we define the boundary of a k -cell $E(\widetilde{w}, \Gamma)$ as a linear combination of the $(k-1)$ -cells which have $E(\widetilde{w}, \Gamma)$ in their co-boundary, with the same coefficient of the co-boundary operator. We make this choice since the boundary operator has a nicer description than co-boundary operator in terms of the elements of \widetilde{W} .*

2.2 Description of the complex

Let \mathcal{S}_W be the CW-complex associated to $\mathcal{T}_{\widetilde{W}}$. By the previous considerations, \mathcal{S}_W admits a description similar to that of $\widetilde{\mathcal{S}}_W$. Indeed each chamber C is in one to one correspondence with an equivalence class $[w] \in \widetilde{W}/\Lambda$ and then with an element $w \in W \simeq \widetilde{W}/\Lambda$ of the finite reflection group W . We will write $C \simeq [w]$.

In the same way, the pair $[C \prec F^k]$ corresponds to the cell $E([w], \Gamma) \in \mathcal{S}_W$ where $C \simeq [w]$ and $\Gamma = \{s_{i_1}, \dots, s_{i_k}\}$ is the unique subset of cardinality k in S such that

$$|F^k| = \bigcap_{j=1}^k H_{[w]s_{i_j}[w^{-1}]}$$

We now want to describe the boundary of each cell: this is done in a standard way by characterizing the cells that are in the boundary of a given cell, and by assigning an orientation to all cells (see, for instance, [18]).

By construction the toric CW-complex is locally isomorphic to the affine one and it can inherit its affine orientation. Then the integer boundary operator for

Weyl toric arrangements can be written as the affine one:

$$\partial_k(E([w], \Gamma)) = \sum_{\sigma \in \Gamma} \sum_{\beta \in W_{\Gamma}^{\Gamma \setminus \{\sigma\}}} (-1)^{l(\beta) + \mu(\Gamma, \sigma)} E([w\beta], \Gamma \setminus \{\sigma\}) \quad (7)$$

where, instead of elements of the affine group \widetilde{W} , we have equivalence classes with representatives in the finite group W .

By the formula above, the complex \mathcal{S}_W can be effectively used for computing homotopy invariants of $\mathcal{R}_{\widetilde{W}}$. For instance we have

Proposition 2.4

$$H^{\bullet}(\mathcal{R}_{\widetilde{W}}, \mathbb{Z}) \simeq H^{\bullet}(\mathcal{S}_W, \mathbb{Z})$$

where the coboundary map is the dual of the map defined in (7).


Example. Let us consider the affine Weyl group \widetilde{B}_2 (see [1]) with Coxeter-Dynkin diagram

$$\begin{array}{ccccc} \circ & \frac{4}{-} & \circ & \frac{4}{-} & \circ \\ s_0 & & s_1 & & s_2 \end{array}$$

and associated finite group B_2

$$\begin{array}{ccc} \circ & \frac{4}{-} & \circ \\ s_1 & & s_2 \end{array}$$

In this case we get translations $t_1 = s_0 s_1 s_2 s_1$ and $t_2 = s_2 s_1 s_0 s_1$ and the affine arrangement is represented as:



and it gives rise to a matrix of 24 columns and 8 rows with entries 0, 1 and -1 . On the other hand, the second boundary map is given by

$$\begin{aligned} \partial_2 E([w], \{s_i, s_{i+1}\}) &= E([w], \{s_i\}) - E([ws_{i+1}], \{s_i\}) + E([ws_i s_{i+1}], \{s_i\}) - \\ &\quad - E([w], \{s_{i+1}\}) + E([ws_i], \{s_{i+1}\}) - E([ws_{i+1} s_i], \{s_{i+1}\}) \end{aligned}$$

$$\partial_2 E([w], \{s_0, s_2\}) = E([w], \{s_0\}) - E([ws_2], \{s_0\}) - E([w], \{s_2\}) + E([ws_0], \{s_2\}).$$

In this way we get that the homology, and hence the cohomology, is torsion free and $H_0(R_{B_2}, \mathbb{Z}) = \mathbb{Z}$, $H_1(R_{B_2}, \mathbb{Z}) = \mathbb{Z}^8$ and $H_2(R_{B_2}, \mathbb{Z}) = \mathbb{Z}^{15}$, which agrees with the Betti numbers computed in [12, Ex. 5.14].

In general we have the following

Conjecture 2.5 *Let \widetilde{W} be an affine Weyl group and $\mathcal{T}_{\widetilde{W}}$ be the corresponding toric arrangement. Then the integer cohomology of the complement is torsion free (and hence it coincides with the De Rham cohomology computed in [3]).*

This conjecture will be proved in a future paper [19].

3 An example from robotics

In this section we give an example of non-thick arrangement: the one obtained from the affine Weyl arrangement $\mathcal{A}_{\widetilde{A}_n}$, by quotienting by the coroot lattice, which we will denote by $\Lambda_{\widetilde{A}_n}$ (see the second part of Remark 2.1).

Indeed in this case the underlying real toric arrangement has a very nice description in terms of Young tableaux. More precisely the facets of $\mathcal{T}_{\widetilde{A}_n, \mathbb{R}}$ are in one to one correspondence with a family of Young tableaux which turn out to be the same tableaux describing the space of all periodic legged gaits of a robot body (see [2]).

It is clear that, in this case, the finite arrangement $\mathcal{A}_{0, \widetilde{A}_n}$ is exactly the braid arrangement \mathcal{A}_{A_n} .

3.1 Tableaux description for the complex $\widetilde{\mathcal{S}}_{A_n}$

We indicate by A_n the symmetric group on $n+1$ elements, acting by permutations of the coordinates. Then $\mathcal{A} = \mathcal{A}_{A_n}$ is the braid arrangement and $\widetilde{\mathcal{S}}_{A_n}$ is the associated CW-complex (even if the arrangement is finite we continue to use the same notation used above for the affine case to distinguish it from the toric one).

Given a system of coordinates in \mathbb{R}^{n+1} , we describe $\widetilde{\mathcal{S}}_{A_n}$ through certain tableaux as follow.

Every k -cell $[\widetilde{C} \prec \widetilde{F}]$ is represented by a tableau with $n+1$ boxes and $n+1-k$ rows (aligned on the left), filled with all the integers in $\{1, \dots, n+1\}$. There is no monotony condition on the lengths of the rows. One has:

- (x_1, \dots, x_{n+1}) is a point in \widetilde{F} if and only if:

1. i and j belong to the same row if and only if $x_i = x_j$,
2. i belongs to a row preceding the one containing j if and only if $x_i < x_j$;

- the chamber \tilde{C} belongs to the half-space $x_i < x_j$ if and only if:

1. either the row which contains i is preceding the one containing j or
2. i and j belong to the same row and the column which contains i is preceding the one containing j .

Notice that the facets of the real stratification are represented by standard Young tableaux, since the order of the entries in each row does not matter, and hence we can assume it to be strictly increasing.

Notice also that the geometrical action of A_n on the stratification induces a natural action on the complex \tilde{S}_{A_n} which, in terms of tableaux, is given by a left action of A_n : $\sigma \cdot T$ is the tableau with the same shape as T , and with entries permuted by σ .

3.2 Tableaux description for the facets of $\mathcal{T}_{\tilde{A}_n, \mathbb{R}}$

Let $\mathcal{A}_{0, \tilde{A}_n} \subset \mathcal{A}_{\tilde{A}_n}$ be the braid arrangement passing through the origin and $\pi : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1} / \Lambda_{\tilde{A}_n} = T_{\mathbb{R}}$ the projection map.

If $\mathbf{F}_{\tilde{A}_n}$ is the stratification of \mathbb{R}^{n+1} into facets induced by the arrangement $\mathcal{A}_{\tilde{A}_n}$, we define the set:

$$\mathbf{F}_{0, \tilde{A}_n} = \{\tilde{F}^k \in \mathbf{F}_{\tilde{A}_n} \mid \text{clos}(F^k) \supset \bigcap_{H \in \mathcal{A}_{0, \tilde{A}_n}} H\}.$$

Obviously $\mathbf{F}_{0, \tilde{A}_n}$ is in one to one correspondence with the stratification \mathbf{F}_{A_n} induced by the braid arrangement \mathcal{A}_{A_n} and the restriction $\pi_{\mathbf{F}_{0, \tilde{A}_n}}$ is surjective on $T_{\mathbb{R}}$.

It follows that in order to understand how $\Lambda_{\tilde{A}_n}$ acts on $\mathbf{F}_{\tilde{A}_n}$ it is enough to study how it acts on $\mathbf{F}_{0, \tilde{A}_n}$. Moreover it is enough to consider facets in the closure of the base chamber \tilde{C}_0 corresponding to $1 \in \tilde{A}_n$; the action on the others will be obtained by symmetry.

Let us remark that a facet \tilde{F}^k is in $\mathbf{F}_{0, \tilde{A}_n}$ if and only if it intersects any ball B_0 around the origin. Let B_0 be a ball of sufficiently small radius and

$$x = (x_1, \dots, x_{n+1}) \in \text{clos}(\tilde{C}_0) \cap B_0$$

be a given point in a facet $\tilde{F}^k \in \mathbf{F}_{0, \tilde{A}_n}$. Then the x_i 's satisfy $x_1 \leq x_2 \leq \dots \leq x_{n+1}$ and the standard Young tableaux $Tb_{\tilde{F}^k}$ associated to \tilde{F}^k will have entries increasing along both, rows and columns.

Let $t_1, \dots, t_n \in \Lambda_{\tilde{A}_n}$ be a base such that t_i translates the reflection hyperplane $H_{i, i+1} = \text{Ker}(x_i - x_{i+1})$ fixing all hyperplanes $H_{j, j+1} = \text{Ker}(x_j - x_{j+1})$ for $j \neq i$ (i.e. each point in $H_{j, j+1}$ is sent in a point still in $H_{j, j+1}$). Then we can assume that translation t_i acts on the entry x_i as $t_i.x_i = x_i + t$ with $x_i + t > x_{i+1}$ and, as $H_{j, j+1}$, for $j \neq i$, are invariant under the action of t_i , it follows that $t_i.x_{i-1} = x_{i-1} + t$ and, by induction, $t_i.x_j = x_j + t$ for all $j < i$, while $t_i.x_j = x_j$ for all $j > i$.

Recall that, by construction, given a standard Young tableaux, a point (x_1, \dots, x_{n+1}) is a point in \tilde{F} if and only if:

1. i and j belong to the same row if and only if $x_i = x_j$,
2. i belongs to a row preceding the one containing j if and only if $x_i < x_j$;

It follows that if Tb is a tableau such that $i \in r_k$ and $i + 1 \in r_{k+1}$ are in two different rows, then t_i acts on Tb sending it in a tableau Tb' with rows $r'_1 = r_{k+1}, \dots, r'_{h-k} = r_h, r'_{h-k+1} = r_1, \dots, r'_h = r_k$. While if $i, i + 1 \in r_k$ are in the same row, then t_i acts sending the corresponding facet in a facet which is not anymore in $\mathcal{A}_{0, \tilde{A}_n}$.

Then $\Lambda_{\tilde{A}_n}$ acts on the h rows of a tableau $Tb_{\tilde{F}}$ as a power of the cyclic permutation $(1, \dots, h)$.

Equivalently let $Y(n + 1, k + 1)$ be the set of standard Young tableaux with $k + 1$ rows and $n + 1$ entries and $Tb \in Y(n + 1, k + 1)$ be a tableau of rows (r_1, \dots, r_{k+1}) . Then we have the following proposition.

Proposition 3.1 *The set of facets F^k of the toric arrangement $\mathcal{T}_{\tilde{A}_n, \mathbb{R}}$ is in one to one correspondence with the set*

$$Y(n + 1, k + 1) / \sim$$

where a tableau $Tb' \sim Tb$ if and only if the rows of Tb' are $(r_{\sigma^s(1)}, \dots, r_{\sigma^s(k+1)})$ for a power σ^s of the cyclic permutation $\sigma = (1, \dots, k + 1)$.

In this way we get exactly the tableaux described in [2].

Finally let us recall that the relation $\tilde{F}^k \prec \tilde{F}^{k+1}$ holds if and only if the tableau $Tb_{\tilde{F}^{k+1}}$ corresponding to \tilde{F}^{k+1} is obtained by attaching two consecutive rows of $Tb_{\tilde{F}^k}$.

As a consequence if F^k and F^{k+1} are facets in the toric arrangement $\mathcal{T}_{\tilde{A}_n, \mathbb{R}}$, $F^k \prec F^{k+1}$ if and only if the tableau $Tb_{F^{k+1}}$ corresponding to F^{k+1} is obtained by attaching two consecutive rows of Tb_{F^k} or attaching the first one to the last one.

References

- [1] N. BOURBAKI, *Groupes et algèbres de Lie (chap. 4,5,6)*, Masson, Paris, 1981.
- [2] F. COHEN, G. C. HAYNES, D.E. KODITSCHKEK, *Gait Transitions for Quasi-Static Hexapedal Locomotion on Level Ground*, International Symposium of Robotics Research, (2009), Lucerne, Switzerland.
- [3] C. DE CONCINI, C. PROCESI, *On the geometry of toric arrangements*, Transformations Groups 10, (2005), 387-422.
- [4] C. DE CONCINI, C. PROCESI, *Topics in hyperplane arrangements, polytopes and box-splines*, Springer 2010.

- [5] C. DE CONCINI, C. PROCESI, AND M. SALVETTI, *Arithmetic properties of the cohomology of braid groups*, *Topology* 40 (2001), 739–751.
- [6] C. DE CONCINI AND M. SALVETTI, *Cohomology of Coxeter groups and Artin groups*, *Math. Res. Lett.*, **7** (2000), 213-232.
- [7] PRIYAVRAT DESHPANDE, *Arrangements of submanifolds and the tangent bundle complement*, in preparation.
- [8] R. EHRENBORG, M. READDY, M. SLONE, *Affine and toric hyperplane arrangements*, arXiv:0810.0295v1 (math.CO), 2008.
- [9] G. I. LEHRER, *The cohomology of the regular semisimple variety*, *J. Algebra* 199 (1998), no. 2, 666-689.
- [10] E. LOOIJENGA, *Cohomology of $M3$ and $M1$ 3. Mapping class groups and moduli spaces of Riemann surfaces*, (Gottingen, 1991/Seattle, WA, 1991), *Contemp. Math.*, 150, Amer. Math. Soc., 1993, 205-228.
- [11] C. MACMEIKAN, *Modules of derivations for toral arrangements*, *Indag. Math. (N.S.)* 15 (2004), no. 2, 257-267.
- [12] L. MOCI, *Combinatorics and topology of toric arrangements defined by root systems*, *Rend. Lincei Mat. Appl.* 19 (2008), 293-308.
- [13] L. MOCI, *A Tutte polynomial for toric arrangements*, arXiv:0911.4823 [math.CO], to appear on *Trans. AMS*.
- [14] L. MOCI, *Zonotopes, toric arrangements, and generalized Tutte polynomials*, FPSAC 2010 Proceedings, DMTCS.
- [15] L. MOCI, *Wonderful models for toric arrangements*, arXiv:0912.5461 [math.AG], to appear on *Int. Math. Res. Not.*
- [16] P. ORLIK, M. TERAQ, *Arrangements of hyperplanes*, Springer-Verlag 300 (1992)
- [17] M. SALVETTI, *Topology of the complement of real hyperplanes in \mathbb{C}^n* , *Inv. Math.*, 88 (1987), no.3, 603-618.
- [18] M. SALVETTI, *The homotopy type of Artin groups*, *Math. Res. Lett.* 1 (1994), no. 5, 565-577.
- [19] S. SETTEPANELLA, *The integer cohomology of toric Weyl arrangements*, arXiv:1008.0631 [math.GT]
- [20] E. H. SPANIER, *Algebraic topology*, McGraw-Hill (1966)
- [21] A. RAM, *Alcove walks, Hecke algebras, spherical functions, crystals and column strict tableaux.*, *Pure Appl. Math. Quart.* 2 (2006), 963-1013.