



TÜBİTAK

Research Note

Turk J Bot
33 (2009) 191-226
© TÜBİTAK
doi:10.3906/bot-0805-12

Check-list of Additional Taxa to the Supplement Flora of Turkey IV

Neriman ÖZHATAY^{1,*}, Şükran KÜLTÜR¹, Serdar ASLAN²

¹Istanbul University, Faculty of Pharmacy, Department of Pharmaceutical Botany, 34116, İstanbul - TURKEY

²Gazi University, Faculty of Sciences and Arts, Department of Botany, 06500, Ankara - TURKEY

Received: 15.05.2008

Accepted: 29.04.2009

Abstract: The fourth check-list of the series entitled "Additional taxa to the Flora of Turkey" comprises a total of 175 taxa based on data given in 162 papers published up until the end of 2007. It provides details of taxa not recorded in the 11 volumes of the Flora of Turkey and the East Aegean Islands nor in the there previously published supplementary check-lists.

Key Words: Turkish Flora, new species, additional taxa

Türkiye Florası Ek Ciltlerine İlave Edilen Taksonların Listesi IV

Özet: "Türkiye Florasına ilave taksonlar" adlı seri halinde yayınlanmakta olan çalışmanın dördüncüsüdür. Bu çalışma 2005-2007 yılları arasında yayınlanan 162 çalışmada verilen takson ile Türkiye Florası ve eklerinde yer almayan taksonlar ile bu seri çalışmanın ilk üçünde bulunmayan toplam 175 taksonu içermektedir.

Anahtar Sözcükler: Türkiye Florası, yeni türler, ilave taksonlar

Introduction

The Second Supplement of *Flora of Turkey and the East Aegean Islands* was published in 2000. Subsequent to its publication (Güler et al., 2000) and up until the end of 2007, 470 additional taxa have been added to the flora, published in over 358 papers in various journals. In seven years, 470 taxa have been added to the flora of Turkey in check-lists III and IV; many are new to science.

Since the publication of the *Flora of Turkey* (Davis, 1965-1985; Davis et al., 1988) the Turkish flora has received considerable attention from botanists from both Turkey and abroad. Ongoing library work revealed that we inevitably had overlooked a certain number of taxa. We were fully aware that subsequent to the publication of the first paper of the series, further additions would be

discovered to Turkey's exceptionally rich flora. For these reasons, the first author realised that a series of papers would be necessary every few years. Nearly 500 papers on the flora of Turkey published in a wide range of journals and other publications since 1988 have been reviewed by the authors. They are now stored as a collection in the library of the Herbarium of Faculty of Pharmacy, İstanbul University (ISTE). The first check-list of the taxa additional to the *Flora of Turkey* was published in 1994 (Özhatay et al., 1994) and the second in 1999 (Özhatay et al., 1999), with a third one following in 2006 (Özhatay & Kültür, 2006). The present fourth part aims to compile all the published taxa added to the *Flora of Turkey* from 2005 to 2007. After the publication of check-list III, since 2005, 162 further papers dealing

* E-mail: s_kultur@istanbul.edu.tr

with the flora of Turkey have been published and many additional taxa have been recorded.

The present list contains a total of 175 taxa, of which 121 taxa were added as taxa new to science (comprising 85 species, 23 subspecies, 9 varieties, and 3 hybrids), whilst a further 54 known taxa have previously been unrecorded from Turkey (40 species, 13 subspecies, and 1 variety) (Table 1). The latter include one genus new for science (*Dumaniana*) and seven genera previously unreported from Turkey (*Asperuginoides*, *Caucasalia*, *Leptaleum*, *Loncomelos*, *Scopolia*, *Stroganowia*, and *Tetradiclis*). The remaining 295 taxa represent new combinations, new status indications, or newly suggested synonyms for previously known taxa (Table 2).

The list of the additional taxa is arranged in alphabetical order within families, genera, and species under four main systematic groups: *Pteridophyta*, *Gymnospermae*, *Dicotyledonae*, and *Monocotyledonae*. The format is in accordance with that in Davis' *Flora of Turkey* and references to original papers appear in brackets at the end of the taxon name. It is hoped that this list will provide useful information for researchers currently working on the Turkish flora.

We have endeavoured to trace all additional taxa to the flora published until the end of 2007. Nevertheless, we are certain that further taxa remain unreported either in Davis' *Flora of Turkey* or in our supplementary check-lists: accordingly, we invite additions or comments for future issues in the series.

Table 1. The list of the additional taxa to the Flora of Turkey after publication of the Second Supplement (check-list III + check-list IV).

Taxa new to science			Taxa new to Turkey		
	Check-list III	Check-list IV		Check-list III	Check-list IV
genus	3	1	genus	6	7
species	154	85	species	75	40
subspecies	12	23	subspecies	16	13
variety	15	9	variety	13	1
hybrid	8	3	hybrid	2	--
form	--	1	form	--	--
Total	189	121	Total	106	54
Total	310	Total		160	

TOTAL After the second supplement was published $310 + 160 = 470$ taxa were added to the Turkish flora until the end of 2007.

Table 2. The list of taxa whose taxonomic status changed after publication of the Second Supplement (Check-list III and Check-list IV).

	Check-list III	Check-list IV
comb. nov.	11	87
comb. + stat. nov.		63
stat. nov.	13	5
syn. nov.		138
nomen nov.	4	2
Total	28	295

The list of additional taxa

e = endemic species; * = new record for Turkish flora;

** = new taxon for science

■ = new genus for Turkish flora, ■ ■ = new genus for science

□ = genus previously known by different name

PTERIDOPHYTA

ISOETACEAE (1: 36)

Isoetes L. (1: 37)

e **[*I. anatolica* Prada et Rolleri in Bot. J. Linn. Soc. 147: 215-222 (2005)].

(Prada & Rolleri, 2005)

Type: (Turkey) A3 Bolu, al S de Abant Gölü, 18.vi.2001, borde de charca, C. Aedo et al. 6185 (holo. MA)

Isotypes: Idem., -plants B, C, D, E-; Turkey: Bolu, al S de Abant Gölü, 18.vi.2001, borde de charca, C. Aedo et al. 6181 (MA 688427).

SPERMATOPHYTA

GYMNOSPERMAE

CUPRESSACEAE (1: 76)

Juniperus L. (1: 78)

* *J. deltoides* R.P.Adams in Phytologia 86: 49-51 (2004).

(Adams, 2004)

Type: (Greece) 14 km E. of Archova, 22.v.2001, R.P.Adams 9436 (holo. BAYLU)

(Turkey). **B3** Eskisehir: 30 km N Eskisehir, 1064 m, 20.v.2001, R.P.Adams 9430, 9431, 9432 (BAYLU). **C6** Hatay: 8 km from Belen towards Antakya, 600 m, 6.v.1965, M.J.E. Coode & B.M.G.Jones 524 (K). **A7** Trabzon: Macka, 300 m, 2.iv.1960, Stainton 8152 (K). **C2** Antalya: Elmalı, 1800 m, 19.iv.1936, T.A.Tengwall, 400 (K). **A8** Artvin: 700 m, 27.vi.1957, Davis & Hedge 30095 (K). **B4** Ankara: above Ankara, 400 m, 5.vii.1974, P.H.Davis 13062 (K).

J. oxycedrus L. Sp. Pl. 1038 (1753) subsp. *oxycedrus*

e ** var. *spilinanus* Yalt., Eliçin et Terzioğlu in Turk J. Bot. 31: 38 (2007).

(Yaltırık et al., 2007)

Type: (Turkey) **B1** Manisa: Spildağı National Park, in forest clearings of *Pinus brutia* and *P. nigra* subsp. *nigra* var. *caramanica* and on stony slopes, 800-1400 m, 28.v.1995, KATO 13371 (holo. KATO; iso. KTUB).

ANGIOSPERMAE

DICOTYLEDONES

BORAGINACEAE (6: 237)

Alkanna Tausch (6: 414)

* *A. stribrnyi* Velen. in Fl. Bulg. Suppl. 647 (1891).

(Greuter & Raus, 2001)

European Turkey. **A1** (E) Tekirdağ: Kumbaga (Rodosto, Tekirdağ), by "St. Anna" Chapel, 4.5.1913, Nikolov (SO).

Cynoglossum L. (6: 306)

**C. vanense* Sutorý in Edinb. J. Bot. 61: 119-124 (2005).

(Sutorý, 2005)

Type: (Turkey) **C10** Hakkari: [Şemdinli]; Cheljane-çaj, 11.vi.1916, A.B.Schelkovnikov & V.Schiptschinskij 100-16, Expeditio Urmiensis Musei Caucasici 1916 (holo. LE).

Onosma L. (6: 326)

e ***O. mersinana* Riedl, Binzet et Orcan in Edinb. J. Bot. 61: 127-129 (2005).

(Riedl et al., 2005)

Type: (Turkey) **C5** Mersin: Erdemli, Müglü deresi, 15.vi.2001 Binzet 119 (holo. GAZI; iso. E, W, Mersin University, Turkey).

e ***O. riediana* Binzet et Orcan in Novon 17: 8-9 (2007).

(Binzet & Orcan, 2007)

Type: (Turkey) **C5** Mersin: Gülnar, Gülnar-Ermenek 42 km, 1450 m, roadside, fallow fields, 24.v.2003. Binzet G7 (holo. GAZI; iso. GZU, M).

Paracaryum DC. Boiss. (6: .282)

e ***P. hedgei* Aytaç et R.R.Mill in Edinb. J. Bot. 61: 109-111 (2005).

(Aytaç & Mill, 2005)

Type: (Turkey) **C4** Konya: Ereğli, 3 km N of İvriz, around Ayrancı Dam. 122-1300 m, protected *Abies cilicica* and *Pinus nigra* forest, 22.vi.1997, Z.Aytaç 7595 (holo. GAZI; iso. ANK, E).

Rindera Pallas (6: 300)

e ***R. dumanii* Aytaç et R.R.Mill in Edinb. J. Bot 61: 113-115 (2005).

(Aytaç & Mill, 2005)

Type: (Turkey) **C3** Antalya: Akseki-Beyşehir: Çamlık kasabası, Kızıldağ doğu yamaçları, 1600-1700 m, *Pinus nigra* orman açıklığı, 30.vi.1998, Z.Aytaç 7868 & H.Duman (holo. GAZI; iso. ANK, E).

CAMPANULACEAE (6: 2)

Campanula L. (6: 2)

**C. cecilii* Chitt. in Gardeners' Chronicle Ser. 3, 89: 397 Fig. 210 (1931).

(Sáez & Aldasoro, 2003)

S.E. Anatolia. **C8/9** Mardin, Senar, 6.vi.1888, Sintenis 921 (W); **C8** Siirt: Sirnak to Gizre, 14 km from Sirnak, 8.v.1966, Davis 42678 (W); **C8** Mardin: NW of Mardin, Diyarbakir, 30.v.1983, Sorger 83162 (W).

e ***C. demirsoyi* Kandemir in Nord. J. Bot. 25: 53-55 (2007).

(Kandemir, 2007)

Type: (Turkey) **B7** Erzincan: Munzur mountains, along Mercan river, 1968 m, limestone crevices 04.vii.2005, A.Kandemir 6992 (holo. GAZI; iso. ANK).

**C. hierosolymitana* Boiss., Diagn. Pl. Orient. Ser. 1, 11: 62 (1849).

(Sáez & Aldasoro, 2003)

Lectotype: [Palestine], aux pieds des rochers à Jérusalem, Naplouse, iii-iv. 1846 Boissier s.n. (G).

Turkey: Syria, prope Alexandrette (İskenderun), 1834, Montbret 18 (W).

C. pamphylica (Contandr. et Pamukç.) E.Akçicek et Vural in Ann Univ Sci 46 (54) (1972).

e **subsp. *afyonica* E.Akçicek et Vural in Ann. Bot. Fennici 42: 408-409 (2005).

(Akçicek et al., 2005)

Type: (Turkey) **B3** Afyon: Şuhut, Akyatak mahallesi, 1300-1350 m, crevices of volcanic rocks, 16.vii.2002, E.Akçicek 3758 (holo. GAZI; iso. ANK. ISTF)

CARYOPHYLLACEAE (2: 15)

Arenaria L. (2: 17)

A. dianthoides Smith, Ic.Pi.Ined. 16, t. 16 (1789).

e **subsp. *tuncbasi* Eren et Parolly in Willdenowia 36: 834-835 (2006).

(Parolly & Eren, 2006)

Type: (Turkey) **C3** Antalya: Beydağları, Çalbalı Dağı, ascent from Feslikan Yaylası, 1900 m thorn-cushion community, 6.vi.2003, Eren 37/3 & Şirin (holo. GAZI; iso. AKDU, AYDN, B, herb. Parolly).

Bolanthus (Ser.) Reichb. (2: 171)

e ***B. huber-morathii* C.Simon in Bauhinia 6:83 (1977).

(Simon, 1977)

Type: (Turkey) **A2 (A)** Bursa: South side of the Ulu Dağ, at the road between Soğukpinar and Keles, 3 km after Soğukpinar, ca. 900 m, 3.viii.1976 (holo. Hb. C.Simon 76-178).

Cerastium L. (2: 73)

C. brachypetalum Pers., Synops. pl. 1: 520 (1805)

*subsp. *tauricum* (Sperengel) in Murb. Lunds. Univ. Arsskr. 27 (5): 159 (1892).

(Keskin & Çırıcı, 2006)

Turkey: **A2(A)** İstanbul Burgazada, V. 1910, S.Selinka, ISTF 16; Tuzla istasyonu kenarı, 29. iv. 1962, B.Tutel 62/12, ISTF 18374 **C5** İçel: Tarsus, Nomrun bölgesi, Karanlıkdere serisi, Kazağacı, Kurukatran mevkisi, güney orta yamaçlar, 1500 m, 20.v.1979, F.Bozkuş 25536, Mersin: Gülnar-Ermenek arası, Gülnar dan 18 km, 1100 m, 10.vi.1950, A.Atilla, ISTF 10403.

Gypsophila L. (2: 149)

e ***G. osmangaziensis* E. Ataşlar et A.Ocak in Ann. Bot. Fennici 42: 57-59 (2005).

(Ataşlar & Ocak, 2005)

Type: (Turkey) **B3** Eskişehir: Osmangazi University Meşlik Campus area, 810 m, steppe, open rocky places, 28.viii.2002, E. Ataşlar & A. Ocak (holo. OUFE).

Saponaria L. (2: 138)

**S. bargyliana* Gombault in Bulletin Societe Botanique de France 109: 265 (1962).

(Mutlu, 2006)

S. Anatolia. **C6** Hatay: Erzin, 30 km from Kuzuculu village towards Ufacık area, 1200-1350 m, 17.vi.2002, *F. orientalis*-*C. orientalis* mixed forest, B. Mutlu 8034.

e ***S. karapinarensis* Vural et N. Adığüzel in Belg. Jour. Bot. 139: 252-254 (2006).

(Vural et al., 2006)

Type: (Turkey) **C4** Konya: Karapınar, Meke Gölü, 1000 m, volcanic scree, 30.vii.1997, M. Vural 7969, N. Adığüzel & F. Ertuğ (holo. GAZI; iso. ANK, HUB, EGE, Hb. YILDIRIMLI).

Silene L. (2: 179)

S. acaulis (L.) Jacq., Enum. Stirp. Vindob. 78 et 242 (1762).

e **subsp. *vanensis* Ozgökçe et Kit Tan in Ann. Bot. Fennici 42: 144-147 (2005).

(Özgökçe et al., 2005)

Type: (Turkey) **B9** Van: Ahta Dag, south-east of Kalecik village near Ozalp, 2500-2700 m, 15.v.1998 F.Özgökçe 5911 (holo. VANF; iso. BEOU, hb. Kit).

e ***S. nerimaniae* G.Ecevit Genc, Kandemir et İ.Genc in Nord. J. Bot. 25: 58-59 (2007).

(Ecevit Genc et al., 2007)

Type: (Turkey) **B7** Erzincan: Kemaliye, 4 km from Erzincan by Salihli village, 1502 m, 12.v. 2005, ISTE 83459 (holo. ISTE).

e ***S. sumbuliana* İ.G.Deniz et O.D.Düşen in Ann. Bot. Fennici 41: 293-295 (2004).

(Deniz & Düşen, 2004)

Type: (Turkey) **C2/3** Antalya: Elmalı Çamkuyu, the top of Avlankuzu mountain, 1850-1880 m, clearing in *Cedrus libani* forest and on calcareous slopes, 21.v.2001, İ.G. Deniz 1573 (holo. AKDU; iso. ANK, HUB, GAZI).

COMPOSITAE (5: 1)

Bidens L. (5: 46)

**B. frondosa* L., Sp. Pl. 832 (1753).

(Coşkunçelebi et al., 2007)

N.E. Anatolia. **A7** Trabzon: Kanuni Kampüsü, ditches, 12.vii.2006, 50-100 m, Coşkunçelebi & Vladimirov (SOM), 28.viii.2006, 50-100 m, Coşkunçelebi 629 (KTUB). **A8** Rize: İyidere, İyidere merkez kavşak, roadsides, 19.vii.2006, 5-10 m, Coşkunçelebi 627 (KTUB); Artvin: on the roadsides to Artvin city centre, 24.viii.2006, 200-300 m, Coşkunçelebi 628 (KTUB).

■ *Caucasalia* B.Nord. in Pl. Syst. Evol. 206 (1-4): 22, (1997).

e ***C. pontica* (K.Koch) Greuter in Willdenowia 35: 227 (2005).

(Greuter & Raab-Straube, 2005)

Ref. note: Koch did not mention his collecting locality, but stated the general area, the mountains of Pontus ("im pontischen Gebirge"). According to Lack (in Notes Roy. Bot. Gard. Edinburgh 37: 89. 1978) this area is entirely situated in Turkey, A7 Trabzon to A8 Çoruh, where the species has not been collected since, but ought to be looked for.

Centaurea L. (5: 465)

e ***C. elazigensis* Kaya et Vural in Novon 17: 198-201 (2007).

(Kaya & Vural, 2007)

Type: (Turkey) **B7** Elazığ: Elazığ-Palu, Baltaşı village, dry and rocky slopes, 1400-1500 m, 20.vii. 2005, Z.Kaya 17 (holo. GAZI).

e ***C. glabro-auriculata* Uysal et Demirelma in Ann. Bot. Fennici 44: 219-221 (2007).

(Uysal et al., 2007)

Type: (Turkey) **C3** Konya: Derebucak, Tekneliyatak Tepesi, 25.vii.2003, Demirelma 2770 (holo. KNYA).

e ***C. goeksunense* Aytaç et H.Duman in Pak. J. Bot. 37: 563-565 (2005).

(Aytaç & Duman, 2005)

Type: (Turkey) **C6** Kahramanmaraş: Göksun, Binboğa dağı, 1600-1650 m, between Yeşilkent and Göksun, near fields, 25.vi.2001, Duman 8606. (holo. GAZI, iso. ANK, HUB, E, hb YILDIRIMLI).

e ***C. kizildaghensis* E.Uzunhisarcıklı, E.Doğan et H.Duman in Bot. J. Linn. Soc. 153: 61-64 (2007).

(Uzunhisarcıklı et al., 2007)

Type: (Turkey) **C3** Konya: Derebucak, Çamlık, Kızıl Dağ, 1600-1700 m, serpentine rocky slopes, open area of a *Pinus nigra* forest, 12. viii. 2003, H.Duman 9129 (holo. GAZI; iso. ANK, HUB).

e ***C. marashica* E.Uzunhisarcıklı, M.Teksen et E.Doğan in Ann. Bot. Fennici 42: 309-311 (2005).

(Uzunhisarcıklı et al., 2005)

Type: (Turkey) **B6** Kahramanmaraş: Göksun, Armut Mountain, Kartallık Kayası, 1700 m, rocky slopes, 23.vii.2003, M.E.Uzunhisarcıklı 1901 & M.Tekşen, E.Doğan (holo. GAZI; iso. ANK, HUB).

e ***C. tuzgoluensis* Aytaç et H.Duman in Belg. Jour. Bot. 139: 256 (2006).

(Vural et al., 2006)

Type: (Turkey) **C4** Aksaray: Eskil, Karagöz district, 1000 m, 29.vii. 1998, halophytic steppe, Duman 6826 & Aytaç (holo. GAZI, iso. ANK, HUB, Hb, YILDIRIMLI).

e ***C. ulrichiorum* Wagenitz, Hellwig et Parolly in Willdenowia 36: 424-427 (2006).

(Wagenitz et al., 2006)

Type: (Turkey) C4 Antalya: NE Demirtaş (Richtung pass), 950 m, steiler Erdhang, Exp. W, 7.9.2001, R.Ulrich 1/25 (holo. GOET).

e ***C. wernerii* Wagenitz, Hellwig et Parolly in Willdenowia 36: 429-431 (2006).

(Wagenitz et al., 2006)

Type: (Turkey) C3 Antalya: Finike-Korkuteli, N Finike, 440 m, senkrechte Kalkfelsen, SE-Exposition, 29.4.2004, R.Ulrich 4/1 (holo. GOET; iso. B).

Cirsium Miller (5: 370)

**C. eriophorum* (L.) Scop., Fl. Carn. Ed.2: 130 (1772).

(Daşkin et al., 2006)

W. Anatolia. A2(A) Bursa: Uludağ: İnegöl, alongside Kiran plateau, 1700 m, 20.vii.2004, G.Kaynak, R.Daşkin 20390 & Ö.Yılmaz (BULU); Bursa: Uludağ Natural Park, between Softaböğan stream and Yörükmezarlığı, 1750 m, 05.viii.2004, G.Kaynak 23195 (BULU).

Jurinea Cass. (5: 439)

**J. cypria* Boiss., Fl. Orient., Suppl., 311 (1888).

(Greuter & Raus, 2002)

S. Anatolia. C4 Mersin: Mut, high plateau of Kozlar, valley of Ayt river, in limestone cliff, alt. 1380 m, 20.vii.1999, Everest 882 (hb. Univ. Mersin).

Onopordum L. (5: 356)

**O. canum* Eig in Palestine J. Bot., Jerusalem Ser. 2: 193 (1942).

(Greuter & Raab-Straube, 2005)

S.E. Anatolia. C6 Urfa: c. 10 km N of Birecik, Wadi, Strassenrand, 380 m, 25.v.1983, Sorger 83-6-2 (W).

□ *Psephellus* Cass. (previously *Centaurea*)

e ***P. turcicus* A.Duran et Hamzaoğlu in Bot. J. Linn. Soc. 148: 496-499 (2005).

(Duran & Hamzaoğlu, 2005)

Type: (Turkey) A5 Yozgat: Aydıncık, Kazankaya town, Kazankaya Canyon passes, 750-950 m, crevices of limestone rocks, 28.vi.2000, A.Duran 5461 & Hamzaoğlu (holo. KNYA; iso. GAZI, ANK, HUB, GOET).

Ptilostemon Cass. (5: 415)

P. *gnaphaloides* (Cyr.) Sojak in Novit. Bot. Del. Sem. Horti Bot. Univ. Carol. Prag 1962: 46 (1962).

*subsp. *pseudofruticosus* (Pomp.) Greuter in Candollea 24: 48 (1969).

(Tuzlaci, 2005)

S.W. Anatolia. C1 Muğla: Bodrum; Türkbüyü köyünün çevresi, Türkbüyü koyunun kuzeybatısı, deniz kıyısındaki yamaçlar, 20 m, 24.iii.2000, E.Tuzlaci, MARE 6185.

□ *Rhaponticoides* Vaill. (previously *Centaurea*)

e ***R. hierroi* Ö.Eren in Plant Systematics and Evolution 267: 15-17 (2007).

(Eren, 2007)

Type: (Turkey) C3 Antalya: between Antalya and Saklıkent road, below Bakacak pass. Boyunduruk 1100 m, rocky slope, open *Pinus brutia* Ten. forest, limestone, 17.vii.2004, Eren 252/04 & Şirin (holo. AYDN; iso. B, E, GAZI, HUB).

Senecio L. (5: 145)

e ***S. salsuginea* H.Duman et Vural in Belgian Jour. Bot. 139: 254 (2006).

(Vural et al., 2006)

Type: (Turkey) B4 Konya: Gölyazı, around Tersakan Gölü, 950 m, 25.x.1999, salt steppe Duman 7084 & F.A.Karavelioğulları (holo. GAZI; iso. ANK, HUB, Hb. YILDIRIMLI).

**S. sylvaticus* L. in Sp. Pl. 2: 868. (1753).

(Greuter & Raus, 1995)

European Turkey A2 (E): İstanbul, km 9 from Binkılıç, in a forest clearing, 350 m, 04.vii.1974, A.Baytop (ISTE 30188).

□ *Tephroseris* (Rchb.) Rchb. (previously *Senecio*)

**T. cladobotrys* (Ledeb.) Griseb. et Schenk, Wiegmann Archiv. 18 (1): 343 (1852).

(Budak et al., 2007)

N.E Anatolia. A8 Artvin: Borçka, Balçı, Kayadibi, 1710 m, 17.vii.1977, Düzenli 964 (HUB); A9 Ardahan: Yalnızçam Plateau, Katmerlikaya, 2300 m, 19.viii.1984, Demirkuş 2756 (HUB); A9 Ardahan: from Şavşat to Ardahan, Yalnızçam Mountains, Çam Pass, below Kocabey Plateau, Budak 1951, Hamzaoğlu & Aksoy (herb. BOZOK).

CRASSULACEAE

Sempervivum L. (4:244)

e ***Sempervivum × feigeanum* Neeff in Kakteen Sukk. 56: 72 (2005).

(Neeff, 2005)

(*Sempervivum davisii* Muirhead × *Sempervivum staintonii* Muirhead)

Type: (Turkey) A8 Rize, Kaçkar Dağları, Yusufeli to Sarıgöl, 927 m, 24.vii.2003, Neeff H073. (holo. E).

e ***S. herfriedianum* Neeff in Kakteen Sukk. 56: 71 (2005).

(Neeff, 2005)

Type: (Turkey) A8 Rize: Kaçkar Dağları, Yusufeli to Sarıgöl, 2086 m, 24.vii.2003, Neeff H075. (holo. E; iso. K, HUB).

CRUCIFERAE (1: 248)

Alyssum L. (1: 362)

**A. idaeum* Boiss. et Heldr. Boiss. Diagn. Pl. Or. Nov. 2 (8): 35 (1849)

(Orcan & Binzet, 2006)

S. Anatolia. C5 Mersin: Yeniköy entrance, 28.iv.2003, R. Binzet AK-781; 11.v.2003, R. Binzet AK-782.

e ***A. mughlai* Orcan in Nordic J. Bot. 23: 703-705 (2006).

(Orcan, 2006)

Type: (Turkey) C2 Muğla: Marmaris-Muğla, 10 km to Marmaris, 52 km to Muğla, roadside, serpentine, 100 m, 06.v.1993, N.Orcan 750 (holo. Hb of Mersin University).

■ *Asperuginoides* Rauschert

**A. axillaris* (Boiss. et Hohen.) Rauschert in Taxon 31 (3): 558 (1982).

(Bani & Adıgüzel, 2006)

E. Anatolia. C9 Van: Çatak, Dalbastı village, around Şeytan deresi, 1320 m, 16.v.2004, B. Bani 2644.

Barbarea R. Br. (1: 433)

B. brachycarpa Boiss. Diagn. ser. 1 (5): 80 (1844). subsp. *minor* (K.Koch) Parolly & Eren

e **var. *pilicarpa* Parolly et Eren in Willdenowia 36: 832 (2006).

(Parolly & Eren, 2006)

Type: (Turkey) C3 Antalya: Beydağıları, Feslikan Yaylası above Geyikbayırı, 1800-1900 m, doline with Trifolio-Polygonetalia community, 10.v.2006, Eren 54/6 (holo. AYDN; iso. B, E, EGE, GAZI, HUB, hb. Parolly)

B. brachycarpa Boiss. Diagn. ser. 1 (5): 80 (1844)

e **var. *ilicifolia* Parolly, Nordt et Eren in Willdenowia 36: 828 (2006).

(Parolly & Eren, 2006)

Type: (Turkey) C3 Konya: kurz nach der Abzweigung von der Hauptstraße Manavgat-Seydişehir in Richtung Bozkır, 1460 m, offener Serpentinhang, Exp. NE, 30.vi.2003, Ulrich 3/10 (holo. B; iso. AYDN, GAZI, hb Parolly).

Hesperis L. (1: 452)

e ***H. kuerschneri* Parolly et Kit Tan in Willdenowia 36: 851-854 (2006).

(Parolly & Tan, 2006)

Type: (Turkey) C2 Denizli: Fethiye-Çameli, N Fethiye, unterhalb des (below) Tuzla Beli, 1300 m, Serpentinhang (serpentine slope), SE-Exp., 7.v.2002, Ulrich 2/1 (holo. B; iso. C, hb. Parolly).

e ***H. turkmendaghensis* A.Duran et A.Ocak in Bot. J. Linn. Soc. 147: 240-242 (2005).

(Duran & Ocak, 2005)

Type: (Turkey) B3 Eskişehir: Türkmendağlı, Efsunbaba tepesi, 1650 m, 01.vi.2003, A.Ocak 3482 (holo. KNYA; iso. ANK, E., GAZI, HUB, Osmangazi University Hb.).

Iberis L. (1: 309)

**I. saxatilis* L., Cent. Pl. 2: 23 (1756).

(Dirmenci et al., 2005)

B1 Balıkesir: İter trojanum, Mt. Ida in marmar, mont. Szus-Dagh, 17.vii.1883, P.Sintenis (photo). Kazdağ (Ida), Nanekırı, limestone, 1500 m, 19.v.2004, T.Dirmenci (2516 a) & F.Satılı (GAZI, HUB, Yıldırımlı) ibid 05.vi.2004, T.Dirmenci (2523) & F.Satılı (EGE), ibid Zığındere valley, limestone, 1600 m, 19.v.2004, T.Dirmenci 2516 b) & F.Satılı (ANK), Susuz dağ-Kuştepe, 1500 m, 02.vii.2004, T.Dirmenci (2872 b) & F.Satılı (Hb. T.Dirmenci).

Lepidium L. (1: 279)

**L. pinnatifidum* Ledeb., Fl. Ross. 1: 206 (1842).

(Ünal et al., 2007)

Type: Russia, Hab. In regione Astrachanensi, Blume, LE.

N.E. Anatolia. **B9** Muş: Bulanık, North-west of Şatırlar village, around Kırmızı Tuzla, saline places, 03.vi.2006, L.Behçet & F.Özgökçe and M.Ünal (LFM 742 in flower), 05.vi.2006 (LFM 792-in fruit).

■ **Leptaleum** DC.

**L. filifolium* (Willd.) DC., System. Veg. 2: 510 (1821).

(Ünal & Özgökçe, 2007)

Syn.: *Sisymbrium filifolium* Willd., Sp. Pl. Ed. 4, 3: 495 (1800).

Type: Habitat in Sibiria ad Kumam fluvium, B.

N.E. Anatolia. **A9** Kars, Halılışlak village, around River Aras, sandy banks, 06.v.2006, M.Ünal & F.Özgökçe 9295 (VANF).

Malcolmia R. Br. (1: 460)

**M. exacoides* (DC.) Spreng., Syst. Veg.Ed. 16: 2: 899 (1825).

(Özgökçe & Ünal, 2007)

Type: in Libano loco dicto Zaale, Labillardiere.

S.E. Anatolia. **C9** Siirt: Siirt to Şırnak, 30 km, around River Dicle, steppe and fields, 11.iii.2006, F.Özgökçe & M.Ünal 9138 (in flower); 14.iv.2006, F.Özgökçe & M.Ünal 9149 (in fruit).

Matthiola R. Br. (1: 447)

e ***M. trojana* T.Dirmenci, F.Satılı et G.Tümen in Bot. J. Linn. Soc. 151: 431-433. (2006).

(Dirmenci et al., 2006)

Type: (Turkey) **B1** Balıkesir: Edremit, Kazdağ (Ida), Naneköprü place, rocky slopes, limestone, 1550 m, 5.vi.2004, T.Dirmenci (2522) & F.Satılı (holo. ISTE; iso. ANK, EGE, GAZI).

□ **Noccaea** Moench (previously *Thlaspi*)

e ***N. camlikensis* Aytaç, Nordt et Parolly in Bot. J. Linn. Soc. 150: 410-412 (2006).

(Aytaç et al., 2006)

Type: (Turkey) **C3** Konya: Derebucak, Çamlık village, Kızıldağ, 1400-1600 m, clearings in *Pinus nigra* forest, 6.vi.2000, Duman 7306 (holo. GAZI; iso. ANK).

Physoptychis Boiss. (1: 356)

e ***P. purpurascens* Çelik et Akpulat in Bot. J. Linn. Soc. 154: 393-395 (2007).

(Çelik et al., 2007)

Type: (Turkey) **B6** Sivas: Deliilyas district, serpentine, chalky steppe and slopes, 23.v.2004, Çelik 6232; Akpulat & Dönmez. (holo. CUFH; iso. ISTE).

■ **Stroganowia** Kar. et Kir.

e ***S. leventii* V.I.Dorof. in Byull. Moskovsk. Obshch. Isp. Prir., Otd. Biol. 109: 72(2004).

(Dorofeyev et al., 2004)

Type: (Turkey) **B8** Erzurum ad meridiem 42 km ab Köprüköy, in parte superiore vallii Araks, 1800 m, 18.vi.2002, V.I. Dorofeev 999 (holo. LE).

DIPSACACEAE (4: 582)

Cephalaria Schröder ex Roemer et Schultes (4: 585)

e ***C. duzceensis* N.Aksoy et R.S.Göktürk in Nord. J. Bot. 25: 65-67 (2007).

(Aksoy et al., 2007)

Type: (Turkey) **A3** Düzce: Gölyaka, Elmacık mountain, Balıklı area, behind of Konaş, Melik deresi, open rock area of *Pinus sylvestris* forest, on disturbed ground, 30.vii.2004, N.Aksoy 5339 (holo. ISTO; iso. GAZI, AIBO, AKDU).

C. gaziapensis H.Sümbül in Edinb. J. Bot. 48: 33. f.3 (1990)

e **subsp. *pilifera* Parolly, Kürscher et Eren in Willdenowia 37: 247-249 (2007).

(Parolly & Eren, 2007)

Type: (Turkey) **C4** Antalya: Alanya, slopes of Akdağ (2461 m), between Şekerkahve and Gökbelen Yayları, 1100-1300 m, oak forest, limestone and dolomite, 12.vii.2000, Eren & Parolly 7947 (holo. B; iso. AYDN, E, Hb. Parolly).

e ***C. tutelianiana* S.Kuş et R.S.Göktürk in Nordic J. Bot. 23: 427-429 (2005).

(Kuş & Göktürk, 2005)

Type: (Turkey) **A2(E)** İstanbul: Kiraç, Bahçeşehir, north-west slopes, clearing macchie scrub, 10.ix.1997,

S.Kuş et al., ISTF 37158 (holo. ISTF; iso. AKDU, ISTE, HUB, GAZI).

Lomelosia Raf. (= *Scabiosa* sect. *Trochocephalus* Mert. & Koch).

e ***L. solymica* Parolly, Ö.Eren et Nordt in Willdenowia 35: 108-110 (2005).

(Parolly et al., 2005)

Type (Turkey) C3 Antalya: Tahtalı Dağı, SW Kemer, ascent from Beycik, vertical limestone cliffs, 6.viii.2004, Eren 387/04 & Taylan (holo. hb. Adnan Menderes University; iso. B, E, GAZI, HUB, hb. Parolly).

FAGACEAE (7: 657)

Quercus L. (7: 659)

Q. trojana P.B.Webb in London, Gard. Mag. 15: 590 (1839)

e **subsp. *yaltirikii* Ziel., Petrova et D.Tomaszewski in Willdenowia 36: 846-848 (2006).

(Zielinski et al., 2006)

Type: (Turkey) C5 İçel: between Haci Ahmetli and Çukurbağ, just near Çukurbağ, 18-20 km NNE of Mut, 1200 m, stony-clayey slope, edge of *Pinus brutia* forest, 31.v.1991, Boratynski, Browicz & Zieliski 6863 (holo. KOR).

Paratypes: (S Turkey), C4/5 İçel: near Abanoz, by the road to Ermenek, small meadow among rocks, 29.v.1991, Boratynski, Browicz & Zielinski 6780 (KOR 31026); Boratynski, Browicz & Zieliski 6780 (KOR 25063); C2/3 Antalya, Geyik Dağları, 1500 m, *Abies cilicica-Pinus nigra* forest on calcareous rocks, frequently, 5.viii.2005, Boratynski & Boratynska TU-05/32 (B, KOR 44631).

GERANIACEAE (2: 450)

Geranium L. (2: 451)

e ***G. kalenderianum* A.Ilçim et L.Behçet in Ann. Bot. Fennici 43: 451-453 (2006).

(İlçim & Behçet, 2006)

Type: (Turkey) B6 Kahramanmaraş: Afşin, Tanır Kasabası, Güvek Yaylası (high-land), Binboğa Dağı (Binboğa Mt.) rocky places, between rock clefts, 1915-

1950 m, 20.vi.2001, A.Ilçim 1373 (holo. KSUH; iso. GAZI, VANF).

GUTTIFERAEE (2: 355)

Hypericum L. (2: 355)

H. saxifragum Robson et Hub.-Mor. in Notes R.B.G. Edinb. 27: 194 (1967).

e **subsp. *eglandulosum* Parolly et Eren in Willdenowia 36: 836-837 (2006).

(Parolly & Eren, 2006)

Type: (Turkey) C3 Antalya: W Antalya, Antalya-Saklıkent road, 300 m, vertical limestone cliffs, E-exp., 15.ix.2004, Ulrich 4/28 (holo. hb. Parolly).

LABIATAE (7: 36)

Calamintha Miller (7: 323)

C. pamphylica Boiss. et Heldr. in Boiss., Diagn. Ser. 1 (12) 52 (1853)

e **subsp. *alanyense* S.Alan et Ocak in Ann. Bot. Fennici 44: 309-313 (2007).

(Alan et al., 2007)

Type: (Turkey) C4 Antalya: Alanya, Kargı brook, rock crevices, 2.vii.2002, S.Alan 1005 (holo. ESSE; iso. GAZI, HUB, hb. Osmangazi Univ.).

Micromeria Bentham (7: 335)

e ***M. maritima* Yıldırımlı, Sadıkoglu et Keskin in Ot Sistematisk Botanik Dergisi, 13: 29-30 (2006).

(Yıldırımlı et al., 2006)

Type: (Turkey) C2 Muğla: Marmaris vegetation, near coast, c. 5 m, 23.vii.1983, Ş.Yıldırımlı 5647 (holo. hb. Yıldırımlı; iso. GAZI).

**M. persica* Boiss., Diagn. Pl. Or. Nov. Ser. 1, 7: 48 (1846)

(Keskin & Sadıkoglu, 2007)

S. Anatolia. C6 Hatay: Antakya, Habib-i Neccar Mountain, slopes, 28.vi.2004, N.Sadıkoglu, ISTE 81697 ibid., 31.vii.2005, N.Sadıkoglu & M.Keskin 4470, ISTE 83175., St. Pierre s church, 31.vii.2005, N.Sadıkoglu & M.Keskin 4455, ISTE 83192.

Nepeta L. (7: 264)

N. sibthorpii Benth. Lab. Gen. Sp. 474 (1834)

e **subsp. *tumeniana* T.Dirmenci in Bot. J. Linn. Soc. 147: 229-232 (2005).

(Dirmenci, 2005)

Type: (Turkey) **B1** Balıkesir: Edremit, Kazdağı (Ida), Dökük, rocky slopes, 1300 m, 10.vii.1999, T.Dirmenci 1024 (holo. GAZI; iso. ANK, EGE, HUB, ISTE).

Phlomis L. (7:102).

e ***P. dinci* Yıld. in Ot Sist. Bot. Dergisi 13: 3 (2006) (Yıldırımlı, 2006)

Type: (Turkey) **B6** Malatya: Doğanşehir, Eskiköy, between Çoban deresi-ihtiyar mağarası, *Quercus-Juniperus excelsa* forest, 1600-1800 m, 25.vii.1971, H.Peşmen 2683 (holo. HUB).

e ***P. isiliae* Yıld. in Ot Sist. Bot. Dergisi 13:3(2006) (Yıldırımlı, 2006)

Type: (Turkey) **B7** Tunceli: Ovacık, above Tornova, Munzur dağları, 1100-1450 m, 26.vii.1979, Ş.Yıldırımlı 2127 (holo. HUB; iso. GAZI).

Salvia L. (7: 400)

e ***S. anatolica* Hamzaoğlu et A.Duran in Ann. Bot. Fennici 42: 216-218 (2005)

(Hamzaoğlu et al., 2005)

Type: (Turkey) **B7** Sivas: 22 km from Divriği to Kemaliye, 1580 m, 30.v.2003 A.Duran 6159, Hamzaoğlu & Sağıroğlu (holo. KNYA; iso. GAZI, ANK, HUB, hb. YILDIRIMLI).

Stachys L. (7: 199)

e ***S. yildirimlii* M.Dinç in Ann. Bot. Fennici 43: 143-145 (2006)

(Dinç & Doğan, 2006)

Type: (Turkey) **C5** Mersin: Arslanköy, Cocakdere, *Cedrus libani-Pinus nigra* forest, clearings, rocky slopes, crevices of limestone rocks, 1950 m, 15.vi.2004 M.Dinç 2150 & H.H.Doğan (holo. KNYA; iso. GAZI).

Teucrium L. (7: 53)

**T. chasmophyticum* Rech.f., Pl. Syst. Evol. 134: 287 (1980)

(Dönmez, 2006)

E. Anatolia. **C9** Siirt: Eruh, above Gölgelikonak village, *Quercus brantii* opening, limestone crevices, 19.vi.2005, AAd 121-19-1. Al-Shehbaz & M.Menke.

e ** *T. microphyllum* Desf. in Ann. Mus. Par. x. 300 (1807).

(Greuter & Raus, 2006)

SW Turkey: Datça Peninsula

e ***T. pseudaroanum* Parolly, Erdağ et Nordt, in Willdenowia 37: 252-257 (2007).

(Parolly & Eren, 2007)

Type: (Turkey) **C2** Antalya: Kasaba, SE Dirgenler, Schlucht des (gorge of) Demre Çayı, 140 m, Kalkfelsen (Marmor) (limestone rock marble), NW-exp., 24.iv.2006, Ulrich 6/5 (holo. B; iso. AYDN, hb. Parolly).

LEGUMINOSAE (3: 1)

Astragalus L. (3: 49)

e ***A. babacianum* A.S.Ertekin in Ann. Bot. Fennici 43: 291-293 (2006)

(Ertekin, 2006)

Type: (Turkey) **C8** Mardin: Mazıdağı, Mazidagi to Derik 6 km, 970 m, fields, 15.iv.2004, A.S.Ertekin 2004-126 (holo. DUF).

e ***A. friedericianus* Kit Tan & Zeitl. in Wulfenia 14: 133 (2007).

(Tan & Zeitlinger, 2007)

Type: (Turkey) **C4** Konya: dry south-facing slopes of Sertoval-Gedici, roadside steppe south of Karaman, c. 1080 m, 27.v.1988, J.Zeitlinger 4126 (holo. KL; iso. Hb. Kit Tan).

e ***A. nezaketae* A.Duran et Aytaç in Ann. Bot. Fennici 42: 381-385 (2005)

(Duran & Aytaç, 2005)

Type: (Turkey) **A8** Erzurum: Erzurum-İspir yolu 50. km, Eğerti köyü, Kelkaya mevkii, step, 2000-2200 m, 19.vii.1990, H.Duman & Z.Aytaç 3195 (holo. GAZI; iso. KNYA).

A. pinetorum Boiss. Diagrb zedb 1 (2): 77 (1843).

e **subsp. *multifoliatus* Akpulat et Çelik in Bot. J. Linn. Soc. 155: 227-229 (2007)

(Akpulat & Çelik, 2007)

Type: (Turkey) **B6** Sivas: Topçuyeniköy district, steppe, 1450 m, 27.vi.2004, Çelik 7025 & Akpulat (holo. CUFH; iso. ISTE).

**A. wagneri* Bunge, in Mem. Acad. Imp. Sc. Petersb. 15 (1): 125 (1868)

(Ekici et al., 2005)

E. Anatolia. B10 Ağrı: Doğubeyazıt, around Tavla village, steppe, 2400 m, 15.vii.2002, M.Ekici 3179, Z.Aytaç & H. Akan (GAZI).

e ***A. turkmenensis* Dural, O.Tugay et Ertugrul in Ann. Bot. Fennici 44: 399-401 (2007)

(Dural et al., 2007)

Type: (Turkey) C4 Konya: Karapınar-Hotamış arası, Kamişağıl köyü çevresi, step, 1050 m, 18.v.2003 Ertuğrul 2764, O.Tugay, Dural, Uysal & Demirelma (holo. KNYA; iso. GAZI, K.).

Chamaecytisus Link (3: 16)

**C. jankae* (Velen.) Rothm.

(Vladimirov et al., 2006)

European Turkey. A2(E) İstanbul: grassy places around Constantinopol (İstanbul), Bijuk-Han station, 16.v.1915, coll. B.Davidov (SOM 43786).

Genista L. (3: 24)

G. januensis Viv. Elench. Pl. Horti Bot. 19 (1802).

*subsp. *januensis* in Ann. Bot. Fennici 41: 455-456 (2004).

(Zielinski et al., 2004)

G. spatulata Spach, Ann. Sci. Nat. Bot. Ser. 3: 128. 1845.

Syntypes: [NW Turkey] "in Olympo Bithynico (Uludağ) et prope Byzantium", Aucher-Eloy (P). NW and W Anatolia

Lupinus L. (3: 38)

L. angustifolius L. Sp. Pl. 722 (1753).

*subsp. *reticulatus* (Desv.) Coutinho in Fl. Port. 315 (1913).

(Özen & Aka, 2007)

W. Anatolia. B1 Balıkesir; Ayvalık, Sarımsaklı (Küçükköy), Badavut, military coastal zone, 0 m, 19.05.2003, F.Özen 1736.

Onobrychis Adanson (3: 560)

O. *alba* (Waldst. & Kit.) Desv., Jour. Bot. Appl. 3: 83 (1814).

*subsp. *calcarea* (Vandas) P.W. Ball, Feddes Repert. 79, 41 (1968).

(Vladimirov et al., 2007)

European Turkey A1(E) Edirne: Sinanköy village, in the meadows, 14.vi.2007, M.Aybeke, C.Kurt & A.Semerçi (EDTU 9506), Centre: Budakdoğanca village, meadows, 14.vi.2007, M.Aybeke, C.Kurt & A.Semerçi (EDTU 9507).

**O. meschetica* Groshh. in Flora U.R.S.S., vol. XIII, p.554 (1948).

(Aktoklu, 2001)

Type: (S. Transcaucasus) prope Borzhomi, 2.vi.1865, Radde 35 (LE).

N.E. Anatolia. A9 Kars: Posof, Binbaşı Eminbey-Yurtbeyler köyü arası, yolboyu korunmuş alanlar, 1500-1900 m, 5.viii.1982, N.Demirkuş-1805 (HUB, Hb. Mustafa Kemal University).

Trifolium L. (3: 384)

T. tumens Stev. et Bieb. Fl. Taur.-Cauc. 2: 217 (1808).

*var. *talychense* (Chalilov) Zoh. in Israel J. Bot. 19: 322 (1970).

(Keskin, 2005)

S. Anatolia. C6 Osmaniye: Zorkun yaylası, *Pinus nigra* orman altı, Amanos dağı, 14.vi.1968, c. 1400 m, Y.Akman ANK 7846.

**T. vavilovii* Eig. in Bull. Appl. Pl. Breed. (Leningrad) 7, 1: 108 (1934).

(Keskin, 2005)

E. Anatolia. B7 Malatya: Kubbe dağı eteği, Köpekli köyü, Pötürge yolu üzeri, çayır, 1300 m, 22.vi.1982, Y.Altan, ANK 2650.

Trigonella L. (3:452)

e** *T. velutinoides* Hub.-Mor. in Bauhinia 4: 197 (1971).

(Huber-Morath, 1971)

Type: (Turkey) C4 Konya: Karaman, Karaman-Ermenek, Quercetum 34 km nach Karaman, 1300 m, 8.vi.1948, Huber-Morath 8197; do., Steppe 15 km nördlich von Karaman, an der Strasse nach Konya, 1030 m, 7.vi.1969, Ch.Simon 69-983 (holo. Hb. Simon; iso. Hb. Hub.-Mor.).

MALVACEAE (2:401)

Alcea L. (2: 411)

e ***A. karsiana* (Bordzil.) Grossg. in Zhurn. Russk. Obshch. VII (1922) 1924.

(Greuter & Raus, 1989).

Type: (Turkey) A9 Kars: Ardahan, Karaurgan and Karakurt. (holo.LE).

* *A. sophiae* Iljin in Zametki Sist. Geogr. Rast. 15: 42, (1949).

(Greuter & Raus, 1989)

E. Anatolia. A9 Kars: Sürmeli and Karakale. (holo. Leningrad Herbarium (LE)).

PLANTAGINACEAE (7:504)

Plantago L. (7:505)

P. uliginosa F.W.Schmidt

*subsp. *mandakii* Chrtek in as. Nár. Muz. Praze, Rada P ír. 173: 89 (2004)

(Chrtek, 2004)

C. Anatolia B5 Kayseri: Secundum viam sub cacumine montis Erciyes Dagi, prope campum, prope oppidum Kaiser, c. 3000 m, 20.ix.2000, B.Mandák (holo. PR 11860).

PLUMBAGINACEAE (7: 463)

Acantholimon Boiss. (7: 478)

A. *acerosum* (Willd.) Boiss., Diagn. Ser. 1 (7): 80 (1846).

e **subsp. *longibracteolorum* Doğan et Akaydın in Bot. J. Linn. Soc. 154: 410-411 (2007)

(Doğan & Akaydın, 2007)

Type: (Turkey) B3 Eskişehir: Sündiken Dağı, 400-500 m, 22.vi.1974, Ekim 923 (holo. ANK).

A. *calvertii* Boiss., Diagn. Ser.2 (4): 65 (1859).

e **var. *glabrum* Akaydın et Dogan in Bot. J. Linn. Soc. 154: 403-404 (2007).

(Doğan & Akaydın, 2007)

Type: (Turkey) A8 Erzurum: between Pasinler and Narman, Çimenli Yayla, 2160 m, 17.vii.2001, Doğan & Akaydın 6930 (holo. ANK).

A. *capitatum* Sosn., Monit. Jard. Bot. Tiflis 11: 11 (1915).

e **subsp. *sivasicum* Dogan et Duman in Bot. J. Linn. Soc. 154: 401 (2007).

(Doğan & Akaydın, 2007)

Type: (Turkey) B6 Sivas: Ulaş, between Kutlukaya and Boğazdere village, 1400-1500 m, serpentine, steppe, 7.vii.2003, H.Duman 8983 (holo. GAZI).

e ***A. ekimii* Dogan et Akaydın in Bot. J. Linn. Soc. 154: 404 (2007).

(Doğan & Akaydın, 2007)

Type: (Turkey) B9 Ağrı: between Akçay and Cumaçay, high mountain steppe, 2300 m, 18.vii.2001, Doğan & Akaydın 6962 (holo. ANK).

e ***A. evrenii* Doğan et Akaydın in Bot. J. Linn. Soc. 149: 352-353 (2005).

(Doğan & Akaydın, 2005)

Type: (Turkey) B7 Elazığ: Palu, Yayılm Mountain, near Baltaşı, steppe, 1500-1600 m, 29.vi.1983, H.Evren 1585 (holo. FUH).

A. *halophilum* Bokhari, Notes R.B.G. Edinb. 30: 297, fig.2 (B) (1970).

e **var. *coloratum* Dogan et Akaydın in Bot. J. Linn. Soc. 154: 412 (2007).

(Doğan & Akaydın, 2007)

Type: (Turkey) B4 Konya: between Cihanbeyli and Tuzla, near Tuz Gölü, 950 m, 21.vi.2002, Doğan & Akaydın 7398 (holo. ANK).

A. *huetii* Boiss., Fl. Or. 4: 835 (1879).

e **var. *breviscapum* Akaydın et Dogan in Bot. J. Linn. Soc. 154: 404 (2007).

(Doğan & Akaydın, 2007)

Type: (Turkey) B6 Kahramanmaraş: Göksun, above Keklikoluk, 2100 m, 9.vii.2001, Doğan & Akaydın 6808 (holo. ANK).

A. *lycaonicum* Boiss.et Heldr. in DC., Prodr. 12: 627 (1848).

e **subsp. *cappadocicum* Dogan et Akaydın in Bot. J. Linn. Soc. 154: 412-413 (2007).

(Doğan & Akaydın, 2007)

Type: (Turkey) A4 Çankırı: Kalecik, Karabasan Da., 850 m, 21.viii.2000, Doğan and Akaydın 6474. (holo. ANK).

Limonium Miller (7:465)

e ***L. adilguneri* Yıld. & Doğru-Koca in Ot Sist. Bot. Dergisi 13: 13 (2006).

(Yıldırımlı & Doğru-Koca, 2006)

Type: (Turkey) B4 Aksaray: Eskil, 15 km from Eskil to Tuzgölü, c. 900 m, 24.ix.2006, .Yıldırımlı & A.Doğru-Koca 32268 (holo. Hb. YILDIRIMLI; iso. K, HUB, GAZI).

RANUNCULACEAE (1: 95)

Delphinium L. (1: 108)

**D. balcanicum* Pawl. in Fragm. Fl. Geobot. 9: 439 (1963).

(Dane & Ekici, 2006)

European Turkey. A1(E): Tekirdağ, Edirne, Çanakkale Eceabat, Seddulbahir, 07.vii.1987, F.Dane (EDTU 1733); Edirne: Budak Doganca. 08.vii.1988, F.Dane (EDTU 2398); Edirne: Süleoğlu, road side, 14.vii.1988, F.Dane (EDTU 2469); Tekirdağ, Şarköy, road side, 15.vii.1988, F.Dane (EDTU 2573).

Ranunculus L. (1:146)

e ***R. anatolicus* Ü.Akkemik, E.Akalin & N. Özhata in Nordic J. Bot. 25:311-313 (2007).

(Akkemik et al., 2007)

Type: (Turkey) A9 Erzurum: Şenkaya-Karinca Düzü-Eşek Meydani Plateau, 2350 m, 26.vi.2002, Ü.Akkemik (holo. ISTO 29271; iso. ISTE 81440).

RESEDAEAE (1: 498)

Reseda L. (1: 499)

R. germanicopolitana Hub.-Mor. in Bauhinia 2: 300 (1965).

e **subsp. *glabra* Karavel. et Vural in Israel J. Plant Sci. 54: 109-110 (2006).

(Karavelioğulları et al., 2006)

Type: (Turkey) B5 Kırşehir: Çiçekdağı, sekili, Tuz tepesi, 900-950 m, 06.ix.1994, salt soil, H.Polat 1631 & F.A.Karavelioğulları (holo. GAZI; iso. ANK, HUB).

Paratypes: B5 Kırşehir: Çiçekdağı, Küçükteflek-Sekili, 900 m, 18.ix.1994, salt soil, H.Polat 2003 &

F.A.Karavelioğulları, Kırşehir: Çiçekdağı, Küçükteflek, Çorakdere, 900 m, 05.viii.1995, salt soil, F.A.Karavelioğulları 2353 & H.Polat.

ROSACEAE (4: 1)

Crataegus L. (4: 133)

e ***C. christensenii* Dönmez in Bot. J. Linn. Soc. 155: 234-236 (2007).

(Dönmez, 2007)

Type: (Turkey) C10 Hakkari: Şemdinli, Altınsu village, 4.4 km from Şemdinli to Yüksekova, among *Quercus* forest in the valley, 3.x.2001, A.A.Dönmez 10314 (fruiting material) (HUB).

e ***C. peshmenii* Dönmez in Bot. J. Linn. Soc. 155: 232-234 (2007).

(Dönmez, 2007)

Type: (Turkey) C10: Hakkari: 2.5 km from Şemdinli to Yüksekova, Altınsu village, among *Quercus* opening, 3.x.2001. A.A.Dönmez 10313 (holo. HUB; iso. EGE).

C. rhipidophylla Gand. Bull. Soc. Bot. France 18: 447 (1871).

e **var. *kutahyaensis* Dönmez in Bot. J. Linn. Soc. 155: 237-238 (2007).

(Dönmez, 2007)

Type: (Turkey) B6 Kütahya: Eskişehir road, 9.4 km, Ahmetoluk village, 23.viii.2001, A.A.Dönmez 9972 (holo. HUB).

e ***C. turcicus* Dönmez in Ann. Bot. Fennici 42: 62-64 (2005).

(Dönmez & Dönmez, 2005)

Type: (Turkey) A9 Artvin: Ardanuç pasture of Yolüstü village, opening of the deciduous forest, 41°06 N, 42°11 E, 1800 m, 30.ix.2002, A.A.Dönmez 11150 (holo. HUB; iso. L.).

Paratype: (Turkey) A9 Artvin, Ardanuç, pasture of Yolüstü village, opening of the deciduous forest, 21.vi.2003, A.Dönmez 11456 (HUB).

e ***C. yaltirkii* Dönmez in Bot. J. Linn. Soc. 148: 245-246 (2005).

(Dönmez, 2005)

Type: (Turkey) C9 Şırnak: Beytüşşebap, above Günyüzü village, opening of deciduous forest, 28.ix.2002; A.A.Dönmez (cited as AAD) 11143 &

B.Mutlu; 18.vi.2003, A.A.Dönmez 11261; 26.v.2004, A.A.Dönmez 11922. **C9** Şırnak, Beytüşşebap, from Dağdibi village to main road, alongside the road, 26.v.2004, A.A.Dönmez 11912 (holo. HUB; iso. E, ISTO).

Padus Miller (4: 6)

P. avium Mill. Gard. Dict. Ed. 8: no 1 (1768).

*subsp. *petraea* (Tausch) Pawłowski ex Holub. in Preslia (Praha) 36: 251-255 (1964).

(Terzioğlu & Coşkunçelebi, 2005)

N.E. Anatolia. **A8** Trabzon, Uzungöl, Demirkapı Köyü, 2175 m, Terzioğlu, 24.vii.2004, KATO 13368, Arpaözü Köyü, 2250 m, Terzioğlu, 30.viii.1995, KATO 12064 & KTUB 506.

SAXIFRAGACEAE (4: 249)

Saxifraga L. (4: 249)

S. adscendens L.

*subsp. *parnassica* (Boiss. et Heldr.) Hayek

(Tuzlaci & Bulut-Emre, 2006)

A2(E) Kırklareli: Kıyıköy, deniz kıyısındaki kayalıklar, **A2 (A)** İstanbul: Riva'nın 1 km doğusu, deniz kenarı (MARE 2661, MARE 4636).

SCROPHULARIACEAE (6: 458)

Verbascum L. (6: 461)

e ***V. cicekdagensis* Karavel. et Vural in Israel J. Plant Sci. 54: 106-109 (2006).

(Karavelioğulları et al., 2006)

Type: (Turkey) **B5** Kırşehir: Çiçekdağı, between Küçükteflek and Sekili, 900 m, 18.v.1995, roadside, F.A.Karavelioğulları 2365 and H.Polat (holo. GAZI; iso. ANK).

Paratypes: **B5** Çiçekdağı, between Küçükteflek and Sekili, 900 m, 18.v.1994, calcareous stony slopes, roadside, F.A.Karavelioğulları 2103 and H.Polat; Çiçekdağı, between Ömeruşağı and Kalevcı, 900-950 m, 19.v.1995, calcareous stony slopes, road-side, H.Polat 1994 and F.A.Karavelioğulları

e ***V. lindae* Parolly et Kit Tan in Willdenowia 37: 279-281 (2007).

(Parolly & Tan, 2007)

Type: (Turkey) **C3** Isparta: Sütçüler, Eğirdir-Sütçüler, 16 km N of Çandır, 1 km SE of Sığırlık Köyü, W-facing limestone cliff, 860 m, 21.v.2003, Ulrich 3/4 (holo. hb. Parolly; iso. B, herb. Kit Tan).

**V. samniticum* Ten. Fl. Nap. 3: 219 (1824-1829).

(Yılmaz & Dane, 2006)

European Turkey. **A1(E)** Kırklareli: Pınarhisar: Kaynarca village, roadside, 27.vi.2002. G.Yılmaz & F.Dane (EDTU 9498).

e ***V. yurtkuranianum* Kaynak, Daşkın et Yılmaz in Ann. Bot. Fennici 43: 457-459 (2006).

(Kaynak et al., 2006)

Type: (Turkey) **A2** Bursa: Gürsu, Gürsu-Ercek, entrance of Ercek village, 4.vi.2003 G.Kaynak, R.(Günay) Daşkın, Ö.Yılmaz, E.Erdoğan 16079 (holo. BULU; iso. ANK, GAZI).

SOLANACEAE (6: 437)

■ *Scopolia* Jacq.

**S. carniolica* Jacq. obs. Bot. 1: 32 (1764).

(Başer et al., 2006)

N. E. Anatolia. **A8** Artvin: Borçka, Macahal

Solanum L. (6: 438)

* *S. elaeagnifolium* Cav. Icon. Descr. 3: 22, tab.243 (1795).

(İlçim & Behçet, 2007)

S. Anatolia. **C6** Kahramanmaraş: Türkoğlu, around Fatih small town, Gavurgölü Marsh, 450-500 m, 10.vii.2006, A İlçim 1632. (KSUH and VANF).

**S. sisymbriifolium* Lam. Tabl. Encycl. 2: 25. 1794. Turk J. Bot. 31: 1-3 (2007).

(Karaer & Kutbay, 2007)

North Anatolia. **A5** Amasya: vicinity of Bahçeleri along the valley S slopes, 400 m, 25.vii.1999, F.Karaer 8250, ibid. 20.ix.1999, F.Karaer 8900 (fruiting material), **A6** Samsun: between Toptepe and Directory of Forestry along the valley east-facing slopes, 50-100 m, 7.vi.1995, F.Karaer 6016 ibid. 18.viii.1995, F.Karaer 6495 (fruiting material).

UMBELLIFERAE (4: 265)

■ ■ *Dumaniana* Yıldırımlı & Selvi

(Yıldırımlı & Selvi, 2006)

e **D. gelendostensis Yıldırımlı & Selvi in Ot Sistematis Botanik Der. 13: 5-6 (2006).

Type: (Turkey) B3 Isparta: Gelendost, Akdağ köyü, Kirişli or Çatalgöl dağı, from SE part of mountain to Çatalgöl, association of *Quercus coccifera*, 1000-1500 m, 12.vi.2002, B. Selvi 2420 (holo. Hb. YILDIRIMLI; iso. GAZI).

Ferula L. (4: 440)

e **F. coskunii H.Duman et M.Sağiroğlu in Bot. J. Linn. Soc. 147: 357-359 (2005).

(Duman & Sağıroğlu, 2005)

Type: (Turkey) C6 Hatay: From Hatay to Yayladağı road, West of Şenköy, Ziyaret dağı, 1150-1200 m, clearing forest, calcareous rock, 2.ix.2002, H.Duman 8779 & C.S.Erdurak (holo. GAZI; iso. ANK, HUB, E).

e **F. mervynii M.Sağiroğlu et H.Duman in Bot. J. Linn. Soc. 153: 357-360 (2007).

(Sağıroğlu & Duman, 2007)

Type: (Turkey) A8 Artvin: Yusufeli-Sarıgöl road, from Yusufeli 1-3 km, 650 m, 26.vii.2002, M.Sağiroğlu, 2207 & A.Duran (holo. GAZI; iso. ANK, HUB, Hb YILDIRIMLI, E).

*F. parva Freyn et Bornm. in Öst. Bot. Zeitschr.42: 121 (1892).

(Sağıroğlu & Duman, 2006)

Inner Anatolia. B5 Kırşehir: between Yeşilyurt and Dağ çiftliği, 1400 m, 22.vi.1994, Hamzaoğlu 1553 (ADO); ibid., 29.vi.2001, M.Sağiroğlu 2241 (GAZI); Kırşehir: Çayağzı town, Höyüktepe, 1320 m, 29.vi.2001, M.Sağiroğlu 1723 (GAZI).

Ferulago W. Koch (4: 453)

e **F. glareosa Kandemir et Hedge in Willdenowia 37: 273-275 (2007).

(Kandemir & Hedge, 2007)

Type: (Turkey) B7 Erzincan: Erzincan-Kemah, Sürek, 3.vi.2005, Kandemir 6901 (holo. E; iso. ANK, GAZI).

Peucedanum L. (4: 473)

e **P. longibracteolatum Parolly et Nordt in Willdenowia 35: 98-101 (2005).

(Parolly & Nordt, 2005)

Type: (Turkey) C4 Antalya: Alanya, Mahmutlar-Taşkent, etwa 50 km NE Mahmutlar, Tal des Gökdere

Çayı nahe Çayarası, 1130 m, Steilhang, Exp. S, 1.vii.2003, Ulrich 3130 (holo. B; iso. hb. Parolly).

Prangos Lindley (4: 382)

e **P. ilanae Pimenov, Akalin et Kljuykov in Candollea 60: 379-380 (2005).

(Pimenov et al., 2005)

Type: (Turkey) B1 Balıkesir: Zeytinli, Kazdağı, Mehmetalan village-Sarıkız summit, Ziindere, Girne pass, 800 m, 16.vii.1994, Özhatay & Byfield 70164 (holo. ISTE; iso. G, MW).

e **P. turcica A.Duran, M.Sağiroğlu et H.Duman in Ann. Bot. Fennici 42: 69-71 (2005).

(Duran et al., 2005)

Type: (Turkey) C6 Osmaniye: between Yarpuz and Yağlıpınar, 1500-1700 m, under mixed forest, shady slopes, 13.ix.2003, A.Duran 6362 & M.Sağiroğlu (holo. GAZI; iso. KNYA, ANK, HUB).

URTICACEAE (7: 633)

Urtica L. (7: 633)

U. dioica L., Sp. Pl. 984 (1753).

*subsp. *kurdistanica* Chrtek, Fl. Iranica 105: 3. 1974.

(Weigend, 2006)

Type: Iran, Kurdistan, Chehel Chashmeh, 44 km NE of Dezh-Shahpur (Merivan), ad rivulum, 2000 m, 7.7.1971, K.H.Rechinger 43026 (W 1972-00387).

Turkey: Cappadocia, Mt Argaeus (= Erciyas Dağı), 2100 m, 19.6.1890, Bornmüller 2507 (B); [S Turkey, Taurus,] Iter Cilicum in Tauri alpes Bulgar Dagh, in alpinis vallis Gusguta, 25.6.1853, T.Kotschy exs. Iter Cilicum 7 (G-DC microfiche & scan).

*subsp. *pubescens* (Ledeb.) Domin

(Weigend, 2006)

Turkey: Lydia, Mount Mesogis above Tire, 600-700 m, 14.6.1906, Bornmüller 9946 (B).

**U. fragilis* Thiébaut in Bull. Soc. Bot. France 82: 192. 1935

(Weigend, 2006)

Turkey: (Prov. Hatay, Amanus Mountains) "Bityas, été 1937", Delbés 81 (G).

ZYGOPHYLLACEAE (2: 491)

■ *Tetradiclis* Stev. ex M. Bibb.

**T. tenella* (Ehrenb.) Litw., Trav. Mus. Bot. Acad. Petersb. 3: 122 (1907).

(Hamzaoğlu et al., 2005)

Type: (Egypt) Alexandria (1824). Ehrenberg (K; G, n.v.)

Inner Anatolia. B5 Kırıkkale: Delice, around Tekel Tuzla İşletmesi; 650 m, salty areas, 2.vi.2002, Hamzaoğlu 2859 & A.Duran (ADO); Delice, Tekel Tuzları, abandoned salt ponds, 19.v.2004, Hamzaoğlu 3569 & A.Duran (ADO, GAZI, ANK, HUB, KNYA, hb. Akhani, hb. YILDIRIMLI).

MONOCOTYLEDONES

ARACEAE (8: 41)

Arum L. (8: 43)

**A. megobrebi* Labin, M.Neumann, Bogner et P.C. Boyce in Willdenowia 37: 447-448 (2007).

(Labin et al., 2007)

Type: Georgia, Imeretia, west side of Surami Mountains, near Chewi, W of street-tunnel, W of the pass, 25.v.2007, A.Gröger & W.Lobin (holo. TBI; iso. B, BM, BONN, E, FR, K, LE, M).

N.E. Anatolia. A9 Artvin: above Çamlı, 2420 m, 26.v.1990, M.Koenen 20*90*K (cult. in BG Bonn, accession number 8404-1-1990); Trabzon - Gümüşhane borders, between Kayaçi Köy and Yağmudere, 1550 m, 14.vi.1991, V.Haller & M.Koenen 36*91*TR H*K (cult. in BG Bonn, accession number 9824-1-1991).

CYPERACEAE (9:32)

Carex L. (9: 73)

C. serotina Mérat

*subsp. *serotina* Nouv. Fl. Env. Paris, ed. 22: 54 (1821).

(Greuter & Raus, 1999)

Inner Anatolia. C4 Karaman: N side of Oyüklü Dağ N of Ermene 24.vii.1992, Raab-Straube 0 2-4 (B, hb. P. Hein, hb. Parolly, hb. Raab-Straube).

GRAMINEAE (9: 158)

Arundo L. (9: 561)

* *A. mediterranea* Danin in Willdenowia 34: 362 (2004).

(Greuter & Raus, 2007)

Type: Israel, Nahal Sorek, 8 km east of Gedera, 18.9.2004, blooming culms from present year and previous year's culms with branches, Danin (holo.HUJ; iso. B, BEO, BM, E, G, K, MPU, PAL).

European Turkey. A1 (E) Tekirdağ: 7 km E of Tekirdağ towards İstanbul, between the coastal road and the beach, 3.7.2007, Danin (B, HUJ, PAL).

Glyceria R. Br. (9:535)

e ***G. holubii* Tzvelev in Bot. Zhurn. (Moscow & Leningrad) 91:266 (2006).

(Tsvelev, 2006)

Type: (Turkey) B7 Tunceli: Ovacık-Hozat, 2000 m, flush in oak scrub, 22.vii.1957, Davis a. Hedge N 31535 (holo. LE).

e ***G. lazistanica* Holub ex Tzvelev in Bot. Zhurn. (Moscow & Leningrad) 91: 264 (2006).

(Tsvelev, 2006)

Type: (Turkey) A8 Rize: Khabakhor, vers 1800 m alt., 30.vii.1866, B.Balansa N 1554 (holo. LE; iso. LE).

G. lithuanica (Gorski) Lindm.

e ** var. *pontica* Tzvelev

Bot. Zhurn. (Moscow & Leningrad) 91:260 (2006). (Tsvelev, 2006)

Type: (Turkey) A8 Rize: Khabakhor, 30.vii.1866, B.Balansa N 1555 (holo. LE; iso. LE).

Poa L. (9: 470)

* *P. densa* Troitsky in Bull. Jard. Bot. Princ. URSS, xxvii. 619 (1928).

(Greuter & Raus, 2002)

S.W. Anatolia. C2 Denizli: Honaz Dağı, summit region, 19.vi.2000, Eren & Parolly 7481 (B, ISTE, herb. Parolly); ibid., steep, open, scree-rich limestone swards on raw soil (Astragalo-Brometalia), alt. 2550 m, 20.6.2000, Eren & Parolly 7514 (B, herb. Parolly); ibid., ascent from Honaz to the summit region, 23.6.2000,

Eren 3048 & Parolly (AKDU, GAZI) **C3** Antalya: Bakırlı Dağı above Saklıkent, rocky slopes, thorny cushion and dwarf shrub communities, alt 2200-2550 m, 24.7.1998, Eren 6507 (AKDU, B, herb. Parolly); id., **C3** Isparta: Eğirdir, Barla Dağı, lateral summit above Çamdağı, 30.6.2000, Eren 3342 & Parolly (AKDU, B).

* *P. maroccana* Nannf. in Svensk Bot. Tidskr. 1938, xxxii. 296.

(Greuter & Raus, 1998)

W. Anatolia. **C1** Muğla: Bafa-Gölü/Besparmak Dağı (Latmos-Gebirge), Kaprikiri, Ruinenstätte Herakleia am Latmos, alt. c. 150 m, 17.iii.1998, Kürschner & Parolly 98-26 (B, hb. Parolly).

Secale L. (9: 255)

S. segetale (Zhuk.) Roshev.

* subsp. *segetale* Trudy in Bot. Inst. Akad. Nauk S.S.S.R., Ser. 1, Fl. Sist. Vyssh. Rast. 6: 143 (1947).

(Greuter & Raus, 2002)

E. Anatolia. **A9** Erzurum: E of Horasan, road verge in abandoned agricultural area.

Setaria P. Beauv. (9.597)

**S. pallide-fusca* (Schumach.) Stapf. et Hubb.

(Başak & Guler, 2006)

European Turkey. **A1(E)** Edirne: Trakya University Güllapoğlu Campus, grassy places, R. 50, 01.ix.2001, EDTU 9617.

IRIDACEAE (8: 381)

Crocus L. (8: 413)

C. biflorus Miller, Gard. Dict. ed. 8 no.4 (1768).

e **subsp. *caelestris* Kerndorff et Pasche in Linzer biol. Beitr. 38: 179-181 (2006).

(Kerndorff & Pasche, 2006)

Type: (Turkey) **B2** Uşak: surrounding areas of the city of Uşak, 1200-1400 m, 13.iii.2004, HKEP 0409 (holo. LI).

C. biflorus Miller, Gard. Dict. ed. 8 no.4 (1768).

e ** subsp. *caricus* Kerndorff & Pasche in Linzer Biol. Beitr. 36:7(2004).

(Kerndorff & Pasche, 2004)

Type: (Turkey) **C2** Muğla: Batı Menteşe dağları, 900-1200 m, HKEP 0307, 14.iii.2003 (holo. LI).

C. biflorus Miller, Gard. Dict. ed. 8 no.4 (1768).

e ** subsp. *ionopharynx* Kernd. & Pasche in Linzer Biol. Beitr. 36:6(2004).

(Kerndorff & Pasche, 2004)

Type: (Turkey) **C2** Muğla: Batı Menteşe dağları, 800-1200 m, HKEP 0306, 14.iii.2003 (holo. LI).

C. biflorus Miller, Gard. Dict. ed. 8 no.4 (1768).

e **subsp. *yataganensis* Kerndorff et Pasche in Linzer biol. Beitr. 38:178-179 (2006).

(Kerndorff & Pasche, 2006)

Type: (Turkey) **C2** Muğla, Doğu Menteşe Dağları, 1000-1200 m, 18.iii.2004, HKEP 0402 (holo. LI).

Gladiolus L. (8: 441)

e ***G. attilae* Kit Tan, B.Mathew et A.Baytop in Phytologia Balcanica 12: 71-73 (2006).

(Tan et al., 2006)

Type (Turkey) **B7** Elazığ: on 22.vi.1991, A.Baytop ISTE 62995 (holo. ISTE; iso. K)

Iris L. (8: 382)

e ***I. celikii* Akpulat et K.I. Chr. in Nordic J. Bot. 24: 207 (2006).

(Christensen & Akpulat, 2006)

Type: (Turkey) **B6** Sivas: along the road from Celalli to Hafik, approx. 1 km before the village of Hafik, 4.iv.2003, Akpulat 3205 (holo. C; iso. CUFH).

e ***I. nezahatiae* Güner et H.Duman in Turk J. Bot. 31: 311-313 (2007).

(Güner & Duman, 2007)

Type: (Turkey) **A8** Artvin: Yusufeli, Kılıçkaya (Ersis) yukarısı [above Kılıçkaya (Ersis)], A.Güner 14024-M.Johnson, M.Öztekin, R.Unwin, P.Ravenhill, NGBB accession 2006-401 (holo. Hb. of NGBB; iso. GAZI).

I. reticulata M. Bib.

e **var. *kurdica* Rukšāns in Buried Treasures 363 (2007).

(Rukšāns, 2007)

Type: (Turkey) **B9** Van: 27 km south of Bashkale, 60 km before Hankari, 27.v.2004, Rukšāns, Seisums & Zetterlund BATM 04-118 (holo. GB).

- LILIACEAE (8: 67)
- Allium L. (8: 98)
- e ***A. antalyense* Eren, Çinbilgel et Parolly in Willdenowia 37: 259-265 (2007).
- (Parolly & Eren, 2007)
- Type (Turkey) C3 Antalya: Teke Dağı, ascent to summit region from Belen Yayla (above Akçağal village), gappy *Elytrigia divaricata* swards with thorn-cushions, limestone, 13.7.2004, Eren 272/04 & Çingibel (holo. AYDN; iso. B, GAZI, hb. Parolly).
- A. circinnatum* Sieber
- e **subsp. *evae* R.M.Burton in Ann. Musei Goulandris 11: 81-83 (2006).
- (Burton, 2006)
- Type: (Turkey) C2 Muğla: SE of Öludeniz, near road to Kadak, E.Pirie s.n., iv.1992 (holo. BM).
- Bellevalia Lapeyr* (8: 264)
- e ***B. leucantha* K.Perss. in Bot. J. Linn. Soc. 150: 253-256 (2006).
- (Persson, 2006)
- Type: (Turkey) B7 Elazığ: Kuruca pass 35 km W of Bingöl in grass in very wet meadow near melting snow, 2000 m, 23.v.1998, Sønderhousen 1100 (holo GB).
- Colchicum* L. (8: 329)
- e ***C. antepense* K.Perss. in Bot. Jahrb. Syst. 127: 169-171 (2007).
- (Persson, 2007)
- Type: (Turkey) C6 Gaziantep: In apric. Calc. Aintab, Killis, 2-3000 ped. , l. 1867, Haussknecht 924 (holo. G; iso. FI, M, W, WU.-SC Turkey).
- e ***C. chlorobasis* K.Perss. in Edinb. J. Bot. 62: 182-185 (2005).
- (Persson, 2005)
- Type (Turkey) C4 Konya: SW of Sorkun (WSW of Bozkır), 1730 m, on limestone with *Juniperus*, *Berberis*, *Acantholimon*, *Astragalus*, and *Cyclamen cilicicum*, 1.x.1996, Kerndorff & Pasche 96-09 (holo.GB).
- **C. crocifolium* Boiss, Diagn. Pl. Or. Nov. Ser. 1, 5: 67 (1844).
- (Akan & Eker, 2005)
- S.E. Anatolia. C7 Şanlıurfa: North of Direkli Hill, 19.i.2003, 640 m, rock slopes, Eker 274 & Akan; ibid., 02.ii.2003, 726 m, Eker 278 & Akan.
- e ***C. ignescens* K.Perss., in Bot. Jahrb. Syst. 127: 191-192 (2007).
- (Persson, 2007)
- Type: (Turkey) C7 Şanlıurfa: Karacadağ, northern side, road to radar station, deep volcanic soil, 1650 m, 8.iv.1990, Kammerlander et al. 90-149 (holo. GB.)
- e ***C. lagotum* K.Perss., in Bot. Jahrb. Syst. 127: 197-198 (2007).
- (Persson, 2007)
- Type: (Turkey) B8 Erzurum: Çat to Bingöl, 73 km from Erzurum, 7 km before Cirisli Geçidi, steep unstable serpentine slopes, 1900 m, 1.vi.2004, Zetterlund et al. BATM 223 (holo. GB)
- *Loncomelos* Raf., Fl. tellur. II/1:24 (1837).
- e ***L. tardus* Speta in Phyton 46: 11 (2006).
- (Speta, 2006)
- Type: (Turkey) C2/3 Antalya: Taurus, Montes Cebireis Dağı, prope pagum Gümüşkavak, 500-700m, cult. LI: 17.vi.2000, V.Vašák TO 229 (holo. LI).
- Merendera* Ramond (8: 351)
- e ***M. figalii* Varol in Belg. Jour. Bot. 138: 89-92 (2005).
- (Varol, 2005)
- Type: (Turkey) C2 Muğla: Sandras Dağ, around Kartal Gölü, 1900-2100 m, 05.v.2001, serpentine rocks, open stony places, Varol 4016 (holo. MUĞLA; iso. GAZI).
- Muscaria* Miller (8: 245)
- e ***M. turicum* Uysal, Ertuğrul et Dural in Bot. J. Linn. Soc. 154: 233-236 (2007).
- (Uysal et al., 2007)
- Type: (Turkey) C4 Konya: Bozkır, Tufan Deresi, Avdan yaylası, alpine steppe, 1950-2100 m, 11.v.2005, Ertuğrul 3430, T.Uysal, M.Küçüködük, H.Dural (holo. KNYA; iso. GAZI).
- Ornithogalum* L. (8: 227)
- **O. boucheanum* (Kunth) Asch.
- (Dalgıç et al., 2006)

European Turkey. A1(E): Edirne: Hacıdanışment-Sarıdanışment village, 30.iv.1989, G.Dalgıç (EDTU 3150), Karakasım: Tayakadın village entrance, 29.iv.2004, G.Dalgıç and F.Dane (EDTU 9499).

e ***O. chetikianum* Uysal, Ertuğrul et Dural in Bot. J. Linn. Soc. 148: 501-503 (2005).

(Uysal et al., 2005)

Type: (Turkey) C4 Konya: Bozkır, Yazı Yaylası, 2000 m, 10.v.2003, T.Uysal 207, K.Ertuğrul & H.Dural (holo. KNYA; iso. GAZI).

e ***O. mekseliniae* Varol in Nord. J. Bot. 23: 607-608 (2005).

(Varol, 2005)

Type: (Turkey) C2 Muğla: Muğla to Denizli, 20. km, Yaraş local, 600 m, 09.iii.2003, damp-fields and water meadows, Varol 4249, (holo. MUKLA; iso. GAZI)

e ***O. sumbulianum* O.D.Düşen et I.G.Deniz in Pak. J. Bot. 37: 33-34 (2005).

(Düşen & Deniz, 2005)

Type: (Turkey) C2 Antalya: Elmalı, Çamkuyusu, 1600-1650 m, sparse *Cedrus libani* forests, 15.v.2002 I.G.Deniz (2127) (holo. AKDU; iso. ANK, HUB, GAZI).

Puschkinia Adams (8: 226)

e ***P. peshmenii* Rix et B.Mathew in Curtis Bot. Mag. 24: 56-57 (2007).

(Rix & Mathew, 2007)

Type: (Turkey) C10 Hakkari: Yüksekova, in the gorge leading into the valley from the west, among dwarf shrubs, E.M.Rix et al. 1926 (holo. ISTE; iso. K.).

Scilla L. (8:214)

e ***S. seismsiana* Rukšāns & Zetterlund in Buried Treasures 363 (2007).

(Rukšāns, 2007)

Type: (Turkey) C8/9 Siirt: Siirt-Baykan road, 6 km south from the Silvan-Tatvan junction, 950 m, 25.v.2004, Ruksans, Seisms & Zetterlund BATM 04-055 (holo. GB).

ORCHIDACEAE (8: 450)

Dactylorhiza Necker ex Nevski (8: 535)

D. maculata (L.) Soó in Nom. Nova Gen. *Dactylorhiza*, 7 (1962).

*subsp. *maculata*

(Daşkin et al., 2007)

= *Orchis maculata* L. in Sp. Pl. 2:942 (1753).

Lectotype: Habitat in Europae pratis succulentis, Herb. Linn. No. 1054.36 (LINN, photo).

W. Anatolia. B2 Bursa: Uludağ, between İnegöl and Oylat; Mesruriye village, around Alaçam forest depot, marshy ground, 24.vii.2005, G.Kaynak, Ö.Yılmaz (holo. BULU).

Himantoglossum W.D. Koch (8:507)

H. caprinum Spreng.

e **subsp. *robustissimum* Kreutz in Jour. Eur. Orch. 38:122 (2006).

(Kreutz, 2006)

Type: (Turkey) A3 Bolu: etwa 15 km NE Abant, UTM UL60.49; lockerer Mischwald mit Eichengebüsch und grasigen Freiflächen, 980 m, 23.vi.1994, C.A.J.Kreutz (holo. L.).

Ophrys L. (8:476)

O. aramaeorum P.Delforge

e **f. *akcakarae* (Kreutz) P.Delforge in Naturalistes Belges 87 (Orchid. 19): 260 (2006).

(Delforge, 2006)

basionym: *Ophrys oestrifera* M.Bieb. subsp. *akcakarae* Kreutz Jour. Eur. Orch. 38: 162 (2006).

Type: (Turkey) B8 Diyarbakır: etwa 10 km W Kulp, UTM 37S FC 65300/59670, Eichengebüsch mit niederem Laubwald und grasigen Freiflächen, 840 m, 20.v.2004, C.A.J.Kreutz (holo. L.).

O. argolica H.Fleischm. ex Vierh.

e **subsp. *mandalyana* (B.& H.Baumann) Kreutz in Orchidee 57: 100 (2006).

(Kreutz, 2006)

basionym: *Ophrys ferrum-equinum* Desf. subsp. *mandalyana* B.Baumann & H.Baumann Jour. Eur. Orch. 37: 949 (2005).

basionym: *Ophrys argolica* H.Fleischm. ex Vierh. subsp. *mandalyana* (B..Baumann & H.Baumann) Kreutz Orchidee (Hamburg) 57: 100 (2006)

Type: (Turkey) C2 Muğla: Bodrum, 15.iv.1982,
B.Baumann & H.Baumann (holo. STU).

O. *fusca* Link

e **subsp. *thracica* Kreutz in Ber. Arbeitskrs. Heim.
Orchid. 24: 102 (2007).

(Kreutz, 2007)

Type: (Turkey) A1(E) Çanakkale: Tayfur (Gelibolu-Halbinsel), 28.iv.1997, C.A.J.Kreutz (holo. L).

e **O. x *gennarii* Rchb.f. nothosubsp. *kusadasiensis*
B.Baumann & H.Baumann in Jour. Eur. Orch. 37: 961
(2005).

(Baumann & Lorenz, 2005)

(*Orchis morio* L. subsp. *caucasica* (K.Koch)
E.G.Camus, Bergon & A.Camus x *O. papilionacea* L. subsp.
heroica (E.D.Clarke) H.Baumann)

Type: (Turkey) C1 Aydın: Kusadasi, 05.iv.1982,
B.Baumann & H.Baumann (holo. STU).

e **O. x *rueckbrodtiana* W.Hahn in Ber. Arbeitskrs.
Heim. Orchid. 24: 125 (2007).

(Hahn, 2007)

Type: (Turkey) C2 Muğla: prope urbem Kara aören;
Mezarlık büyük, 8.v.2007, W.Hahn (holo. Hb.
C.A.J.Kreutz).

O. *transhyrcana* Czerniak.

e **subsp. *paphlagonica* Kreutz in Ber. Arbeitskrs.
Heim. Orchid. 24: 119 (2007).

(Kreutz, 2007)

Type: (Turkey) A5 Sinop: Umgebung von Türkeli,
18.v.1996, C.A.J.Kreutz (holo. L).

Serapias L. (8:502)

S. *vomeracea* Briq.

e **subsp. *artemisiae* W.Hahn, Kreutz & Passin in
Ber. Arbeitskrs. Heim. Orchid. 24: 134 (2007).

(Kreutz, 2007)

Type: (Turkey) C2 Muğla: Ortaca (Dalaman),
16.iv.1997, C.A.J.Kreutz (holo. L).

New Taxonomic Status

(○ new synonym, ● new combination, ◆ new status,
■ nomen nova)

CAMPANULACEAE

Campanula L. (6: 2)

● ◆ C. *pamphylica* (Contandr., Quezel & Pamukç.)
E.Akçicek & Vural.

(Akçicek et al., 2005)

Campanula argaea Boiss. & Bal. subsp. *pamphylica*
Contandr., Quezel & Pamukç., in Ann. Univ. Provence Sci.
46: 54. 1972.

● ◆ C. *pamphylica* (Contandr., Quezel & Pamukç.)
subsp. *tokuri* (A.Ocak) E.Akçicek & Vural.

(Akçicek, et al., 2005)

Campanula tokuri A.Ocak, in Israel J.Pl. Sci. 51:
321.2003.

CARYOPHYLLACEAE

Arenaria L. (2: 17)

A. *angustifolia* McNeill

(Parolly & Eren, 2006)

○ *Arenaria yunus-emrei* Aytaç & H.Duman in Ann.
Bot. Fenn. 41: 213-214. 2004.

Minuartia L. (2: 38)

● ◆ M. *elmalia* (Aytaç) Aytaç, Parolly & Eren.

(Parolly & Eren, 2006)

M. dianthifolia subsp. *elmalia* Aytaç in Ann. Bot. Fenn.
41: 216. 2004.

Silene L. (2: 179)

S. *pratensis* (Rafn) Godron & Gren. in Gren. et
Godron, Fl. Fr. 1: 216 (1847).

(McNeill & Prentice, 1981)

○ *Silene alba* (Mill.) E.H.L. Krause in Sturm, Deutschl.
Fl. ed. 2, 5: 98 (1901) (non *S. alba* Muhl. Ex Britton
(1893).)

● S. *pratensis* (Rafn) Gren. & Godr. subsp.
divaricata (Reichnb.) McNeill & H.C.Prent.

(McNeill & Prentice, 1981)

○ *Silene alba* (Mill.) E.H.L. Krause subsp. *divaricata*
(Reichnb.) Walters, Feddes Repert. 69: 48 (1964).

● S. *pratensis* (Rafn) Gren. & Godr. subsp.
eriocalycina (Boiss..) McNeill & H.C.Prent.

(McNeill & Prentice, 1981)

○ *Silene alba* (Mill.) E.H.L.Krause subsp. *eriocalycina* (Boiss.) Walters, Feddes Repert. 69: 48 (1964).

COMPOSITAE

Achillea L. (5: 224)

● *A. santolinoides* Lag. subsp. *wilhelmsii* (K.Koch) Greuter.

(Greuter et al., 2003)

○ *A. wilhelmsii* K.Koch in Linnaea 24: 328 (1851).

Centaurea L. (5: 465)

◆ *C. kotschyi* (Boiss. & Heldr.) Hayek subsp. *persica* (Boiss.) Greuter.

(Greuter, 2003-b)

○ *Centaurea kotschyi* (Boiss. & Heldr.) Hayek var. *persica* (Boiss.) Wagenitz in Bot. Jahrb. Syst. 82: 169 (1963).

Cirsium Miller (5:370)

C. leucocephalum Spreng. subsp. *leucocephalum*.

(Greuter & Raus, 2005)

○ *C. lappaceum* (M.Bieb.) Fisch. subsp. *anatolicum* Petr.

○ *C. penicillatum* K.Koch subsp. *anatolicum* (Petr.) Soldano

C. leucocephalum Spreng. subsp. *penicillatum* (K.Koch) Greuter.

(Greuter & Raus, 2005)

○ *C. lappaceum* (M.Bieb.) Fisch. Cat. Jard. Gorenki, ed. 1812: 35. 1812 (non Lam. 1779).

● *C. leucocephalum* subsp. *tenuilobum* (K.Koch) Greuter.

(Greuter & Raus, 2005)

○ *Cirsium lappaceum* (M.Bieb.) Fisch. subsp. *tenuilobum* (K.Koch) P.H.Davis & Parris in Notes Roy. Bot. Gard. Edinburgh 33: 430 (1975).

Conyza Tamasch. in Fl. URSS 25: 185, 583, (1959).

● *C. squamatus* (Spreng.) Tamamsch.

(Greuter & Raus, 1995)

○ *Aster squamatus* (Spreng.) Hieron

Cota J.Gay in Guss. Fl. Sic. Syn. ii. 866 (1844).

● *C. halophila* (Boiss. & Balansa) Oberprieler & Greuter.

(Greuter et al., 2003)

○ *Anthemis halophila* Boiss. & Balansa in Boissier, Fl. Orient. 3: 285 (1875).

● *C. melanoloma* (Trautv.) Holub subsp. *trapezuntica* (Grierson) Oberprieler & Greuter.

(Greuter et al., 2003)

○ *Anthemis melanoloma* Trautv. subsp. *trapezuntica* Grierson in Notes Roy. Bot. Gard. Edinburgh 33: 411 (1975).

● *C. tinctoria* J.Gay subsp. *euxina* (Boiss.) Oberprieler & Greuter.

(Greuter et al., 2003)

○ *Anthemis tinctoria* L. var. *euxina* (Boiss.) Grierson in Notes Roy. Bot. Gard. Edinburgh 33 (3): 428 (1975).

Cyanus Mill. in Gard. Dict. Abr., ed. 4. (1754); Druce in Rep. Bot. Exch. Cl. Brit. Isles, 3: 431, (1913).

● *C. bourgaei* (Boiss.) Wagenitz & Greuter.

(Greuter, 2003-b)

○ *Centaurea bourgaei* Boiss., Fl. Orient. 3: 637, (1875).

● *C. pichleri* (Boiss.) Holub subsp. *extrarosularis* (Hayek & Siehe) Wagenitz & Greuter

(Greuter, 2003-b)

○ *Centaurea pichleri* Boiss. subsp. *extrarosularis* (Hayek & Siehe) Wagenitz in Notes Roy. Bot. Gard. Edinburgh 33: 231, (1974).

● *C. tchihatcheffii* (Fisch. & C.A.Mey.) Wagenitz & Greuter.

(Greuter, 2003-b)

○ *Centaurea tchihatcheffii* Fisch. & C.A.Mey. in Ann. Sci. Nat., Bot., ser. 4, 1: 31, (1854).

Echinops L. (5: 609)

● *E. spinosissimus* Turra subsp. *bithynicus* (Boiss.) Greuter.

(Greuter, 2003-b)

○ *Echinops viscosus* subsp. *bithynicus* (Boiss.) Rech. f. in Akad. Wiss. Wien, Math.-Naturwiss. Kl., Denkschr. 105 (1): 641(1944).

- Helichrysum** Gaertner (5: 80)
H. chasmolyicum P.H.Davis
(Parolly & Eren, 2006)
○ *Helichrysum orbicularifolium* Sümbül & al. in Bot. J. Linn. Soc. 141: 251. 2003.
- Hieracium** L. (5: 696)
H. transiens Freyn
●♦ subsp. *autranii* (Post) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium autranii* Post in Bull. Herb. Boissier 1: 25, (1893).
●♦ subsp. *artabirense* (Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium artabirense* (Zahn) Üksip in Komarov, Fl. SSSR 30: 74 (1960).
●♦ subsp. *brandisianum* (Prain) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium brandisianum* [Zahn in Magyar Bot. Lapok 5: 83. 1906, nom. inval., ex] Prain, Index Kew., Suppl. 4: 109 (1913).
●♦ subsp. *caloprasinum* (Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium caloprasinum* (Zahn) Üksip in Komarov, Fl. SSSR 30: 73, (1960).
●♦ subsp. *cryptonaevum* (Bornm. & Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium cryptonaevum* (Bornm. & Zahn) P.D.Sell & C.West in Notes Roy. Bot. Gard. Edinburgh 33: 430, (1975).
●♦ subsp. *erythrocarpum* (Peter) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium erythrocarpum* Peter in Nachr. Königl. Ges. Wiss. Göttingen, Math.-Phys. Kl. 1898: 44, (1898).
●♦ subsp. *insolitum* (Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium insolitum* (Zahn) Üksip in Komarov, Fl. SSSR 30: 73 (1960).
●♦ subsp. *karagoellense* (Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium karagoellense* (Zahn) P.D.Sell & C.West in Notes Roy. Bot. Gard. Edinburgh 33: 431 (1975).
●♦ subsp. *karakolense* (Bornm. & Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium karakolense* (Bornm. & Zahn) P.D.Sell & C.West in Notes Roy. Bot. Gard. Edinburgh 33: 431 (1975).
●♦ subsp. *laxifurcans* (Bornm. & Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium laxifurcans* (Bornm. & Zahn) P.D.Sell & C.West in Notes Roy. Bot. Gard. Edinburgh 33: 431 (1975).
●♦ subsp. *mallopoderes* (Bornm. & Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium mallopoderes* (Bornm. & Zahn) P.D.Sell & C.West in Notes Roy. Bot. Gard. Edinburgh 33: 431 (1975).
●♦ subsp. *Sintenisii* (Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium marmoricola* P.D.Sell & C.West in Notes Roy. Bot. Gard. Edinburgh 33: 244 (1974).
●♦ subsp. *stellidorsum* (Bornm. & Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium stellidorsum* (Bornm. & Zahn) P.D.Sell & C.West in Notes Roy. Bot. Gard. Edinburgh 33: 431 (1975).
●♦ subsp. *variegatisquamum* (Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium variegatisquamum* (Zahn) Üksip in Komarov, Fl. SSSR 30: 75 (1960).
- H. jurassicum** Griseb.
●♦ subsp. *barbeyi* (Post) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium barbeyi* Post in Bull. Herb. Boissier 1: 24 (1893).
●♦ subsp. *tamderense* (Hub.-Mor.) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium tamderense* Hub.-Mor. in Notes Roy. Bot. Gard. Edinburgh 33: 246 (1974).

- H. chalcidicum Boiss. & Heldr.
- ♦ subsp. *paphlagonicum* (Freyn & Sint.) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium paphlagonicum* Freyn & Sint. in Österr. Bot. Z. 44: 260 (1894).
- ♦ subsp. *cappadocicum* (Freyn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium cappadocicum* Freyn in Österr. Bot. Z. 41: 55 (1891).
- ♦ subsp. *koenigianum* (Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium koenigianum* (Zahn) Üksip in Komarov, Fl. SSSR 30: 200 (1960).
- ♦ H. *pannosum* Boiss. subsp. *strigulosum* (Post) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium strigulosum* Post in Bull. Herb. Boissier 1: 25 (1893).
- ♦ H. *caucasiense* Arv.-Touv. ex Lipsky subsp. *biebersteinii* (Litv. & Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium biebersteinii* Litv. & Zahn in Repert. Spec. Nov. Regni Veg. 4: 329 (1907).
- ♦ H. *brevifolium* Tausch subsp. *noëanum* (Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium noëanum* Zahn in Engler, Pflanzenr. 79: 1006 (1922).
- ♦ H. *camkoriense* Zahn subsp. *sarykamyschense* (Üksip) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium sarykamyschense* Üksip in Bot. Mater. Gerb. Bot. Inst. Komarova Akad. Nauk SSSR 19: 487 (1959).
- H. *lachenalii* Suter in Helvet. Fl. 2: 145 (1802).
- ♦ subsp. *argillaceoides* (Litv. & Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium argillaceoides* (Litv. & Zahn) Üksip in Komarov, Fl. SSSR 30: 269 (1960).
- ♦ subsp. *macrophyllum* (Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium macrophyllum* (Zahn) Üksip in Komarov, Fl. SSSR 30: 255, (1960).
- ♦ subsp. *subhastulatum* (Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium subhastulatum* (Zahn) Üksip in Komarov, Fl. SSSR 30: 257 (1960).
- ♦ H. *laevigatum* Willd. subsp. *hypopityforme* (Üksip) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium hypopityforme* Üksip in Bot. Mater. Gerb. Bot. Inst. Komarova Akad. Nauk SSSR 19: 495 (1959).
- ♦ H. *levicaule* Jord. subsp. *leptogrammoides* (Üksip) Greuter
(Greuter & Raab-Straube, 2007)
- *Hieracium leptogrammoides* Üksip in Bot. Mater. Gerb. Bot. Inst. Komarova Akad. Nauk SSSR 19: 502 (1959).
- ♦ H. *murorum* L. subsp. *gentiliforme* (Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium gentiliforme* (Zahn) P.D.Sell & C.West in Notes Roy. Bot. Gard. Edinburgh 33: 431 (1975).
- ♦ H. *racemosum* Waldst. & Kit. ex Willd. subsp. *subartvinense* (Üksip) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium subartvinense* Üksip in Bot. Mater. Gerb. Bot. Inst. Komarova Akad. Nauk SSSR 19: 468 (1959).
- ♦ H. *schmidtii* Tausch subsp. *argaeum* (Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium argaeum* (Zahn) P.D.Sell & C.West in Notes Roy. Bot. Gard. Edinburgh 33: 430 (1975).
- ♦ subsp. *giresunense* (Hub.-Mor.) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium giresunense* Hub.-Mor. in Notes Roy. Bot. Gard. Edinburgh 33: 245 (1974).

- ♦ subsp. *huber-morathii* (P.D.Sell & C.West) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium huber-morathii* P.D.Sell & C.West in Notes Roy. Bot. Gard. Edinburgh 33: 244 (1974).
- ♦ subsp. *labillardierei* (Arv.-Touv.) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium labillardierei* Arv.-Touv. in Annuaire Conserv. Jard. Bot. Genève 1: 101 (1897).
- ♦ subsp. *lasiochaetum* (Bornm. & Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium lasiochaetum* (Bornm. & Zahn) P.D.Sell & C.West in Notes Roy. Bot. Gard. Edinburgh 33: 431 (1975).
- ♦ subsp. *leucothecum* (Freyn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium leucothecum* Freyn in Österr. Bot. Z. 42: 343, (1892).
- ♦ subsp. *pseudodontotrichum* (Hub.-Mor.) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium pseudodontotrichum* Hub.-Mor. in Notes Roy. Bot. Gard. Edinburgh 33: 246 (1974).
- ♦ subsp. *subvandasii* (Bornm. & Zahn) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium subvandasii* (Bornm. & Zahn) P.D.Sell & C.West in Notes Roy. Bot. Gard. Edinburgh 33: 431 (1975).
- ♦ *H. sericophyllum* Nejceff & Zahn subsp. *tmoleum* (Hub.-Mor.) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium tmoleum* Hub.-Mor. in Notes Roy. Bot. Gard. Edinburgh 33: 247 (1974).
- ♦ *H. sparsum* Friv. subsp. *beypazariense* (Hub.-Mor.) Greuter.
(Greuter & Raab-Straube, 2007)
- *Hieracium beypazariense* Hub.-Mor. in Notes Roy. Bot. Gard. Edinburgh 33: 425 (1975).
- Jacobaea* Mill. in Gard. Dict. Abr., ed. 4. (1754).
- *J. cilicia* (Boiss.) B.Nord.
(Greuter & Raab-Straube, 2007)
- *Senecio cilicius* Boiss., Diagn. Pl. Orient., ser. 2, 3: 37, (1856).
- *J. inops* (Boiss. & Balansa) B.Nord.
(Greuter & Raab-Straube, 2007).
- *Senecio inops* Boiss. & Balansa, Diagn. Pl. Orient., ser. 2, 6: 101, (1859).
- ♦ *J. maritima* (L.) Pelser & Meijden subsp. *bicolor* (Willd.) B.Nord. & Greuter.
(Greuter & Raab-Straube, 2006)
- *Senecio bicolor* (Willd.) Tod., Ind. Sem. Horti Panorm. 1859: 30, (1860) (non Viv. 1802).
- *J. sandrasica* (P.H.Davis) B.Nord. & Greuter.
(Greuter & Raab-Straube, 2006)
- *Senecio sandrasicus* P.H.Davis in Notes Roy. Bot. Gard. Edinburgh 22: 75, (1956).
- *J. trapezuntina* (Boiss.) B.Nord.
(Greuter & Raab-Straube, 2007).
- *Senecio trapezuntinus* Boiss., Fl. Orient. 3: 393, (1875).
- Klasea* Cass. in Dict. Sc. Nat. xxxv. 173 (1825); xli. 321.
- *K. aznavouriana* (Bornm.) Greuter & Wagenitz.
(Greuter, 2003-b)
- *Serratula aznavouriana* Bornm. in Bull. Herb. Boissier, ser. 2, 6: 233 (1906).
- *K. bornmuelleri* (Azn.) Greuter & Wagenitz.
(Greuter, 2003-b)
- *Serratula bornmuelleri* Azn. in Repert. Spec. Nov. Regni Veg. 11: 397 (1912).
- *K. cerinthifolia* (Sm.) Greuter & Wagenitz.
(Greuter, 2003-b)
- *Serratula cerinthifolia* (Sm.) Boiss., Fl. Orient. 3: 585, (1875).
- *K. erucifolia* (L.) Greuter & Wagenitz.
(Greuter, 2003-b)
- *Serratula erucifolia* (L.) Druce in Bot. Exch. Club Soc. Brit. Isles 3: 424 (1914).

- *K. grandifolia* (P.H.Davis) Greuter & Wagenitz.
(Greuter, 2003-b)
- *Serratula grandifolia* P.H.Davis in Notes Roy. Bot. Gard. Edinburgh 33: 293 (1974).
- *K. hakkiarica* (P.H.Davis) Greuter & Wagenitz.
(Greuter, 2003-b)
- *Serratula hakkiarica* P.H.Davis, Fl. Turkey 10: 235 (1988).
- *K. kotschyi* (Boiss.) Greuter & Wagenitz.
(Greuter, 2003-b)
- *Serratula kotschyi* Boiss., Fl. Orient. 3: 590 (1875).
- *K. kurdica* (Post) Greuter & Wagenitz.
(Greuter, 2003-b)
- *Serratula kurdica* Post, Pl. Post. 4: 8, (1892).
- *K. lasiocephala* (Bornm.) Greuter & Wagenitz.
(Greuter, 2003-b)
- *Serratula lasiocephala* Bornm. in Repert. Spec. Nov. Regni Veg. 5: 167, (1908).
- *K. oligocephala* (DC.) Greuter & Wagenitz.
(Greuter, 2003-b)
- *Serratula oligocephala* DC., Prodr. 6: 669, (1838).
- *K. quinquefolia* (Willd.) Greuter & Wagenitz.
(Greuter, 2003-b)
- *Serratula quinquefolia* Willd., Sp. Pl. 3: 1639, (1803).
- *K. serratuloides* (DC.) Greuter & Wagenitz.
(Greuter, 2003-b)
- *Serratula serratuloides* (DC.) Takht. in Tahtadzjan & Fedorov, Fl. Erevana: 323, (1945).
- *K. haussknechtii* (Boiss.) Holub (Martins, 2006)
- *Serratula haussknechtii* Boiss., Fl. Orient. 3: 589 (1875).
- *K. radiata* (Waldst. & Kit.) A.Löve & D. Löve (Martins, 2006)
- *Serratula radiata* (Waldst. & Kit.) M.Bieb., Fl. Taur.-Caucas. 3: 545 (1819).
- *K. radiata* (Waldst. & Kit.) A.Löve & D. Löve subsp. *biebersteiniana* (Grossh.) Greuter.
- (Greuter & Raab-Straube, 2005)
- *Serratula radiata* (Waldst. & Kit.) M.Bieb. subsp. *biebersteiniana* Grossh., Fl. Kavk. 4: 194 (1934).
- *K. coriacea* (Fisch. & C.A. Mey. ex DC.) Holub (Martins, 2006)
- *Serratula coriacea* Fisch. & C. A. Mey. ex DC., Prodr. 6: 668 (1838).
- Podospermum* DC. in Fl. Franc. (DC. & Lamarck), ed. 3. 4: 61, (1805).
- *P. laciniatum* DC. subsp. *decumbens* (Guss.) Gemeinholzer & Greuter.
(Greuter & Raab-Straube, 2006)
- *Scorzonera laciniata* L. subsp. *decumbens* (Guss.) Greuter in Willdenowia 33: 237, (2003).
- ♦ *P. radicosum* (Boiss.) Gemeinholzer & Greuter.
(Greuter & Raab-Straube, 2006)
- *Scorzonera cana* (C.A.Mey.) Hoffm. var. *radicosa* (Boiss.) Chamberlain in Notes Roy. Bot. Gard. Edinburgh 33(3): 433, (1975).
- *Scorzonera radicosa* Boiss., Diagn. Pl. Orient. 11: 43, (1849).
- Psephellus* Cass (5: 466)
- *P. goeksunensis* (Aytaç & H.Duman) Greuter & Raab-Straube.
(Greuter & Raab-Straube, 2006)
- *Centaurea goeksunensis* Aytaç & H.Duman in Pakistan J. Bot. 37: 563, (2005).
- *P. yildizii* (Civelek et al.) Greuter.
(Greuter, 2003-d)
- Centaurea yildizii* Civelek et al. in Bot. J. Linn. Soc. 143: 207 (2003).
- Reichardia* Roth (5:693)
- R. dichotoma* (DC.) Freyn.
(Greuter, 2003-c)
- *Reichardia glauca* V.A. Matthews in Notes Roy. Bot. Gard. Edinburgh 34: 125. 1975, nom. illeg.
- Rhaponticoides* Vaill. in Phys. Abh. Königl. Akad. Wiss. Paris 5: 165 (1754).
- *R. amasiensis* (Bornm.) M.V.Agab. & Greuter.
(Greuter, 2003-b)

- *Centaurea amasiensis* Bornm. in Repert. Spec. Nov. Regni Veg. 3: 54 (1906).
- *R. amplifolia* (Boiss. & Heldr.) M.V.Agap. & Greuter.
(Greuter, 2003-b)
- *Centaurea amplifolia* Boiss. & Heldr. in Boissier, Diagn. Pl. Orient., ser. 2, 3: 68 (1856).
- *R. iconiensis* (Hub.-Mor.) M.V.Agap. & Greuter.
(Greuter, 2003-b)
- *Centaurea iconiensis* Hub.-Mor. in Bauhinia 7: 77 (1981).
- *R. mykalea* (Hub.-Mor.) M.V.Agap. & Greuter.
(Greuter, 2003-b)
- *Centaurea mykalea* Hub.-Mor. in Bauhinia 6: 370 (1979).
- Xanthium L.* (5: 47)
- *X. orientale* L. subsp. *italicum* (Moretti) Greuter.
(Greuter, 2003-d)
- Xanthium italicum* Moretti in Giorn. Fis., ser. 2, 5: 326. 1822
- Xanthium italicum* Moretti subsp. *italicum* [per elak., Prodr. Fl. Böhmen: 186. 1871]
- Xanthium saccharatum* subsp. *italicum* (Moretti) Hayek in Repert. Spec. Nov. Regni Veg. Beih. 30(2): 613.1931
- Xanthium strumarium* subsp. *italicum* (Moretti) D.Löve in Bot. J. Linn. Soc. 71:271.1976
- Xanthium echinatum* subsp. *italicum* (Moretti) O.Bolòs & Vigo in Collect. Bot. (Barcelona) 17: 90.1988.
- Xanthium saccharatum* Wallr., Beitr. Bot.: 238.1844;
= *Xanthium cavanillesii* Schouw in Ann. Sci. Nat., Bot., ser. 3, 12: 357. 1849; & Ind. Sem. Hort. Acad. Haun. 1849: 14. 1849-1850
- Xanthium strumarium* subsp. *cavanillesii* (Schouw) D.Löve & Dans. in Canad. J. Bot. 37: 205. 1959.
- Scorzonera L.* (5: 632)
- *S. laciniata* L. subsp. *decumbens* (Guss.) Greuter.
(Greuter, 2003-c)
- *Scorzonera laciniata* L. subsp. *calcitrapifolia* (Vahl) Maire in Bull. Soc. Hist. Nat. Afrique N. 22: 302 (1931).
- Senecio L.* (5: 145)
- ♦ *S. leucanthemifolius* Poir. subsp. *vernalis* (Waldst. & Kit.) Greuter.
(Greuter, 2003-d)
- *Senecio vernalis* Waldst. & Kit., Descr. Icon. Pl. Hung. 1: 23, (1800).
- Sonchus L.* (5: 690)
- *S. bulbosus* (L.) N.Kilian & Greuter subsp. *microcephalus* (Rech.f.) N.Kilian & Greuter.
(Greuter, 2003-c)
- *Aetheorhiza bulbosa* L. subsp. *microcephala* Rech. f. in Phyton (Horn) 16: 217.
- Tragopogon L.* (5: 657)
- ♦ *T. porrifolius* L. subsp. *longirostris* (Sch. Bip.) Greuter.
(Greuter, 2003-c)
- *Tragopogon longirostris* Sch. Bip. in Webb & Berthelot, Hist. Nat. Iles Canaries 3(2,2): 469 (1850).
- Tripolium* Nees in Gen. Sp. Aster. 10, 152 (1832).
- *T. pannonicum* (Jacq.) Dobrocz. subsp. *tripolium* (L.) Greuter.
(Greuter, 2003-a)
- *Aster tripolium* L., Sp. Pl.: 872 (1753).
- CRUCIFERAE** (1: 248)
- Barbarea R.Br.* (1: 433)
- B. brachycarpa* Boiss.
- ♦ subsp. *anfractuosa* (Hartvig & Strid) Parolly & Eren.
(Parolly & Eren, 2006)
- Barbarea minor* K.Koch var. *anfractuosa* Hartvig & Strid in Bot. Jahrb. Syst. 108: 316 (1987).
- B. brachycarpa* Boiss.
subsp. *brachycarpa*
var. *brachycarpa*
(Parolly & Eren, 2006)
- *Barbarea minor* K.Koch var. *libanotica* Bornm. in Verh. Zool.-Bot. Ges. Wien 48: 550 (1898).
- *Barbarea minor* K.Koch var. *eriopoda* N.Busch, Fl. Cauc. Crit. 3 (4): 318 (1909).

○ *Barbarea hedgeana* Kit Tan & Gemici in Edinb J Bot 47: 287 (1990).

B. brachycarpa Boiss.

● ♦ subsp. *robusta* (Coode & Cullen) Parolly & Eren.

(Parolly & Eren, 2006)

Barbarea minor K.Koch var. *robusta* Coode & Cullen in Notes Roy. Bot. Gard. Edinburg 26: 190 (1965).

B. brachycarpa Boiss.

● ♦ subsp. *minor* (K.Koch) Parolly & Eren.

(Parolly & Eren, 2006)

Barbarea minor K.Koch in Linnaea 19: 55.1847.

var. *minor* (Parolly & Eren, 2006)

○ *Barbarea alpina* K.Koch in Linnaea 19: 319 (1847).

Draba L. (1: 413)

D. imeretica (Rupr.) Rupr.

(Parolly & Eren, 2007)

○ *D. terekemensis* Yild. in Ot 7: 1-6 (2000).

Noccaea Moench

● *N. cariensis* (Carlström) Parolly, Nordt & Aytaç (Aytaç et al., 2006)

(basionym: *Thlaspi cariense* Carlström in Willdenowia 16 (1): 73. 1986)

● ♦ *N. sintenisii* subsp. *crassum* (P.H.Davis) Parolly.

(Parolly & Eren, 2006)

Thlaspi crassum P.H.Davis in Fl. Turk. 10: 235. 1988

Thlaspi sintenisii Hausskn. ex Bornm. subsp. *crassum* (P.H.Davis) Parolly

Zuvanda (F.Dvorak) Askerova

♦ *Z. exacoides* (DC.) Askerova.

(Parolly & Eren, 2007)

Sisymbrium exacoides DC., Syst. Veg. 2: 463. 1821

Malcolmia exacoides (DC.) Spreng., Syst. Veg., ed. 16, 2: 899.1825.

Conringia nana Boiss. & Kotschy in Boissier, Diagn. Pl. Orient., ser. 2, 5: 25.1856.

Malcolmia conringioides Boiss., Fl. Orient. 1: 230.

1867.

DIPSACACEAE

Cephalaria Schroder ex Roemer et Schultes (4: 585)

C. balansae Raus in Willdenowia 35(1): 59 (2005).

(Greuter & Raus, 2005)

○ *C. dipsacoides* Boiss. & Balansa (non Kar. & Kir. in Bull. Soc. Imp. Naturalistes Moscou 14: 434. 1841) in Boissier, Diagn. Pl. Orient., ser. 2, 5: 107, (1859).

EUPHORBIACEAE

Chamaesyce S.F.Gray (7: 571)

● *C. canescens* (L.) Prokh. (Parolly & Eren, 2007)

Euphorbia canescens L., Sp. Pl. Ed. 2: 652 (1762)

Euphorbia chamaesyce L., Sp. Pl. 455 (1753)

● *C. forskalii* (J.Gay) Parolly.

(Parolly & Eren, 2007)

Euphorbia forskalii J.Gay in Webb & Berth., Hist. Nat. Canaries 3 (2, 3): 240(1847).

Euphorbia aegyptiaca Boiss., Cent. Euphorb.: 13. 1860, nom. illeg

● *C. humifusa* (Willd.) Prokh.

(Parolly & Eren, 2007)

Euphorbia humifusa Willd. in Enum. Pl. Horti Berol. Suppl. 27 (1813).

● *C. maculata* (L.) J.K.Small.

(Parolly & Eren, 2007)

Euphorbia maculata sensu auct. Plur., non L. (1753).

Euphorbia supina Rafin. in Amer. Monthly Mag. 2: 119 (1817).

● *C. nutans* (Lag.) J.K.Small.

(Parolly & Eren, 2007)

Euphorbia nutans Lag., Gen. Sp. Pl.: 17. (1816)

● *C. peplis* (L.) Prokh.

(Parolly & Eren, 2007)

Euphorbia peplis L. in Sp. Pl. 455 (1753).

- *C. prostrata* (Aiton) J.K.Small.
(Parolly & Eren, 2007)
Euphorbia prostrata Aiton, Hort. Kew. ed. 1, 2: 139
(1789)
- *C. serpens* (Kunth) J.K.Small.
(Parolly & Eren, 2007)
Euphorbia serpens Kunth in Humb., Bonpl. & Kunth,
Nov. Gen. Sp. 2: 52 (1817).
- LABIATAE**
- Salvia* L. (7: 400)
- *S. absconditiflora* Greuter & Burdet in Willdenowia
15: 77 (1985).
(Greuter & al., 1986)
S. cryptantha Montbret & Aucher ex Benth. in Ann.
Sci. Nat. Bot. Ser. 2, 6: 40 (1836) (non *Salvia*
cryptanthos Schultes, Observ. Bot.: 12. 1809).
- LEGUMINOSAE**
- Astragalus* L. (3: 49)
- ◆ *A. lycaonicus* Hub.-Mor. & Reese, Feddes Report
48: 282 (1940).
(Ekici et al., 2005)
A. mesogitanus Boiss., Diagn. Ser. 1 (2): 41 (1843).
Cytisopsis Jaub. & Spach (3: 511)
- *C. pseudocytisus* (Boiss.) Fertig (Greuter & Raus,
1989)
- *C. dorycnifolia* Jaub. & Spach subsp. *reeseana*
(Guyot) Huber-Morath in P.H.Davis, Fl. Turkey 3: 511
(1970).
- Genista* L. (3: 24)
- ◆ *Genista januensis* subsp. *lydia* (Boiss.) Kit Tan
& Zielinski.
(Zielinski et al., 2004)
G. lydia Boiss., Diagn. Pl. Orient. ser. 1, 2: 8. 1843.
G. lydia Boiss. var. *lydia* in Flora of Turkey and the
East Aegean Islands 3: 26. 1970.
- G. leptophylla* Spach, Ann. Sci. Nat., ser. 3, 3: 127.
1845.
- G. antiochia* Boiss., Diagn. Pl. Orient. 9: 4. 1849.
- G. patula* M.Bieb. var. *antiochia* (Boiss.) Boiss., Fl.
Orient. 2: 44. 1867.
- G. lydia* Boiss. var. *antiochia* (Boiss.) P.E.Gibbs, Flora
of Turkey and the East Aegean Islands 3: 26. 1970.
- G. lydia* Boiss. subsp. *antiochia* (Boiss.) Ponert, Feddes
Repert. 83: 619. 1973.
- Medicago* L. (3: 483)
- *M. monantha* Blatt subsp. *noëana* (Boiss.)
Greuter & Burdet.
(Greuter & Raus, 1989)
- *Trigonella monantha* C.A.Mey. subsp. *noëana*
(Boiss.) Hub.-Mor. in P.H.Davis, Fl. Turkey 3: 475
(1969).
- Ononis* L. (3: 373)
- ◆ *O. spinosa* L. subsp. *arvensis* (L.) Greuter & Burdet
(Greuter & Raus, 1989)
- *Ononis arvensis* L. in Syst. Nat. ed. 10: 1159,
(1759).
- LINACEAE**
- Linum* L. (2: 425)
- L. hirsutum* L., Sp. Pl. ed. 2, 398 (1762).
◆ subsp. *platyphyllum* (P.H.Davis) Yilmaz & Kaynak
(Yilmaz & Kaynak, 2006)
- L. hirsutum* L. subsp. *anatolicum* (Boiss.) Hayek var.
platyphyllum P.H.Davis, Notes Royal Bot. Garden
Edinburgh 22: 142. (1957).
- L. pamphylicum* Boiss. & Heldr. ex Planch., London
J. Bot. vii. 511 (1848).
- ◆ subsp. *papilliferum* (Hub.-Mor. & Reese)
Yilmaz & Kaynak.
(Yilmaz & Kaynak, 2006)
- Linum papilliferum* Hub.-Mor. & Reese in Feddes
Repert. 52 (1): 45. 1943.
- L. mucronatum* Bertol. subsp. *gypsicola* P.H.Davis var.
papilliferum (Hub.-Mor. & Reese) P.H.Davis, Notes Royal
Bot. Garden Edinburgh 22: 158 (1967).

POLYGONACEAE

Persicaria (L.) Miller Gard. Dict. Abr., ed. 4. (1054) 1754.

● *P. bistorta* Sampaio subsp. *carnea* (K.Koch) Greuter & Burdet.

(Greuter & Raus, 1989)

Polygonum bistorta L. subsp. *carneum* (K.Koch) Coode & Cullen in Notes Roy. Bot. Gard. Edinburgh 27: 215 (1967).

PLUMBAGINACEAE

Acantholimon Boiss. (7: 478)

A. acerosum (Willd.) Boiss., Diagn. Ser. 1 (7): 80 (1846).

● ♦ subsp. *brachystachyum* (Boiss.) Dogan & Akaydin.

(Doğan & Akaydin, 2007)

Syn.: *A. acerosum* (Willd.) Boiss. var. *brachystachyum* Boiss. Fl. Or. 4: 838 (1879); *A. tchihatcheffii* Fisch. & Mey., Ann. Sci. Nat. Ser. 4, 1: 30 (1843); *A. phrygium* Boiss., Diagn. Ser. 1 (7): 79 (1846); *A. pinardii* Boiss., loc. Cit. (1846); *A. listoniae* Boiss. in DC., Prodr. 12: 631 (1848); *A. acerosum* (Willd.) Boiss. var. *pinardii* (Boiss.) Mobayen, Revis. Taxon. *Acantholimon*: 179 (1964). Ic: Tchihat., Asia Min. Atlas t.36 (1860).

● ♦ *A. multiflorum* (Bokhari) Dogan & Akaydin.

(Doğan & Akaydin, 2007)

Syn.: *A. spirizianum* Mobayen var. *multiflorum* Bokhari, Notes R.B.G. Edinb. 32: 70 (1972).

● ♦ *A. parviflorum* (Bokhari) Akaydin & Dogan.

(Doğan & Akaydin, 2007)

Syn.: *A. caryophyllaceum* Boiss. var. *parviflorum* Bokhari, Notes R.B.G. Edinb. 30: 299 (1970).

A. puberulum Boiss. & Bal. in Boiss., Diagn, ser, 2 (4): 62 (1859).

● ♦ subsp. *longiscapum* (Bokhari) Dogan & Akaydin.

(Doğan & Akaydin, 2007)

Syn.: *A. puberulum* var. *longiscapum* Bokhari, Notes R.B.G. Edinb. 30: 300 (1970).

A. puberulum Boiss. & Bal. in Boiss., Diagn, ser, 2 (4): 62 (1859).

● ♦ subsp. *peroninii* (Boiss.) Akaydin & Dogan.

(Doğan & Akaydin, 2007)

Syn.: *A. peroninii* Boiss., Fl. Or. 4: 842 (1879).

A. puberulum Boiss. & Bal. in Boiss., Diagn, ser, 2 (4): 62 (1859).

● ♦ subsp. *puberulum*.

(Doğan & Akaydin, 2007)

Syn.: *A. puberulum* Boiss. & Bal. var. *glabrum* Bokhari, Notes R.B.G. Edinb. 30: 300 (1970).

A. quinquelobum Bunge, Mem. Acad. Sci. Petersb. Ser. 7, 18 (2): 56 (1872).

● ♦ var. *curviflorum* (Bunge) Dogan & Akaydin.

(Doğan & Akaydin, 2007)

Syn.: *A. curviflorum* Bunge in Mem. Acad. Sci. Petersb. Ser. 7, 18 (2): 57 (1872).

RANUNCULACEAE

Anemone L. (1: 134)

● *A. narcissifolia* L. subsp. *willdenowii* (Boiss.) Greuter & Burdet.

(Greuter & Raus, 1989)

A. narcissiflora L. subsp. *willdenowii* (Boiss.) P.H.Davis in Notes Roy. Bot. Gard. Edinburgh 26: 175 (1965).

Delphinium L. (1: 108)

● ♦ *D. fissum* Waldst. & Kit. subsp. *albiflorum* (DC.) Greuter & Burdet.

(Greuter & Raus, 1989)

D. albiflorum DC., Syst. Nat. 1: 353 (1817).

Ranunculus L. (1: 146)

R. peltatus Schrank

○ *R. kastamonuensis* A.Dönmez.

(Parolly & Eren, 2007)

■ *R. kurdicus* Greuter & Burdet.

(Greuter & Raus, 1989)

R. diversifolius Boiss. & Kotschy in Boiss., Fl. Orient. 1: 45, 1867 [non Schrank, Baier. Fl. 2: 103, 1789].

ROSACEAE

Crataegus L. (4: 133)

C. rhipidophylla Gand.

var. *rhipidophylla*

(Dönmez, 2007)

○ *C. curvisepta* Lindman in Svensk Fanerogamfl. 307 (1918). (Dönmez, 2004 & 2007)

● ♦ var. *browicziana* (K.I.Chr.) Dönmez

(Dönmez, 2007)

○ *Crataegus x browicziana* K.I.Christ, Syst. Bot. Monogr. 38: 153-154 (1992).

UMBELLIFERAE

Dichoropetalum Fenzl in Pug. Pl. Nov. Syr. 17. (1842).

● *D. carvifolia* (Vill.) Pimenov & Kljuykov.

(Pimenov & al., 2007)

○ *Peucedanum carvifolia* Vill. Prosp. Hist. Pl. Dauphiné 25 (1779).

● *D. alpinum* Fenzl in Pug. Pl. Nov. Syr. 17 (1842).

(Pimenov & al., 2007)

○ *Johrenia alpina* (Fenzl) Fenzl in Russegger Reisen 1: 961. t. 17 (1843).

● *D. alpigenum* (Boiss.) Pimenov & Kljuykov.

(Pimenov & al., 2007)

○ *Peucedanum alpigenum* Boiss., Fl. Orient. 2: 1020 (1872).

● *D. aureum* (Boiss.) Pimenov & Kljuykov.

(Pimenov & al., 2007)

○ *Johrenia aurea* Boiss. & Balansa ex Boiss., in Diagn. Pl. Orient. ser. 2, 6: 81 (1859).

● *D. graminifolium* (Boiss.) Pimenov & Kljuykov.

(Pimenov & al., 2007)

○ *Peucedanum graminifolium* Boiss., in Ann. Sci. Nat., Bot., ser. 3, 1: 314 (1844).

● *D. isauricum* (Parolly & Nordt) Pimenov & Kljuykov.

(Pimenov & al., 2007)

○ *Peucedanum isauricum* Parolly & Nordt, in Willdenowia 34(1): 481 (2004).

● *D. depauperatum* (Boiss. & Balansa ex Boiss.) Pimenov & Kljuykov.

(Pimenov & al., 2007)

○ *Peucedanum depauperatum* Boiss. & Balansa ex Boiss., in Diagn. Pl. Orient., ser. 2, 5: 98 (1856).

○ *Johrenia berytea* Boiss. & Hausskn. ex Boiss., Fl. Orient. 2: 1012 (1872).

D. seseloides (C.A.Mey.) Pimenov & Kljuykov.

(Pimenov & al., 2007)

○ *Peucedanum meyeri* (Boiss.) Boiss., Fl. Orient. 2: 1018 (1872).

● *D. chryseum* (Boiss. & Heldr. ex Boiss.) Pimenov & Kljuykov

(Pimenov & al., 2007)

○ *Peucedanum chryseum* (Boiss. & Heldr. ex Boiss.) D.F. Chamb., in Davis, Fl. Turkey 4: 475 (1972).

○ *Peucedanum chrysanthum* Boiss., Diagn. Pl. Orient., ser. 2, 6: 86 (1859).

● *D. palimboides* (Boiss.) Pimenov & Kljuykov.

(Pimenov & al., 2007)

○ *Peucedanum palimboides* Boiss., Fl. Orient. 2: 1021 (1872).

Johrenia DC.

J. selinoides Boiss. & Balansa ex Boiss., Diagn. Pl. Orient., ser. 2, 5: 99 (1856).

(Pimenov & al., 2007)

Tordylium L. (4: 504)

● *T. trachycarpum* (Boiss.) Al-Eisawi & Jury.

(Al-Eisawi & Jury, 1988)

○ *Ainsworthia trachycarpa* Boiss., Diagn. ser. 1 (10): 43 (1849).

URTICACEAE (7: 633)

Urtica L. (7: 633)

U. dioica L., Sp. Pl. 984 (1753). subsp. *dioica*

○ *U. haussknechtii* Boiss. in Fl. Orient. 4: 1146 (1879).

(Weigend, 2006)

MONOCOTYLEDONES

GRAMINEAE (9: 158)

Elytrigia Desv. (9: 206)

- *E. erosiglumis* (Melderis) Valdés & H.Scholz.
(Parolly & Eren, 2006)

Elymus erosiglumis Melderis in Notes R.B.G. Edinb. 42: 78 (1984).

Zingeria P. Smirnov (9: 363)

- ◆ *Z. trichopoda* (Boiss.) P.Smirn.

(Parolly & Eren, 2006)

Z. biebersteiniana (Claus) P.Smirn. subsp. *trichopoda* (Boiss.) R. R. Mill in Fl. Turkey & E. Aegean Is., 9: 365 (1985).

Bromopsis (Dumont.) Fourr. in Ann. Soc. Linn. Lyon sér. 2, 17: 187 (1869).

- *B. cappadocica* (Boiss. & Balansa) Holub
subsp. *sclerophylla* (Boiss.) H.Scholz & Valdés.
(Valdés & Scholz, 2006)

○ *Bromus cappadocicus* Boiss. & Bal. subsp. *sclerophyllus* (Boiss.) P.M.Smith in Notes RBG Edinb. 42:496 (1985).

- *B. erecta* Fourr.

(Valdés & Scholz, 2006)

○ *Bromus erectus* Huds. in Flora Anglica 39 (1762).

- *B. sipylea* (Boiss.) H.Scholz & Valdés.

(Valdés & Scholz, 2006)

○ *Bromus sipyleus* Boiss., Diagn. Pl. Orient., ser. 1, 3: 65 (1854).

- *B. tomentella* (Boiss.) Holub

(Valdés & Scholz, 2006)

○ *Bromus tomentellus* Boiss. in Diagn. Pl. Orient. ser. 1, 7: 126, (1846).

- *B. tomentella* (Boiss.) Holub subsp. *nivalis* (Bornm.) H.Scholz.

(Valdés & Scholz, 2006)

○ *Bromus tomentellus* Boiss. subsp. *nivalis* (Bornm.) H.Scholz & A.J.Byfield in Turk. J. Bot. 24(4): 263 (2000).

Elytrigia Desv. in Nouv. Bull. Soc. Philom. ii. 190 (1810).

- *E. caespitosa* (K.Koch) Nevski subsp. *gypsicola* (Melderis) Valdés & H.Scholz.

(Valdés & Scholz, 2006)

○ *Elymus nodosus* (Nevski) Melderis subsp. *gypsicolus* Melderis in Notes Roy. Bot. Gard. Edinburgh 42: 82 (1984).

- *E. caespitosa* (K.Koch) Nevski subsp. *platyphylla* (Melderis) Valdés & H.Scholz.

(Valdés & Scholz, 2006)

○ *Elymus nodosus* (Nevski) Melderis subsp. *platyphyllus* Melderis in Notes Roy. Bot. Gard. Edinburgh 42: 80 (1984).

- *E. clivorum* (Melderis) Valdés & H.Scholz.

(Valdés & Scholz, 2006)

○ *Elymus clivorum* Melderis in Notes Roy. Bot. Gard. Edinburgh 42: 77 (1984).

- *E. elongata* (Host ex Beauv.) Nevski subsp. *salsa* (Melderis) Valdés & H.Scholz.

(Valdés & Scholz, 2006)

○ *Elymus elongatus* (Host) Runemark subsp. *salsus* Melderis in Notes Roy. Bot. Gard. Edinburgh 42: 77 (1984).

- *E. elongata* (Host ex Beauv.) Nevski subsp. *turcica* (McGuire) Valdés & H.Scholz.

(Valdés & Scholz, 2006)

○ *Elymus elongatus* (Host) Runemark subsp. *turcicus* (McGuire) Melderis in Notes Roy. Bot. Gard. Edinburgh 42: 81 (1984).

- *E. elongata* (Host ex Beauv.) Nevski subsp. *pontica* (Podp.) Gamisans in J.Gamisans & D.Jeanmonod, Cat. Pl. Vasc. Corse (ed. 2): 243 (1993).

(Valdés & Scholz, 2006)

○ *Elymus elongatus* (Host) Runemark subsp. *ponticus* (Podp.) Melderis in Bot. J. Linn. Soc. 76 (4): 377 (1978). [= *Triticum ponticum* Podp. Verh. Zool.-Bot. Ges. Wien 52: 68 (1902).

- *E. erosiglumis* (Melderis) Valdés & H.Scholz.

(Valdés & Scholz, 2006)

○ *Elymus erosiglumis* Melderis in Notes Roy. Bot. Gard. Edinburgh 42: 78 (1984).

- *E. lazica* (Boiss.) Valdés & H.Scholz subsp. *lazica*.
(Valdés & Scholz, 2006)
- *Elymus lazicus* (Boiss.) Melderis subsp. *lazicus* in Notes Roy. Bot. Gard. Edinburgh 42: 79 (1984).
- *E. lazica* (Boiss.) Valdés & H.Scholz subsp. *divaricata* (Boiss. & Balansa) Valdés & H.Scholz.
(Valdés & Scholz, 2006)
- *Elymus lazicus* (Boiss.) Melderis subsp. *divaricatus* (Boiss. & Balansa) Melderis in Notes Roy. Bot. Gard. Edinburgh 42: 79 (1984).
- *E. lazica* (Boiss.) Valdés & H.Scholz subsp. *lomatolepis* (Melderis) Valdés & H.Scholz.
(Valdés & Scholz, 2006)
- *Elymus lazicus* (Boiss.) Melderis subsp. *lomatolepis* Melderis in Notes Roy. Bot. Gard. Edinburgh 42: 79 (1984).
- Ochlopoa* (Asch. & Graebn.) H.Scholz in Ber. Inst. Landschafts Pflanzenökol. Univ. Hohenheim Beih. 16: 58 (2003).
- *O. speluncarum* (J.R.Edm.) H.Scholz & Valdés.
(Valdés & Scholz, 2006)
- *Poa speluncarum* Edmonson in Davis, Fl. Turkey 9: 623 (1985).
- *O. supina* (Schrad.) H.Scholz & Valdés.
(Valdés & Scholz, 2006)
- *Poa supina* Schrad., Fl. Germ. 289 (1806).
- *Allium elmaliense* I.G.Deniz & Sümbül.
(Parolly & Eren, 2006)
- A. scabriflorum* Boiss.
- *Allium vuralii* Kit Tan in Pl. Syst. Evol. 155: 102 (1987).
(Parolly & Eren, 2007).
- Colchicum* L. (8: 329)
- *C. figalii* (Ö.Varol) Parolly & Eren.
(Parolly & Eren, 2007; Persson, 2007).
- Merendera figalii* Ö.Varol in Belg. J. Bot. 138: 89 (2005).
- *C. manissadjianii* (Azn.) K.Perss.
(Persson, 2007)
- Merendera manissadjianii* Azn., in Bull. Herb. Boissier ser. 2, 8: 248 (1908).
- C. szovitsii* Fish. & C.A.Mey.
subsp. *brachyphyllum* (Boiss. & Hausskn.) K.Perss.
(Persson, 2007)
- C. brachyphyllum* Boiss. & Hausskn., in Boiss., Fl. Orient. 5: 164 (1882)
- C. fasciculare* (L.) R.Br. var. *brachyphyllum* (Boiss. & Hausskn.) Stef., in Sborn. Balg. Akad. Nauk., Klon Prir.-Mat. 22: 29 (1926)
- C. uliginosum* Boiss. & Hausskn., in sched., nom. nud.

Acknowledgements

We are grateful to the following botanists for their valuable contribution: Zeki Aytaç, Hayri Duman, Tuna Ekim, Mustafa Keskin, Mine Koçyiğit, Nermin Orcan, Gerald Parolly, Martyn Rix, and Jerzy Zieliski.

References

- Adams RP (2004). *Juniperus deltoides*, a new species, and nomenclatural notes an *Juniperus polycarpos* and *J. turcomanica* (Cupressaceae). *Phytologia* 86: 49-53.
- Akan H & Eker İ (2005). A new record for Turkey: *Colchicum crocifolium*, with a contribution to the description of the species. *Belg J Bot* 138: 93-96.
- Akçicek E, Vural M, Açık L & Celebi A (2005). Notes on *Campanula argaea* group (Campanulaceae) in Turkey and related species. *Ann Bot Fennici* 42: 405-410.
- Akkemik Ü, Akalin E & Özhatay N (2007). *Ranunculus anatolicus* sp. nov. (Ranunculaceae) from northeast Turkey. *Nord J Bot* 25:311-314.
- Akpulat HA & Çelik N (2007). A new subspecies of *Astragalus* (Section Caprini DC. Fabaceae) from Turkey. *Bot J Linn Soc* 155: 227-230.
- Aksoy N, Göktürk SR, Açık L & Celebi A (2007). *Cephalaria duzceensis* (Dipsacaceae) a new species from the western Black Sea region Turkey. *Nord J Bot* 25: 64-69.

- Aktoklu E (2001). Two new varieties and a new record in *Onobrychis* from Turkey. *Turk J Bot* 25: 359-363.
- Al-Eisawi D & Jury S (1988). A taxonomic revision of the genus *Tordylium* L. (Apiaceae). *Bot J Linn Soc* 97: 357-403.
- Alan S, Ocak A & Duman H (2007). *Calamintha pamphylica* subsp. *alanyense* (Lamiaceae) a new subspecies from South Anatolia, Turkey. *Ann Bot Fennici* 44: 309-314.
- Ataşlar E & Ocak A (2005). *Gypsophila osmangaziensis* (Caryophyllaceae), a new species from Central Anatolia, Turkey. *Ann Bot Fennici* 42: 57-60.
- Aytaç Z & Duman H (2005). A new species of *Centaurea* L. (Compositae) from Turkey. *Pak J Bot* 37: 563-566.
- Aytaç Z & Mill RR (2005). Two new species of Boraginaceae (Tribe Cynoglosseae) from Turkey. *Edinb J Bot* 61: 109-118.
- Aytaç Z, Nordt B & Parolly G (2006). A new species of *Noccea* (Brasicaceae) from South Anatolia, Turkey. *Bot Jour Linn Soc* 150: 409-416.
- Bani B & Adıgüzel N (2006). A new genus record for Turkey: *Asperuginoides* Rauschert. *International J Bot* 2: 440-442.
- Başak N & Guler N (2006). A new record for the flora of Turkey: *Setaria pallide-fusca* (Schumach.) Stapf et Hubb. (Poaceae) IV. Balkan Botanical Congress, 20-26 June 2006, Abstract book, p. 162-163.
- Başer KHC, Koyuncu M & Vural M (2006). Macahel'e Bir Gezi ve Türkiye Florasına Bir Katkı. *Bağbahçe Derg* 5: 28-31.
- Baumann H & Lorenz R (2005). Beiträge zur Taxonomie europäischer und mediterraner Orchideen. *Jour Eur Orch* 37: 939-974.
- Binzet R & Orcan N (2007). A new species of *Onosma* (Boraginaceae) from Southern Turkey. *Novon* 17: 8-10.
- Budak Ü, Hamzaoğlu & Aksoy A (2007). A new for the Flora of Turkey: *Tephroseris cladobotrys* (Ledeb.) Griseb. & Schenk (Senecioneae, Asteraceae). *Turk J Bot* 31: 265-268.
- Burton R (2006). A new subspecies of *Allium circinnatum* Sieber from S.W. Turkey. *Ann. Musei Goulandris* 11: 81-84.
- Christensen K I & Akpulat H A (2006). *Iris celikii* (Iridaceae) a new species from North-eastern Turkey. *Nord J Bot* 24: 207-210.
- Chrték J (2004). A new subspecies of *Plantago uliginosa* from Central Anatolia (Turkey). *Cas Nar Muz Praze, Rada Prir* 173: 89.
- Coşkunçelebi K, Terzioğlu S & Vladimirov (2007). A new alien species for the Flora of Turkey: *Bidens frondosa* L. (Asteraceae). *Turk J Bot* 31: 1-3.
- Çelik N, Akpulat A & Dönmez E (2007). A new species of *Physoptychis* (Brassicaceae) from central Anatolia, Turkey. *Bot J Linn Soc* 154: 393-396.
- Dalgaç G, Dane F & Aksoy OD (2006). A new record for the flora of Turkey: *Ornithogalum boucheanum* (Kunth) Asch. (Hyacinthaceae), IV. Balkan Botanical Congress, 20-26 June 2006, Abstract book, p. 167.
- Dane F & Ekici N (2006). A new record for the flora of Turkey: *Delphinium balcanicum* Pawl. (Ranunculaceae), IVth Balkan Botanical Congress, 20-26 June 2006, Abstract book, p. 167.
- Daşkin R, Yılmaz Ö & Kaynak G (2006). Presence of *Cirsium eriophorum* (L.) Scop. (Asteraceae) in Turkey. *Turk J Bot* 30: 1-5.
- Daşkin R, Yılmaz Ö & Kaynak G (2007). A new record for the flora of Turkey: *Dactylorhiza maculata* (L.) Soó (Orchidaceae). *J Biol Environ Sci* 1: 11-14.
- Davis PH (ed.) (1965-1985). *Flora of Turkey and the East Aegean Islands*. vol. 1-9. Edinburgh: Edinburgh University Press.
- Davis PH, Tan K & Mill RR (1988). *Flora of Turkey and the East Aegean Islands*. vol. 10 (Suppl. I). Edinburgh: Edinburgh University Press.
- Delforge P (2006). Nouvelles contributions taxonomiques et nomenclaturales aux Orchidées d'Europe. *Natural. Belges (Orchid.)* 19: 258-261.
- Deniz İG & Düşen OD (2004). *Silene sumbuliana* (Caryophyllaceae) a new species from SW Anatolia, Turkey. *Ann Bot Fennici* 41: 293-296.
- Dinç M & Doğan HH (2006). *Stachys yıldırımlıi* (Lamiaceae), a new species from south Anatolia, Turkey. *Ann Bot Fennici* 43: 143-147.
- Dirmenci T (2005). A new subspecies of *Nepeta* (Lamiaceae) from Turkey. *Bot J Linn Soc* 147: 229-233.
- Dirmenci T, Satılı F & Tümen G (2005). A new record for the flora of Turkey: *Iberis saxatilis* L. (Brassicaceae). *Turk J Bot* 29: 471-474.
- Dirmenci T, Satılı F & Tümen G (2006). A new species of *Matthiola* R. Br. (Brassicaceae) from Turkey. *Bot J Linn Soc* 151: 431-435.
- Doğan M & Akaydin G (2005). A new species of *Acantholimon* Boiss. sect. *Glumaria* Boiss. (Plumbaginaceae) from Elazığ, Turkey. *Bot J Linn Soc* 149: 351-356.
- Doğan M & Akaydin G (2007). Synopsis of Turkish *Acantholimon* Boiss. (Plumbaginaceae). *Bot J Linn Soc* 154: 397-419.
- Dönmez A A (2004). The genus *Crataegus* L. (Rosaceae) with special reference to hybridisation and biodiversity in Turkey. *Turk J Bot* 28: 29-37.
- Dönmez A A (2005). A new species of *Crataegus* (Rosaceae) from Turkey. *Bot J Linn Soc* 148: 245-249.
- Dönmez A & Dönmez E (2005). *Crataegus turcicus* (Rosaceae), a new species from NE Turkey. *Ann Bot Fennici* 42: 61-65.
- Dönmez A A (2006). *Teucrium chasmophyticum* Rech.f. (Lamiaceae): A new record for the flora of Turkey. *Turk J Bot* 30: 1-4.
- Dönmez A A (2007). Taxonomic notes on the genus *Crataegus* (Rosaceae) in Turkey. *Bot J Linn Soc* 155: 231-240.
- Dorofeyev V I, Korotaev B A & Gültkin L (2004). A new species of the genus *Stroganowia* Kar. et Kir. (Cruciferae) from Northeast Turkey and Rhynchophorous beetles (Coleoptera, Curculionoidea) associated with it. *Byull Moskovsk Obshch Isp Prir, Otd Biol* 109: 72-76.
- Duman H & Sağıroğlu M (2005). A new species of *Ferula* (Apiaceae) from South Anatolia, Turkey. *Bot J Linn Soc* 147: 357-361.
- Dural H, Tugay O, Ertuğrul K, Uysal T & Demirelma H (2007). *Astragalus turkmenensis* (Fabaceae) a new species from Turkey. *Ann Bot Fennici* 44: 399-402.

- Duran A & Aytaç Z (2005). *Astragalus nezaketae* (Fabaceae), a new species from Turkey. *Ann Bot Fennici* 42: 381-385.
- Duran A & Ocak A (2005). *Hesperis turkmendaghensis* (sect. *Hesperis*) (Cruciferae/Brassicaceae) a new species from the Central Anatolia region, Turkey. *Bot J Linn Soc* 147: 239-247.
- Duran A & Hamzaoglu E (2005). *Psephellus turcicus* sp. nova. (Asteraceae), a new chasmophyte species from central Anatolia, Turkey. *Bot J Linn Soc* 148: 495-500.
- Duran A, Sağıroğlu M & Duman H (2005). *Prangos turcica* (Apiaceae) a new species from South Anatolia, Turkey. *Ann Bot Fennici* 42: 67-72.
- Düzen OD & Deniz IG (2005). *Ornithogalum sumbulianum* (Hyacinthaceae), a new endemic species from South West Anatolia. *Pak J Bot* 37: 33-36.
- Ehrendorfer F & Guo YP (2005). Changes in the circumscription of the genus *Achillea* (Compositae-Anthemideae) and its subdivision. *Willdenowia* 35: 49-54.
- Ekici M, Aytaç Z & Akan H (2005). Contribution to the genus *Astragalus* (Fabaceae) in Turkey: *Astragalus wagneri*, *Astragalus lycocinus* and their relations. *Belg J Bot* 138: 192-198.
- Eren Ö (2007). The genus *Rhaponticooides* Vaill. (Asteraceae) in Turkey: a new species and first key. *Pl Syst Evol* 267: 13-23.
- Ertekin AS (2006). *Astragalus babacianum* (Fabaceae), a new species from Turkey. *Ann Bot Fennici* 43: 291-294.
- Genc Ecevit G, Kandemir A & Genc İ (2007). A new species of *Silene* (Caryophyllaceae) from east Anatolia, Turkey. *Nord J Bot* 25(1-2): 58-63.
- Greuter W, Burdet HM & Long G (1986). "Med-Checklist", *conservatoire et Jardin Botaniques, Ville de Genève Med-Checklist Trust of OPTIMA*, Geneve, Vol. 3, 313.
- Greuter W & Raus T (1989). Med-Checklist Notulae, 15. *Willdenowia* 19: 27-48.
- Greuter W & Raus T (1995). Med-Checklist Notulae, 16. *Willdenowia* 25: 171-176.
- Greuter W & Raus T (1998). Med-Checklist Notulae, 17. *Willdenowia* 28: 163-174.
- Greuter W & Raus T (1999). Med-Checklist Notulae, 18. *Willdenowia* 29: 51-67.
- Greuter W & Raus T (2001). Med-Checklist Notulae, 20. *Willdenowia* 31: 319-328.
- Greuter W & Raus T (2002). Med-Checklist Notulae, 21. *Willdenowia* 32: 195-208.
- Greuter W (2003-a). The Euro+Med treatment of *Astereae* (Compositae) – generic concepts and required new names. *Willdenowia* 33: 45-47.
- Greuter W (2003-b). The Euro+Med treatment of *Cardueae* (Compositae) – generic concepts and required new names. *Willdenowia* 33: 49-61.
- Greuter W (2003-c). The Euro+Med treatment of *Cichorieae* (Compositae) – generic concepts and required new names. *Willdenowia* 33: 229-238.
- Greuter W (2003-d). The Euro+Med treatment of *Senecioneae* and the minor (Compositae) – tribes – generic concepts and required new names, with an addendum to *Cardueae*. *Willdenowia* 33: 245-251.
- Greuter W, Oberprieler C & Vogt R (2003). The Euro+Med treatment of *Anthemideae* (Compositae) – generic concepts and required new names. *Willdenowia* 33: 37-43.
- Greuter W & Raus T (2005). Med-Checklist Notulae, 23. *Willdenowia* 35: 55-64.
- Greuter W & von Raab-Straube E (2005). Euro+Med Notulae, 1. *Willdenowia* 35: 223-239.
- Greuter W & Raus T (2006). Med-Checklist Notulae, 24. *Willdenowia* 36: 719-730.
- Greuter W & von Raab-Straube E (2006). Euro+Med Notulae, 2. *Willdenowia* 36: 707-717.
- Greuter W & von Raab-Straube E (2007). Euro+Med Notulae, 3. *Willdenowia* 37: 139-189.
- Greuter W & Raus T (2007). Med-Checklist Notulae, 25. *Willdenowia* 37: 205-213.
- Güner A, Özhatay N, Ekim T & Başer KHC (2000). *Flora of Turkey and the East Aegean Islands* (Supplement II) vol. 11. Edinburgh: Edinburgh University Press.
- Güner A & Duman H (2007). A new juno *Iris* from North-east Anatolia, Turkey. *Turk J Bot* 31: 311-315.
- Hahn W (2007). *Ophrys x rueckbrodtiana*, nothospecies nat. Nova; eine Ophrys-Hybride aus der Südtürkei. *Ber Arbeitskrs Heim Orchid* 24: 122-126.
- Hamzaoglu E, Duran A & Akhani H (2005). A new genus record for the Flora of Turkey: *Tetradiclis* Stev. ex M. Bieb. (Zygophyllaceae). *Turk J Bot* 29: 403-407.
- Hamzaoglu E, Duran A & Pinar NM (2005). *Salvia anatolica* (Lamiaceae), a new species from East Anatolia, Turkey. *Ann Bot Fennici* 42: 215-220.
- Huber-Morath A (1971). Novitiae Florae Anatolicae IX. *Bauhinia* 4: 197.
- İlçim A & Behçet L (2006). *Geranium kalenderianum* (Geraniaceae) a new species from Turkey. *Ann Bot Fennici* 43: 451-455.
- İlçim A & Behçet L (2007). *Solanum elaeagnifolium* Cav. in Turkey. *Turk J Bot* 31: 1-3.
- Kandemir A (2007). A new *Campanula* (Campanulaceae) from east Anatolia, Turkey. *Nord J Bot* 25: 53-57.
- Kandemir A & Hedge IC (2007). An anomalous new *Ferulago* (Apiaceae) from eastern Turkey. *Willdenowia* 37: 273-276.
- Karaer F & Kutbay GH (2007). *Solanum sisymbriifolium* Lam. (Solanaceae): A new record for Turkey. *Turk J Bot* 31: 1-3.
- Karavelioğulları FA, Vural M & Polat H (2006). Two new taxa from Central Anatolia, Turkey. *Isr J Plant Sci* 54: 105-111.
- Kaya Z & Vural M (2007). A new species of *Centaurea* Sect. Acrocentron (Asteraceae) from Turkey. *Novon* 17: 198-201.
- Kaynak G, Daşkin R, Yılmaz Ö & Erdoğan E (2006). *Verbascum yurtkuranianum* (Scrophulariaceae), a new species from northwest Anatolia, Turkey. *Ann Bot Fennici* 43: 456-459.

- Kerndorff H & Pasche E (2004). Two new taxa of the *Crocus biflorus* Aggregate (Liliiflorae, Iridaceae) from Turke. *Linzer biol Beitr* 36/1: 5-10.
- Kerndorff H & Pasche E (2006). *Crocus biflorus* (Liliiflorae, Iridaceae) in Anatolia (Part three). *Linzer biol Beitr* 38/1: 165-187.
- Keskin M (2005). Türkiye de *Trifolium* L. cinsinden yeni bir takson kaydı ve bir türün varlığı. *Ot Sistematisk Botanik Dergisi* 12: 21-24.
- Keskin M & Çırıcı A (2006). İstanbul çevresinin *Cerastium* L. (Caryophyllaceae) türleri üzerinde taksonomik araştırmalar. *Ot Sistematisk Botanik Dergisi* 13: 71-96.
- Keskin M & Sadikoğlu (2007). *Micromeria persica* Boiss. (Labiatae): A new record for Turkey. *J Fac Pharm Istanbul* 39.
- Kreutz CAJ (2006). Bemerkungen zu den Orchideen von Libanon, Syrien, Israel, Zypern und der Türkei. *Jour Eur Orch* 38: 105-160.
- Kreutz CAJ (2006). Orchideen Europas-Beiträge, umkombinationen und kurze Bemerkungen. *The Orchidee* 57: 99-104.
- Kreutz CAJ (2007). Beitrag zur Taxonomic und Nomenklatur europäischer, mediterraner, nordafrikanischer und vorderasiatischer Orchideen. *Ber Arbeitskr Heim Orchid* 24: 77-141.
- Kuş S & Göktürk S (2005). A new *Cephalaria* species (Dipsacaceae) from the European part of Turkey. *Nord J Bot* 23: 427-430.
- Lobin W, Neumann M, Bogner J & Boyce PC (2007). A new *Arum* species (Areae, Araceae) from NE Turkey and Georgia. *Willdenowia* 37: 445-449.
- Martins L (2006). Systematics and biogeography of *Klasea* (Asteraceae-Cardueae) and a synopsis of the genus. *Bot J Linn Soc* 152: 435-464.
- Mc Neill J & Prentice HC (1981). *Silene pratensis* (Rafn.) Godr. & Gren. The correct name for white campion or white cockle (*Silene alba* (Mill.) E.H.L.Krause, nom. illeg. *Taxon* 30: 27-32.
- Mutlu B (2006). *Saponaria bargyliana* Gombault (Caryophyllaceae): A new record for Turkey and analysis of its morphological characters with related species. *Turk J Bot* 30: 63-70.
- Neff P (2005). Neue der Gattung *Sempervivum* (Crassulaceae) in der Türkei. *Kakteen und andere Sukkulanten* 56: 71-73.
- Orcan N (2006). *Alyssum mughlai* (Brassicaceae) a new species from Southwest Anatolia. *Nord J Bot* 23: 703-705.
- Orcan N & Binzet R (2006). A new record for the flora of Turkey: *Alyssum idaeum* Boiss. & Heldr. *Pak J Bot* 38: 931-933.
- Özen F & Aka GE (2007). A new record for the Flora of Turkey: *Lupinus angustifolius* L. subsp. *reticulatus* (Desv.) Coutinho (Fabaceae). *IJNES* 1: 33-35.
- Özgökçe F, Tan K & Stevanovic V (2005). A new subspecies of *Silene acaulis* (Caryophyllaceae) from East Anatolia, Turkey. *Ann Bot Fennici* 42: 143-149.
- Özgökçe F & Ünal M (2007). A new record for Turkey: *Malcolmia exacoides* (DC.) Spreng. (Brassicaceae). *Turk J Bot* 31: 345-347.
- Özhatay N, Kültür Ş & Aksoy N (1994). Check-list of additional taxa to the Supplement Flora of Turkey. *Turk J Bot* 18: 497-514.
- Özhatay N, Kültür Ş & Aksoy N (1999). Check-list of additional taxa to the Supplement Flora of Turkey II. *Turk J Bot* 23: 151-169.
- Özhatay N & Kültür Ş (2006). Check-list of additional taxa to the Supplement Flora of Turkey III. *Turk J Bot* 30: 281-316.
- Parolly G & Nordt B (2005). A further new *Peucedanum* species (Apiaceae) from the Taurus Mts, Turkey. *Willdenowia* 35: 97-105.
- Parolly G, Eren Ö & Nordt B (2005). *Lomelosia solymica* (Dispsacaceae), a new chasmophyte from the Western Taurus Mts, Turkey. *Willdenowia* 35: 107-115.
- Parolly G & Eren Ö (2006). Contributions to the flora of Turkey 1. *Willdenowia* 36: 823-844.
- Parolly G & Tan K (2006). A new species of *Hesperis* (Brassicaceae) from SW Anatolia, Turkey. *Willdenowia* 36: 851-856.
- Parolly G & Eren Ö (2007). Contributions to the flora of Turkey 2. *Willdenowia* 37: 243-271.
- Parolly G & Tan K (2007). *Verbascum lindae* (Scrophulariaceae), a new species from SW Anatolia, Turkey. *Willdenowia* 37: 277-282.
- Persson K (2005). A new Turkish species of *Colchicum* (Colchicaceae) related to *C. boissieri*. *Edinb J Bot* 62: 181-192.
- Persson K (2006). One new and one amended species of *Bellevalia* (Hyacinthaceae) from Turkey. *Bot J Linn Soc* 150: 253-260.
- Persson K (2007). Nomenclatural synopsis of the genus *Colchicum* (Colchicaceae), with some new species and combinations. *Bot Jahrb Syst* 127: 165-242.
- Pimenov MG, Akalın E & Kljukov E (2005). *Prangos ilanae* (Umbelliferae) a new species from Western Turkey. *Candollea* 60: 379-386.
- Pimenov MG, Kljukov EV & Ostromova TA (2007). Critical taxonomic analysis of *Dichoropetalum*, *Johrenia*, *Zeravschania* and related genera of Umbelliferae-Apioideae-Peucedaneae. *Willdenowia* 37: 465-502.
- Prada C & Rolleri CH (2005). A new species of *Isoetes* (Isoetaceae) from Turkey, with a study of microphyll intercellular pectic protuberances and their potential taxonomic value. *Bot J Linn Soc* 147: 213-228.
- Riedl H, Binzet R & Orcan N (2005). A new species of *Onosma* (Boraginaceae-Lithospermeae) from Southern Turkey. *Edinb J Bot* 61: 127-130.
- Rix M & Mathew B (2007). *Puschkinia peshmenii*. *Curtis s Botanical Magazine* 24: 54-57.
- Ruksans J (2007). Buried Treasures, Finding and Growing the world's choicest bulbs, Portland, Oregon: Timber Press.
- Sağiroğlu M & Duman H (2006). *Ferula parva* Freyn & Bornm. (Apiaceae) A contribution to an Enigmatic Species from Turkey. *Turk J Bot* 30: 399-404.
- Sağiroğlu M & Duman H (2007). *Ferula mervynii* (Apiaceae) a distinct new species from north-east Anatolia, Turkey. *Bot J Linn Soc* 153: 357-362.
- Sáez L & Aldasoro JJ (2003). A taxonomic revision of *Campanula* L. subgenus *Sicyocodon* (Feer) Damboldt and subgenus *Megalocalyx* Damboldt (Campanulaceae). *Bot J Linn Soc* 141: 215-241.

- Simon C (1977). *Bolanthus huber-morathii*, nov. spec. Aus Kleinasien. *Bauhinia* 6: 83-84.
- Speta F (2006). Die Gattung *Loncomelos* Raf. (*Hyacinthaceae-Ornithogaloideae*), vorgestellt anhand dreier neuer Arten. *Phytton* 46: 1-25.
- Sutor K (2005). A new species of *Cynoglossum* (Boraginaceae-Cynoglosseae) from Eastern Turkey. *Edinb J Bot* 61: 119-126.
- Tan K, Mathew B & Baytop A (2006). *Gladiolus attilae* (Iridaceae), a new species from East Anatolia, Turkey. *Phytologia Balcanica* 12: 71-73.
- Tan K & Zeitlinger J (2007). *Astragalus friederickeanus* Kit Tan & Zeitlinger (Fabaceae), a new species from south Anatolia, Turkey. *Wulfenia* 14: 133-136.
- Terzioğlu S & Coşkunçelebi K (2005). A contribution to the Flora of Turkey: *Padus avium* Mill. subsp. *petraea* (Tausch) Pawłowski ex Holub. *Turk J Bot* 29: 251-253.
- Tsvelev NN (2006). Synopsis of the managrass genus, *Glyceria* (Poaceae). *Bot Zhurn (Moscow and Leningrad)*. 91: 255-276.
- Tuzlaci E (2005). Bodrumda Bitkiler ve Yaşam. Ali Rıza Başkan Güzel Sanatlar Matbaası A.Ş. İstanbul.
- Tuzlaci E & Bulut-Emre G (2006). Türkiye Florası için yeni bir takson: *Saxifraga adscendens* L. subsp. *parnassica*, XVI. Bitkisel İlaç Hammaddeleri toplantısı (BİHAT 2006) Bildiri Özeti Kitabı, s.53.
- Uysal T, Ertuğrul K & Dural H (2005). A new species of *Ornithogalum* (Liliaceae) from South Anatolia, Turkey. *Bot J Linn Soc* 148: 501-504.
- Uysal T, Ertuğrul K, Dural H & Küçüködük M (2007). *Muscari turicum* (Liliaceae/Hyacinthaceae), a new species from south Anatolia, Turkey. *Bot J Linn Soc* 154: 233-236.
- Uysal T, Demirelma H, Ertuğrul K, Garcia-Jacas N & Susana A (2007). *Centaurea glabro-auriculata* (Asteraceae) a new species from Turkey. *Ann Bot Fennici* 44: 219-222.
- Uzunhisarcıklı ME, Tekşen M & Doğan E (2005). *Centaurea marashica* (Asteraceae), a new species from Turkey. *Ann Bot Fennici* 42: 309-312.
- Uzunhisarcıklı ME, Doğan E & Duman H (2007). A new species of *Centaurea* L. (Cardueae: Asteraceae) from Turkey. *Bot J Linn Soc* 153: 61-66.
- Ünal M & Özgökçe F (2007). A new genus (*Leptaleum* DC.) record for Turkey. *Turk J Bot* 31: 337-339.
- Ünal M, Özgökçe F, Behçet L & Karabacak O (2007). A new record for Turkey: *Lepidium pinnatifidum* Ledeb (Brassicaceae). *Turk J Bot* 31: 1-2.
- Valdés B & Scholz H (2006). The Euro+Med treatment of *Gramineae* – a generic synopsis and some new names. *Willdenowia* 36: 657-669.
- Varol Ö (2005). *Merendera figalii* (Colchicaceae) a new species from Southwestern Anatolia, Turkey. *Belg J Bot* 138: 89-92.
- Varol Ö (2005). *Ornithogalum mekselinae* (Liliaceae), a new species from south-western Anatolia, Turkey. *Nord J Bot* 23: 607-609.
- Vladimirov V, Dane F, Nikolic T, Stevanovic V & Tan K (2006). New floristic records in the Balkans: 2. *Phytologia Balcanica* 12: 279-301.
- Vladimirov V, Dane F, Stevanovic V & Tan K (2007). New floristic records in the Balkans: 6. *Phytologia Balcanica* 13: 305-327.
- Vural M, Duman H, Aytaç Z & Adıgüzel N (2006). *Saponaria karapınarensis*, *Senecio salsuginea* and *Centaurea tuzgoluensis*, three new species from central Anatolia, Turkey. *Belg J Bot* 139: 252-260.
- Wagenitz G, Hellwig FH, Parolly G & Martins L (2006). Two new species of *Centaurea* (Compositae, Cardueae) from Turkey. *Willdenowia* 36: 423-435.
- Weigend M (2006). *Urtica dioica* subsp. *cypria*, with a re-evaluation of the *U. dioica* group (Urticaceae) in western Asia. *Willdenowia* 36: 811-822.
- Yaltırık F, Eliçin G & Terzioğlu S (2007). *Juniperus oxycedrus* L. subsp. *oxycedrus* var. *spilinanus* Yalt., Eliçin & Terzioğlu: A new variety from Turkey. *Turk J Bot* 31: 37-40.
- Yıldırımlı Ş (2006). Two new species of *Phlomis* (Lamiaceae) from Turkey, *P. isilae* Yıld. and *P. dinci* Yıld. *Ot Sistematisk Botanik Dergisi* 13: 1-10.
- Yıldırımlı Ş & Doğru-Koca A (2006). A new species *Limonium adilguneri* Yıld. & Doğru-Koca (Plumbaginaceae) from edge of Tuzgölü, Inner Anatolia Turkey. *Ot Sistematisk Botanik Dergisi* 13: 11-16.
- Yıldırımlı Ş & Selvi B (2006). A new genus *Dumaniana* Yıldırımlı & Selvi and a new species *D. gelendostensis* Yıldırımlı & Selvi (Apiaceae) from Isparta, Southwest Anatolia, Turkey and new combinations. *Ot Sistematisk Botanik Dergisi* 13: 1-8.
- Yıldırımlı Ş, Sadıkoğlu N & Keskin M (2006). A new species from Turkey *Micromeria maritima* Yıldırımlı, Sadıkoğlu & Keskin (Lamiaceae). *Ot Sistematisk Botanik Dergisi* 13: 27-34.
- Yılmaz G & Dane F (2006). A new record for the flora of Turkey: *Verbascum samniticum* Ten. (V. *phlomoides* subsp. *sartorii* (Boiss. & Heldr.) Nyman) (Scrophulariaceae), IVth Balkan Botanical Congress, 20-26 June 2006, Abstract book, p. 205-206.
- Yılmaz Ö & Kaynak G (2006). *Linum hirsutum* subsp. *platyphyllum*, stat. nov. (Linaceae). *Ann Bot Fennici* 43: 62-63.
- Yılmaz Ö & Kaynak G (2006). New combination in *Linum* sect. *Syllinum* (Linaceae). *Ann Bot Fennici* 43: 77-79.
- Zielinski J, Tan K & Tomaszewski D (2004). Notes on the taxonomy of *Genista januensis* and *G. lydia* (Fabaceae). *Ann Bot Fennici* 41: 453-457.
- Zielinski J, Petrova A & Tomaszewski D (2006). *Quercus trojana* subsp. *yaltirikii* (Fagaceae) a new subspecies from southern Turkey. *Willdenowia* 36: 845-849.