

Preprint: January 31, 2010

## AN EXTENSION OF STERN'S CONGRUENCE

ZHI-HONG SUN AND LIN-LIN WANG

School of the Mathematical Sciences, Huaiyin Normal University,  
Huai'an, Jiangsu 223001, PR China  
Email: szh6174@yahoo.com; wanglinlin\_1986@yahoo.cn

**ABSTRACT.** Let  $\{E_n\}$  be the Euler numbers. In the paper we determine  $E_{2^m k + b} - E_b$  modulo  $2^{m+7}$ , where  $k, m$  are positive integers with  $m \geq 4$  and  $b \in \{0, 2, 4, \dots\}$ .

MSC: Primary 11B68, Secondary 11A07

Keywords: Euler number; Congruence

### 1. Introduction.

The Euler numbers  $\{E_n\}$  are given by

$$E_0 = 1, \quad E_{2n-1} = 0, \quad \sum_{r=0}^n \binom{2n}{2r} E_{2r} = 0 \quad (n = 1, 2, 3, \dots).$$

The first few Euler numbers are shown below:

$$\begin{aligned} E_0 &= 1, \quad E_2 = -1, \quad E_4 = 5, \quad E_6 = -61, \quad E_8 = 1385, \quad E_{10} = -50521, \\ E_{12} &= 2702765, \quad E_{14} = -199360981, \quad E_{16} = 19391512145. \end{aligned}$$

Let  $\mathbb{N}$  be the set of positive integers. For  $k, m \in \mathbb{N}$  and  $b \in \{0, 2, 4, \dots\}$ , in 1875 Stern ([2]) proved the following Stern's congruence:

$$(1.1) \quad E_{2^m k + b} \equiv E_b + 2^m k \pmod{2^{m+1}}.$$

There are many modern proofs of (1.1). See for example [1,3,5,6].

Let  $b \in \{0, 2, 4, \dots\}$  and  $k, m \in \mathbb{N}$ . In [4] the first author showed that

$$(1.2) \quad E_{2^m k + b} \equiv E_b + 2^m k \pmod{2^{m+2}} \quad \text{for } m \geq 2,$$

$$(1.3) \quad E_{2^m k + b} \equiv E_b + 5 \cdot 2^m k \pmod{2^{m+3}} \quad \text{for } m \geq 3$$

and

$$(1.4) \quad E_{2^m k + b} \equiv \begin{cases} E_b + 5 \cdot 2^m k \pmod{2^{m+4}} & \text{if } b \equiv 0, 6 \pmod{8}, \\ E_b - 3 \cdot 2^m k \pmod{2^{m+4}} & \text{if } b \equiv 2, 4 \pmod{8} \end{cases} \quad \text{for } m \geq 4.$$

---

The first author is supported by the Natural Sciences Foundation of China (grant No. 10971078).

From [4, Corollary 3.2] we know that for any nonnegative integer  $k$  and positive integer  $n$ ,

$$(1.5) \quad \frac{3^{2k+1} + 1}{4} E_{2k} \equiv \sum_{r=0}^{n-1} (-1)^{n-1-r} \binom{k-1-r}{n-1-r} \binom{k}{r} \frac{3^{2r+1} + 1}{4} E_{2r} \pmod{2^{3n}}.$$

In the paper we use (1.5) to obtain a congruence for  $E_{2^m k+b} - E_b$  modulo  $2^{m+7}$ , where  $k, m \in \mathbb{N}$ ,  $m \geq 4$  and  $b \in \{0, 2, 4, \dots\}$ , see Theorem 2.1. As a consequence, we have

$$(1.6) \quad E_{2^m k+b} \equiv E_b - 2^m k((b+1)^2 + 10) \pmod{2^{m+6}} \quad \text{for } m \geq 6.$$

Throughout the paper we use  $\mathbb{Z}_2$  to denote the set of rational numbers whose denominator is odd.

## 2. The main result.

**Theorem 2.1.** *Let  $b \in \{0, 2, 4, \dots\}$  and  $k, m \in \mathbb{N}$  with  $m \geq 4$ . Then*

$$E_{2^m k+b} \equiv E_b - 2^m k((b+1)^2 + 42 - 32(-1)^{\frac{b(b+2)}{8}} + 2^m k((-1)^{\frac{b}{2}} + 2(-1)^{\frac{b(b+2)}{8}})) \pmod{2^{m+7}}.$$

Proof. Set  $f(k) = \frac{3^{2k+b+1} + 1}{4} E_{2k+b}$  and  $F(k) = 2^{-3k} \sum_{r=0}^k \binom{k}{r} (-1)^r f(r)$ . From (1.5) (with  $n = k$ ) or [4, Lemma 2.3] we know that  $F(k) \in \mathbb{Z}_2$ . By the binomial inversion formula we have  $f(k) = \sum_{r=0}^k \binom{k}{r} (-2)^{3r} F(r)$  and so

$$\begin{aligned} f(2^{m-1}k) &= F(0) + \sum_{r=1}^{2^{m-1}k} \binom{2^{m-1}k}{r} (-2)^{3r} F(r) \\ &= F(0) + \sum_{r=1}^{2^{m-1}k} \frac{2^{m-1}k}{r} \binom{2^{m-1}k-1}{r-1} (-2)^{3r} F(r). \end{aligned}$$

For  $r \geq 3$  we have  $\frac{2^{r-2}}{r} \in \mathbb{Z}_2$  and  $F(r) \in \mathbb{Z}_2$ . Thus

$$\frac{2^{m-1} \cdot 2^{3r} F(r)}{r} = 2^{m+2r+1} \cdot \frac{2^{r-2}}{r} F(r) \equiv 0 \pmod{2^{m+7}}.$$

Hence,

$$\begin{aligned} (2.1) \quad &\frac{3^{2^m k+b+1} + 1}{4} E_{2^m k+b} \\ &= f(2^{m-1}k) \equiv F(0) - 2^{m+2}kF(1) + 2^{m+4}k(2^{m-1}k-1)F(2) \\ &= \frac{3^{b+1} + 1}{4} E_b - 2^{m-1}k(f(0) - f(1)) - 2^{m-2}k(f(0) - 2f(1) + f(2)) + 2^{2m+3}k^2 F(2) \\ &\equiv \frac{3^{b+1} + 1}{4} E_b - 2^{m-2}k(3f(0) - 4f(1) + f(2)) \pmod{2^{m+7}}. \end{aligned}$$

Putting  $n = 3$  in (1.5) we see that

$$\begin{aligned} \frac{3^{b+1} + 1}{4} E_b &\equiv \sum_{r=0}^2 (-1)^{2-r} \binom{b/2 - 1 - r}{2 - r} \binom{b/2}{r} \frac{3^{2r+1} + 1}{4} E_{2r} \\ &= \frac{\left(\frac{b}{2} - 1\right)\left(\frac{b}{2} - 2\right)}{2} - \frac{b}{2} \left(\frac{b}{2} - 2\right) \cdot 7 \cdot (-1) + \frac{\frac{b}{2}\left(\frac{b}{2} - 1\right)}{2} \cdot 61 \cdot 5 \\ &= 40b^2 - 84b + 1 \pmod{2^9}. \end{aligned}$$

Thus,

$$\begin{aligned} (2.2) \quad 3f(0) - 4f(1) + f(2) \\ &\equiv 3(40b^2 - 84b + 1) - 4(40(b+2)^2 - 84(b+2) + 1) + 40(b+4)^2 - 84(b+4) + 1 \\ &= 336 - 320b \equiv -176 - 64b \pmod{2^9}. \end{aligned}$$

For  $r \geq 3$  we see that  $4^{r-3}/r \in \mathbb{Z}_2$  and so

$$\binom{2^{m-2}k}{r} 80^r = 2^{m+4+2r} \cdot \frac{4^{r-3}}{r} \cdot k \cdot 5^r \binom{2^{m-2}k - 1}{r-1} \equiv 0 \pmod{2^{m+9}}.$$

Thus,

$$\begin{aligned} 3^{2^m k} &= (1 + 80)^{2^{m-2}k} = \sum_{r=0}^{2^{m-2}k} \binom{2^{m-2}k}{r} 80^r \\ &\equiv 1 + 5k \cdot 2^{m+2} + 25k \cdot 2^{m+5}(2^{m-2}k - 1) \\ &= 1 - 195 \cdot 2^{m+2}k \equiv 1 + 61 \cdot 2^{m+2}k \pmod{2^{m+9}}. \end{aligned}$$

This together with (2.1) and (2.2) yields

$$\begin{aligned} &\frac{3^{b+1}(1 + 61 \cdot 2^{m+2}k) + 1}{4} E_{2^m k + b} \\ &\equiv \frac{3^{2^m k + b + 1} + 1}{4} E_{2^m k + b} \equiv \frac{3^{b+1} + 1}{4} E_b + 2^{m-2}k(176 + 64b) \pmod{2^{m+7}}. \end{aligned}$$

Thus,

$$(2.3) \quad \frac{3^{b+1} + 1}{4} (E_{2^m k + b} - E_b) \equiv -55 \cdot 3^b \cdot 2^m k E_{2^m k + b} + 2^m k (44 + 16b) \pmod{2^{m+7}}.$$

By [3, Corollary 7.9] or [4, Corollary 3.5] we have

$$(2.4) \quad E_b \equiv \begin{cases} 3b^2 - 11b + 1 \pmod{2^7} & \text{if } 4 \mid b, \\ b^2 - 23b + 41 \pmod{2^7} & \text{if } 4 \mid b - 2. \end{cases}$$

Thus,

$$(2.5) \quad E_{2^m k+b} \equiv \begin{cases} 3b^2 - 11b + 1 + 2^m k(6b - 11) \pmod{2^7} & \text{if } 4 \mid b, \\ b^2 - 23b + 41 + 2^m k(2b - 23) \pmod{2^7} & \text{if } 4 \mid b - 2. \end{cases}$$

As

$$\begin{aligned} 3^b &= (1+8)^{b/2} = 1 + \binom{b/2}{1} 8 + \binom{b/2}{2} 8^2 + \dots \equiv 8b^2 - 12b + 1 \\ &\equiv \begin{cases} -12b + 1 \pmod{2^7} & \text{if } 4 \mid b, \\ 20b - 31 \pmod{2^7} & \text{if } 4 \mid b - 2, \end{cases} \end{aligned}$$

by (2.5) we have

$$3^b E_{2^m k+b} \equiv \begin{cases} (3b^2 - 11b + 1 + 2^m k(6b - 11))(-12b + 1) \pmod{2^7} & \text{if } 4 \mid b, \\ (b^2 - 23b + 41 + 2^m k(2b - 23))(20b - 31) \pmod{2^7} & \text{if } 4 \mid b - 2. \end{cases}$$

If  $4 \mid b$ , then

$$(3b^2 - 11b + 1)(-12b + 1) = -36b^3 + 135b^2 - 23b + 1 \equiv 7b^2 - 23b + 1 \pmod{128}$$

and

$$(6b - 11)(-12b + 1) \equiv -11 \equiv -3 \pmod{8}.$$

If  $4 \mid b - 2$ , then  $32 \mid (b-2)(b+2)$  and so

$$\begin{aligned} (b^2 - 23b + 41)(20b - 31) &= ((b-2)^2 - 19(b-2) - 1)(20(b+2) - 71) \\ &\equiv -20(b+2) - 71(b^2 - 23b + 41) \\ &\equiv -7b^2 + 13b - 7 \pmod{128} \end{aligned}$$

and

$$(2b - 23)(20b - 31) = (2(b-2) - 19)(20b - 31) \equiv 19 \cdot 31 \equiv -3 \pmod{8}.$$

Thus

$$3^b E_{2^m k+b} \equiv \begin{cases} 7b^2 - 23b + 1 - 3 \cdot 2^m k \pmod{2^7} & \text{if } 4 \mid b, \\ -7b^2 + 13b - 7 - 3 \cdot 2^m k \pmod{2^7} & \text{if } 4 \mid b - 2. \end{cases}$$

Substituting this into (2.3) we obtain

$$(2.6) \quad \frac{3^{b+1} + 1}{4} (E_{2^m k+b} - E_b) \equiv \begin{cases} 2^m k(-b^2 + b - 11 - 3 \cdot 2^m k) \pmod{2^{m+7}} & \text{if } 4 \mid b, \\ 2^m k(b^2 + 5b + 45 - 3 \cdot 2^m k) \pmod{2^{m+7}} & \text{if } 4 \mid b - 2. \end{cases}$$

It is easily seen that

$$\frac{3^{b+1} + 1}{4} = \frac{3(1+8)^{b/2} + 1}{4} \equiv \frac{3\left(1 + \frac{b}{2} \cdot 8 + \binom{b/2}{2} \cdot 8^2\right) + 1}{4} = 6b^2 - 9b + 1 \pmod{2^7}.$$

If  $4 \mid b$ , one can easily check that

$$(6b^2 - 9b + 1)(b^3 + 3b^2 + 9b + 1) \equiv 1 \pmod{2^7}, \quad b^3 + 3b^2 + 9b + 1 \equiv 1 + b \pmod{8}$$

and

$$(-b^2 + b - 11)(b^3 + 3b^2 + 9b + 1) \equiv b^3 - b^2 - 2b - 11 \pmod{2^7}.$$

Thus, by (2.6) we have

$$\begin{aligned} E_{2^m k+b} - E_b &\equiv 2^m k(-b^2 + b - 11 - 3 \cdot 2^m k)(b^3 + 3b^2 + 9b + 1) \\ &\equiv 2^m k(b^3 - b^2 - 2b - 11 - 3(b+1) \cdot 2^m k) \\ &\equiv \begin{cases} 2^m k(-(b+1)^2 - 10 - 3 \cdot 2^m k) \pmod{2^7} & \text{if } 8 \mid b, \\ 2^m k(-(b+1)^2 - 10 + 64 + 2^m k) \pmod{2^7} & \text{if } 8 \mid b - 4. \end{cases} \end{aligned}$$

If  $4 \mid b - 2$ , one can easily check that

$$\begin{aligned} (6b^2 - 9b + 1)((b+2)^3 - 3(b+2)^2 + 9(b+2) + 3) &\equiv 1 \pmod{2^7}, \\ (b+2)^3 - 3(b+2)^2 + 9(b+2) + 3 &\equiv b+5 \pmod{8} \end{aligned}$$

and

$$(b^2 + 5b + 45)((b+2)^3 - 3(b+2)^2 + 9(b+2) + 3) \equiv b^3 + 5b^2 - 22b + 61 \pmod{2^7}.$$

Thus, by (2.6) we have

$$\begin{aligned} E_{2^m k+b} - E_b &\equiv 2^m k(b^2 + 5b + 45 - 3 \cdot 2^m k)((b+2)^3 - 3(b+2)^2 + 9(b+2) + 3) \\ &\equiv 2^m k(b^3 + 5b^2 - 22b + 61 - 3(b+5) \cdot 2^m k) \pmod{2^7}. \end{aligned}$$

Since

$$\begin{aligned} b^3 + 5b^2 - 22b + 61 + (b+1)^2 + 10 \\ = b^3 + 6b^2 - 20b + 72 = (b+2)^3 - 32(b-2) \\ \equiv (b+2)^3 \equiv \begin{cases} 0 \pmod{128} & \text{if } b \equiv 6 \pmod{8}, \\ 64 \pmod{128} & \text{if } b \equiv 2 \pmod{8}, \end{cases} \end{aligned}$$

we see that

$$\begin{aligned} E_{2^m k+b} - E_b &\equiv 2^m k(b^3 + 5b^2 - 22b + 61 - 3(b+5) \cdot 2^m k) \\ &\equiv \begin{cases} 2^m k(-(b+1)^2 - 10 - 2^m k) \pmod{2^{m+7}} & \text{if } b \equiv 6 \pmod{8}, \\ 2^m k(-(b+1)^2 - 10 + 64 + 3 \cdot 2^m k) \pmod{2^{m+7}} & \text{if } b \equiv 2 \pmod{8}. \end{cases} \end{aligned}$$

Note that  $(-1)^{\frac{b(b+2)}{8}} = 1$  or  $-1$  according as  $b \equiv 0, 6 \pmod{8}$  or  $b \equiv 2, 4 \pmod{8}$ . Combining all of the above we deduce the result.

**Remark 2.1** Let  $k \in \{0, 1, 2, \dots\}$  and  $b \in \{0, 2, 4, \dots\}$ . Taking  $m = 3$  in [3, Corollary 7.8] we deduce

$$E_{8k+b} \equiv \binom{k}{2} E_{16+b} - k(k-2)E_{8+b} + \binom{k-1}{2} E_b \pmod{1024}.$$

Hence

$$(2.7) \quad E_{8k+b} - E_b \equiv \binom{k}{2} (E_{16+b} - E_b) - k(k-2)(E_{8+b} - E_b) \pmod{1024}.$$

By [4, Corollary 3.7] we have

$$(2.8) \quad E_{4k} \equiv \frac{1+176k+128k^2}{1+220k-160k^2} \pmod{512} \text{ and } E_{4k+2} \equiv \frac{-7-208k+128k^2}{7-68k+96k^2} \pmod{512}.$$

**Corollary 2.1.** Let  $k, m \in \mathbb{N}$  with  $m \geq 4$ . Then

$$E_{2^m k} \equiv 1 - 2^m k(11 + 3 \cdot 2^m k) \pmod{2^{m+7}}$$

and

$$E_{2^m k+2} \equiv -1 + 2^m k(45 + 3 \cdot 2^m k) \pmod{2^{m+7}}.$$

**Corollary 2.2.** Let  $b \in \{0, 2, 4, \dots\}$  and  $k, m \in \mathbb{N}$  with  $m \geq 4$ . Then

$$E_{2^m k+b} \equiv E_b - 2^m k((b+1)^2 + 10 - (-1)^{\frac{b}{2}} 2^m k) \pmod{2^{m+6}}.$$

**Corollary 2.3.** Let  $b \in \{0, 2, 4, \dots\}$  and  $k, m \in \mathbb{N}$  with  $m \geq 5$ . Then

$$E_{2^m k+b} \equiv \begin{cases} E_b - 11 \cdot 2^m k \pmod{2^{m+5}} & \text{if } b \equiv 0, 14 \pmod{16}, \\ E_b + 13 \cdot 2^m k \pmod{2^{m+5}} & \text{if } b \equiv 2, 12 \pmod{16}, \\ E_b - 3 \cdot 2^m k \pmod{2^{m+5}} & \text{if } b \equiv 4, 10 \pmod{16}, \\ E_b + 5 \cdot 2^m k \pmod{2^{m+5}} & \text{if } b \equiv 6, 8 \pmod{16}. \end{cases}$$

Proof. By Corollary 2.2 we have

$$E_{2^m k+b} \equiv E_b - 2^m k((b+1)^2 + 10) \pmod{2^{m+5}}.$$

Thus the result follows.

In the end we pose the following conjecture.

**Conjecture 2.1.** Let  $m, n \in \mathbb{N}$ ,  $m \geq n$  and  $b \in \{0, 2, 4, \dots, 2^{m+n-1} - 2\}$ . Then

$$E_{2^m k+b} - E_b \equiv E_{2^m k+2^{m+n-1}-2-b} - E_{2^{m+n-1}-2-b} \pmod{2^{m+n}}.$$

## REFERENCES

1. R. Ernvall, *Generalized Bernoulli numbers, generalized irregular primes, and class number*, Ann. Univ. Turku. (Ser. A), **178** (1979), 1-72.
2. M.A. Stern, *Zur Theorie der Eulerischen Zahlen*, J. Reine Angew. Math. **79** (1875), 67-98.
3. Z.H. Sun, *Congruences involving Bernoulli polynomials*, Discrete Math. **308** (2008), 71-112.
4. Z.H. Sun, *Euler numbers modulo  $2^n$* , Bull. Aust. Math. Soc. **82** (2010), 221-231.
5. Z.W. Sun, *On Euler numbers modulo powers of two*, J. Number Theory **115** (2005), 371-380.
6. S.S. Wagstaff Jr., *Prime divisors of the Bernoulli and Euler numbers*, in: M.A. Bennett et al. (Eds.), *Number Theory for the Millennium*, vol. III (Urbana, IL, 2000), A K Peters, Natick, MA, 2002, 357-374.