

POSITIVE LYAPUNOV EXPONENT BY A RANDOM PERTURBATION

ZENG LIAN AND MIKKO STENLUND

ABSTRACT. We study the effect of a random perturbation on a one-parameter family of dynamical systems whose behavior in the absence of perturbation is ill understood. We provide conditions under which the perturbed system is ergodic and admits a positive Lyapunov exponent, with an explicit lower bound, for a large and controlled set of parameter values.

Acknowledgements. The authors are indebted to Lai-Sang Young for stimulating discussions. Mikko Stenlund has received funding from the Academy of Finland.

1. INTRODUCTION

1.1. **Background.** The question of the existence of positive Lyapunov exponents for a given dynamical system is one of extreme importance. It turns out to be very hard even in seemingly simple examples, which poses a great challenge to modern mathematics.

For example, consider the logistic family ¹

$$P_a : x \mapsto ax(1 - x), \quad a \in [1, 4].$$

Let the set \mathcal{A} consist of those values of $a \in [1, 4]$ for which P_a admits a unique, finite, ergodic, absolutely continuous invariant measure, with a positive Lyapunov exponent. On the other hand, let \mathcal{B} denote the set of $a \in [1, 4]$ for which P_a has a periodic orbit attracting all orbits in $[0, 1]$. The set \mathcal{A} is known to have positive measure [7] (also [2, 20]) and the set \mathcal{B} is open and dense [6, 16]. Using renormalization arguments, it has moreover been shown in [17] (see also [15]) that almost every value of $a \in [1, 4]$ falls in precisely one of the two sets \mathcal{A} and \mathcal{B} .

The above results on the abundance of parameter values admitting either a positive Lyapunov exponent or a periodic sink have been extended to multimodal situations in which several critical points are allowed; see [22, 25] and [11], respectively. However, it is not known whether the union of the two classes forms a set of full measure.

Taking these considerations into account, we understand why even such one-dimensional systems are notoriously hard to analyze: the nature of the dynamics depends very sensitively on the value of the parameter. Adding random noise to the model simplifies the picture due to averaging effects. For noisy systems, the dependence of a Lyapunov exponent on parameters is regular under mild conditions. Second, it is a well-known

Date: December 21, 2010.

2000 Mathematics Subject Classification. 37H15; 70K60.

Key words and phrases. Lyapunov exponent, random perturbation.

¹Alternatively, as is done in some of the cited references, one could consider the real quadratic family $Q_a : x \mapsto a + x^2$, $a \in [-2, \frac{1}{4}]$.

dichotomy in the random case (see, *e.g.*, [27] for a discussion) that the sign of the Lyapunov exponent indicates in which dynamical category the system belongs to: a negative Lyapunov exponent implies convergence to a random sink consisting of finitely many points for almost all sample paths [1, 13], while a positive one yields a random SRB measure almost surely [12].

Regarding random perturbations of dynamical systems, it is commonplace to start out with systems that are very well controlled in the absence of perturbation. One then goes on to show that control of the system is retained under sufficiently small random perturbations. A system possessing this property is called stochastically stable. For uniformly hyperbolic systems, standard references on stochastic stability include [10] and [26]. For one-dimensional maps admitting critical points, see [8] and [4]. The more recent [3] discusses a two-dimensional case. The preceding list, which of course could be continued much further, is meant to point the reader quickly to a handful of original references.

While results of the above kind are very interesting, it would be much more satisfying if one could reverse the direction. That is, to perturb a dynamical system too hard to analyze by itself, to take advantage of the randomness in the noisy system, and then to infer properties of the zero-noise limit. The idea of doing so can be traced back at least to Pontryagin, Andronov, and Vitt [19], and later to Kolmogorov [21]. As the real world is inherently noisy, say, an invariant measure obtained in that limit could be interpreted as a physically observable (albeit idealized) statistical description of the system. Unfortunately, the zero-noise limit is not always well behaved. For instance, Lyapunov exponents may fail to be continuous at the point of vanishing perturbation; see the figure-eight attractor in [5].

To take steps in the direction of the previous paragraph — and more generally to develop new techniques for proving lower bounds on Lyapunov exponents — we work with a one-parameter family of systems in which the dynamical properties of the unperturbed system for a given parameter value are unknown. Here, a sufficiently large perturbation is required (i) to regularize the parameter dependence of the nature of the dynamics so that (ii) a good lower bound on the Lyapunov exponent can be established for a large and controlled set of parameters.

The paper is organized as follows. In Section 1.2 we introduce our model and the necessary technical notions so that the results of the paper can be formulated in Section 1.3. Theorem 1 concerns ergodicity of the system and is proved in Section 2. Proposition 4 identifies parameter values for which random sinks appear unless the perturbation is large enough. Its proof is given in Section 3. Theorems 2 and 5 give sufficient conditions for a positive Lyapunov exponent together with an explicit lower bound. They are also proved in Section 3.

1.2. Preliminaries. We denote by \mathbb{S} the circle obtained by identifying the endpoints of the unit interval $[0, 1]$ and by m the uniform measure on \mathbb{S} . Let

$$\tau_a : \mathbb{S} \rightarrow \mathbb{S} : x \mapsto a + x + L\psi(x) \pmod{1},$$

where $a \in [0, 1)$ and $L > 0$ are constants, and $\psi : \mathbb{S} \rightarrow \mathbb{R}$ is a twice continuously differentiable map. Although τ_a depends on L , it is notationally convenient not to indicate this explicitly by a subscript. We assume that ψ has $N > 0$ critical points c_1, \dots, c_N where $\psi'(c_i) = 0$, each of which is nondegenerate, *i.e.*, $\psi''(c_i) \neq 0$. Since nondegenerate critical points are isolated and the circle is compact, $N < \infty$.

Notice that the maps τ_a are not unimodal with just one critical point, which is a case studied extensively in the literature. By contrast, the present paper involves a rather general class of multimodal maps for which the number of critical points is arbitrary. This setting arises, for example, in applications pertaining to shear-induced chaos in the theory of rank one attractors [14, 18, 23–25].

As discussed earlier, the parametric dependence of the dynamical nature of such maps can be very complicated. With the exception of some special parameter values, it is practically impossible to determine whether a particular choice of the parameter a results in chaotic or regular motion. To remedy the situation, we add a small amount of random noise to the system, which turns out to have a regularizing effect on the dependence of the dynamics on a .

To be specific, we are interested in the ergodic and chaotic properties of the random circle map $\tau_a + Y$ determined by

$$x \mapsto \tau_a(x) + Y \pmod{1},$$

where Y is a random perturbation, or kick, distributed uniformly on $[-\varepsilon, \varepsilon]$ with some $\varepsilon > 0$. The reader may think of ε as the level of noise present in the system. For simplicity, we take $Y(\omega) = \omega$ for each realization $\omega \in [-\varepsilon, \varepsilon]$. Given a realized sequence $(\omega_n)_{n=1}^\infty \in [-\varepsilon, \varepsilon]^{\mathbb{Z}^+}$ of i.i.d. kicks, the trajectory $(x_n)_{n=0}^\infty$ of any initial point $x_0 = x \in \mathbb{S}$ is determined for each $n \geq 1$ by

$$x_n = \tau_a(x_{n-1}) + \omega_n.$$

Let us write

$$\tau_a^\omega(x) = \tau_a(x) + \omega \quad \forall \omega \in \mathbb{R}$$

and denote the uniform probability measure on $[-\varepsilon, \varepsilon]$ by η . We say that a Borel probability measure μ on \mathbb{S} is *invariant* for the above random map $\tau_a + Y$, if

$$\mu(B) = \int_{[-\varepsilon, \varepsilon]} \mu((\tau_a^\omega)^{-1}B) d\eta(\omega)$$

holds for every Borel set $B \subset \mathbb{S}$. Further, an invariant measure μ is *ergodic*, if the condition $\mu(B \Delta (\tau_a^\omega)^{-1}B) = 0$ for η -a.e. $\omega \in [-\varepsilon, \varepsilon]$ on the set B implies $\mu(B) \in \{0, 1\}$. Here $A \Delta B$ denotes the symmetric difference $(A \setminus B) \cup (B \setminus A)$ of two sets. For example, it is easy to check that for complete smearing ($\varepsilon = \frac{1}{2}$) of the image $\tau_a(x)$, the unique invariant measure — which is always ergodic — is $\mu = m$. In general, there can exist many invariant measures. However, we will see in Lemma 6 that there can be at most one measure which is both ergodic and equivalent to m . For a sufficiently noisy system, such a measure turns out to exist and to rule out the existence of other invariant measures.

It follows from Birkhoff's ergodic theorem that the *Lyapunov exponent*

$$\lambda_a((\omega_n)_{n=1}^\infty, x; L) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \log |\tau'_a(x_k)| \quad (1)$$

exists $(\eta^{\mathbb{Z}^+} \times \mu)$ -a.s., if μ is invariant for the random map $\tau_a + Y$. If μ is also ergodic, then $\lambda_a((\omega_n)_{n=1}^\infty, x; L)$ is $(\eta^{\mathbb{Z}^+} \times \mu)$ -a.s. equal to the constant

$$\lambda_a(L) = \int_{\mathbb{S}} \log |\tau'_a| d\mu. \quad (2)$$

Notice that the Lyapunov exponent measures the exponential rate of separation of initial points infinitesimally close to x , under the same sequence $(\omega_n)_{n=1}^\infty$ of kicks. A positive Lyapunov exponent indicates sensitive dependence of the trajectory on the initial condition.

For any $K > 1$, the map τ_a is uniformly expanding on the complement of the set

$$I_K = \{x \in \mathbb{S} : |\tau'_a(x)| \leq K\}.$$

The non-expanding set I_1 is critical to the dynamics and its structure plays a central role in our analysis.

For any $K \geq 1$, the set I_K is independent of a and consists of those points $x \in \mathbb{S}$ for which $-(K+1)/L \leq \psi'(x) \leq (K-1)/L$. For a sufficiently large L , they form N disjoint intervals, each containing precisely one of the critical points c_i of ψ , which does not depend on L . (For smaller values of L , some of the intervals merge.) The endpoints are obtained by solving $\psi'(c_i + \xi_i) = \mp(K \pm 1)/L$ for ξ_i . Taylor expanding, we see that the length of such an interval is $2K/|\psi''(c_i)|L + O((K/L)^2)$. In particular, the length of the largest component of I_K is

$$b_K = \frac{2K}{L} \frac{1}{\min_{1 \leq i \leq N} |\psi''(c_i)|} + O((K/L)^2) \quad \text{as } L/K \rightarrow \infty, \quad (3)$$

and

$$m(I_K) = \frac{2K}{L} \sum_{1 \leq i \leq N} \frac{1}{|\psi''(c_i)|} + O((K/L)^2) \quad \text{as } L/K \rightarrow \infty. \quad (4)$$

Throughout this paper,

$$B_r(A) = \{x \in \mathbb{S} : \text{dist}(x, A) \leq r\}$$

is the closed r -neighborhood of a set $A \subset \mathbb{S}$ and $B_r(x) = B_r(\{x\})$. In our estimates, C stands for a generic constant whose numerical value may change from one expression to the next.

1.3. Results. We are now in position to state the results of this paper.

Theorem 1. *Given any $L > 0$, $a \in [0, 1)$, and $\varepsilon > m(I_{N+1})/2$, the system admits a unique invariant measure. It is both ergodic and equivalent to m . For any sufficiently large value of L , the same is true for any $a \in [0, 1)$ and $\varepsilon > b_2(L)/2$.*

Theorem 2. *Given constants $C > 0$ and $\beta \in (0, 1]$, a function $\varepsilon = \varepsilon(L) \geq CL^{\beta-1}$, and a sufficiently large L , the system admits a unique ergodic measure for any value of a . With respect to those measures,*

$$\liminf_{L \rightarrow \infty} \inf_{a \in (0,1)} \frac{\lambda_a(L)}{\log L} \geq \beta.$$

Remark 3. *The assumption on the size of the perturbation ε guarantees ergodicity. Indeed, recalling (3), the condition $\varepsilon > b_2(L)/2$ of Theorem 1 is satisfied if L is large.*

Let us pause to discuss the cases $\beta = 1$ and $\beta = 0$. Assuming that the measure μ is ergodic, with no explicit constraints on the parameters, Jensen's inequality implies, for all $L \geq 1$,

$$\exp \lambda_a(L) = \exp \int_{\mathbb{S}} \log |\tau'_a| d\mu \leq \int_{\mathbb{S}} 1 + L|\psi'| d\mu \leq CL,$$

or

$$\limsup_{L \rightarrow \infty} \sup_{a \in (0,1)} \frac{\lambda_a(L)}{\log L} \leq 1. \quad (5)$$

Hence, the lower bound of Theorem 1 for $\beta = 1$ is optimal.

When $\beta = 0$, the theorem above suggests that all values of a may not yield a positive limit for $\lambda_a(L)/\log L$. The reason is that the size of the non-expanding set I_1 scales like L^{-1} , as observed above. If the perturbation is not sufficiently large in comparison, the smeared image $B_\varepsilon(\tau_a(z))$ of a critical point z of τ_a may, for some values of the parameter a , be contained in a small contracting neighborhood of z . By this mechanism, negative Lyapunov exponents appear:

Proposition 4. *There exists a constant $D > 0$ and, for any large enough L and for any $\varepsilon \leq DL^{-1}$, an invariant measure μ , such that*

$$\lambda_a((\omega_n)_{n=1}^\infty, x; L) < 0$$

for $(\eta^{\mathbb{Z}^+} \times \mu)$ -a.e. $((\omega_n)_{n=1}^\infty, x)$.

We point out that, by general results [1, 13], a negative Lyapunov exponent for an ergodic random system implies the existence of a random attracting set consisting of finitely many points. Such a set can in fact be constructed following the proof of the above proposition.

To shed more light on the case $\beta = 0$ in particular, we have the following result, which holds for a *restricted set of values* of the parameter a .

Theorem 5. *There exist sets $A_L \subset [0, 1)$, $L > 0$, such that the following holds. Each A_L consists of a union of at most N^2 intervals and*

$$\lim_{L \rightarrow \infty} m(A_L) = 1.$$

Given a function $\varepsilon = \varepsilon(L) > b_2(L)/2$ the system admits a unique ergodic measure for any sufficiently large value of L and any value of a . With respect to those measures,

$$\liminf_{L \rightarrow \infty} \inf_{a \in A_L} \frac{\lambda_a(L)}{\log L} \geq \frac{1}{2}.$$

The number $\frac{1}{2}$ appearing in Theorem 5 is a technical artifact of the proof. Namely, the set A_L consists roughly speaking of parameter values for which the set I_1 does not intersect its image $B_\varepsilon(\tau_a(I_1))$ under the random map $\tau_a + Y$. Clearly one could exclude fewer parameter values by considering cases in which the trajectory of a point is allowed to visit the set I_1 at several consecutive times. Moreover, the estimate could be improved for many parameter values by more elaborate techniques. In view of the fact that the value 1 could not be exceeded due to the upper bound in (5), we have not pursued such an improvement.

2. ERGODICITY

We will next prove Theorem 1. But first we need to recall some basic facts and definitions.

Notice that we can view the random circle map $\tau_a + Y$ above as the Markov chain generated by the transition kernel

$$p(x, A) = \frac{1}{2\varepsilon} \mathfrak{m}(A \cap B_\varepsilon(\tau_a(x))), \quad (6)$$

for points $x \in \mathbb{S}$ and Borel sets $A \subset \mathbb{S}$. In other words, $p(x, A)$ is the probability that the random image of the initial point x belongs to the set A . A Borel measure μ on \mathbb{S} is stationary if

$$\mu(A) = \int_{\mathbb{S}} p(x, A) d\mu(x). \quad (7)$$

Stationary measures for the Markov chain are precisely the invariant measures for the random map. Because $x \mapsto p(x, A)$ is a continuous function for any A , it is a standard fact [9] that there exists a stationary measure μ . For any $\varepsilon > 0$,

$$\mu(A) \leq \max_x p(x, A) \leq \frac{1}{2\varepsilon} \mathfrak{m}(A) \quad (8)$$

follows immediately from (7). In other words, μ is absolutely continuous with respect to the measure \mathfrak{m} , written $\mu \ll \mathfrak{m}$, and therefore has a density ρ :

$$\mu(A) = \int_A \rho d\mathfrak{m}.$$

Define $P^* : L^1(\mathbb{S}) \rightarrow L^1(\mathbb{S})$ by

$$(P^*f)(x) = \int_{\mathbb{S}} f(y) p(x, dy), \quad f \in L^1(\mathbb{S}).$$

Thus, given an initial state $x \in \mathbb{S}$ of the system, the expected value of a function $f \in L^1(\mathbb{S})$ after one time step is $(P^*f)(x)$. We say that a Borel set A is invariant modulo μ if $(P^*1_A)(x) = 1_A(x)$ for μ -a.e. x . Notice that in this case $p(x, A) = 1$ for μ -a.e. $x \in A$. Finally, the measure μ is *ergodic*, if all invariant sets are trivial, i.e., $\mu(A) \in \{0, 1\}$ whenever A is invariant modulo μ . Again, this definition of ergodicity coincides with the one given earlier for the random map.

Lemma 6. *Let $\varepsilon > 0$ and consider the Markov chain generated by the transition kernel p . If there exists an ergodic stationary measure which is equivalent to \mathfrak{m} , there are no other stationary measures.*

Proof. Let μ_1 and μ_2 be ergodic stationary measures for the Markov chain on the state space \mathbb{S} . Recall that, given an initial measure μ , the chain generates a probability measure P^μ on the space of trajectories, $\mathbb{S}^{\mathbb{N}}$, and

$$P^\mu = \int_{\mathbb{S}} P^x d\mu(x), \quad (9)$$

where P^x is the measure corresponding to an initial point mass at x . In the sense of measure preserving transformations, P^{μ_1} and P^{μ_2} are ergodic with respect to the left shift on $\mathbb{S}^{\mathbb{N}}$. It follows from Birkhoff's ergodic theorem that either $P^{\mu_1} = P^{\mu_2}$ or the measures are mutually singular, written $P^{\mu_1} \perp P^{\mu_2}$. In the first case, $\mu_1 = \mu_2$, as can be seen by considering sets of the form $A \times \mathbb{S}^{\mathbb{Z}_+} \subset \mathbb{S}^{\mathbb{N}}$ with $A \subset \mathbb{S}$ a Borel set. In the second case, there exists a Borel set $\mathcal{A} \subset \mathbb{S}^{\mathbb{N}}$ such that $P^{\mu_1}(\mathcal{A}) = 1$ and $P^{\mu_2}(\mathcal{A}) = 0$. Therefore (9) implies $P^x(\mathcal{A}) = 1$ for μ_1 -a.e. x and $P^x(\mathcal{A}) = 0$ for μ_2 -a.e. x , meaning that $\mu_1 \perp \mu_2$.

In conclusion, two distinct ergodic measures are mutually singular. Assuming now that there exists an ergodic measure μ which is equivalent to m , it must be the only ergodic measure, because by (8) any other candidate would also have a density with respect to m . Since any stationary measure is a convex combination of ergodic ones, μ must in fact be the only stationary measure for the Markov chain. \square

Proof of Theorem 1. Uniqueness of the measure μ with the claimed properties is guaranteed by Lemma 6. Thus, we are left with proving existence.

First, we claim that for m -a.e. $x \in \text{supp } \mu$, it holds true that $p(x, \text{supp } \mu) = 1$. Since $d\mu = \rho dm$ is stationary

$$\mu(\text{supp } \mu) = \int_{\mathbb{S}} p(x, \text{supp } \mu) d\mu(x) = \int_{\text{supp } \mu} p(x, \text{supp } \mu) \rho(x) dm(x).$$

The claim follows from $\mu(\text{supp } \mu) = \int_{\text{supp } \mu} \rho(x) dm(x)$. Moreover, since $\mu \ll m$, we also have $m(\text{supp } \mu) > 0$.

Second, we show that any Borel set A invariant modulo μ with $\mu(A) > 0$ has $m(A) = 1$. By $\mu \ll m$, this also implies $\mu(A) = 1$. Since $\text{supp } \mu$ is invariant modulo μ , we assume without loss of generality that $A \subset \text{supp } \mu$. The idea of the proof is to construct a sequence of intervals J_0, J_1, \dots with $J_{i+1} = B_\varepsilon(\tau_a(J_i))$ such that **(i)** $J_i \subset A \pmod{m}$, meaning $m(J_i \setminus A) = 0$ and **(ii)** $J_i = \mathbb{S}$ for all sufficiently large values of i , provided ε is sufficiently large. As a byproduct, we will have obtained $m(\text{supp } \mu) = 1$, which shows that m and μ are equivalent measures.

Proof of (i). Note that restricted to $\text{supp } \mu \supset A$, the statements “ μ -a.e.” and “ m -a.e.” are equivalent, so we will simply write “a.e.” in such a situation. We fix an arbitrary parameter value $a \in [0, 1)$ for the map τ_a . For a.e. $x \in A$, $p(x, A) = 1$. We pick such an x . Then, by (6), the interval $J_0 = B_\varepsilon(\tau_a(x))$ satisfies $J_0 \subset A \pmod{m}$. Observe that $\mu(J_0 \cap A) > 0$ because $m(J_0 \cap A) = 2\varepsilon$ and $A \subset \text{supp } \mu$. By invariance of A , $p(y, A) = 1$ a.e. $y \in J_0 \cap A$. Denote the set of such y by \tilde{J}_0 . Then $J_0 = \tilde{J}_0 \cup N_0$ for some m -null set N_0 .

We define $J_i = B_\varepsilon(\tau_a(J_{i-1}))$ and $\tilde{J}_i = \{y \in J_i \cap A : p(y, A) = 1\} \subset A$ for $i \geq 1$ inductively. We claim that $J_i = \tilde{J}_i \cup N_i$ for some m -null set N_i . The proof is inductive.

First of all, denoting by ∂J_i the boundary of J_i (consisting of no more than two points), we have

$$J_i = \bigcup_{y \in \hat{J}_{i-1}} B_\varepsilon(\tau_a(y)) \cup \partial J_i,$$

where \hat{J}_{i-1} is a countable dense subset of \tilde{J}_{i-1} . This is so, because J_i and $J_{i-1} = \tilde{J}_{i-1} \cup N_{i-1}$ are closed intervals and $m(N_{i-1}) = 0$. Second, for each $y \in \tilde{J}_{i-1}$ we have $p(y, A) = 1$, which implies $B_\varepsilon(\tau_a(y)) \subset A \pmod{m}$. Since \hat{J}_{i-1} is countable, we conclude $J_i \subset A \pmod{m}$. Hence also $\tilde{J}_i = J_i \pmod{m}$ by invariance of A .

Proof of (ii). Recall that N is the number of critical points of the map ψ and that $I_K = \{x \in \mathbb{S} : |\tau'_a(x)| \leq K\}$. Suppose first that $L > 0$ is arbitrary. For any $K > N + 1$, any $\varepsilon \geq Km(I_K)/2(N + 1)$, and any interval $J \subset \mathbb{S}$, we have that

$$\begin{aligned} m(B_\varepsilon(\tau_a(J))) &\geq \min\{1, 2\varepsilon + m(\tau_a(J \cap (I_K)^c))\} \\ &\geq \min\left\{1, 2\varepsilon + \frac{K}{N+1}m(J \cap (I_K)^c)\right\} \\ &\geq \min\left\{1, 2\varepsilon + \frac{K}{N+1}(m(J) - m(I_K))\right\} \\ &\geq \min\left\{1, \frac{K}{N+1}m(J)\right\}. \end{aligned}$$

The maximum number N of critical points enters the argument, because although the map τ_a is locally expanding on $(I_K)^c$, the graph of τ_a has a fold at each critical point. The above estimate shows that the interval J_i grows exponentially with i until it covers \mathbb{S} .

Now, assume instead that $L > 0$ is so large that τ_a wraps each of the intervals (z_i, z_{i+1}) around \mathbb{S} twice, where z_i are the critical points of τ_a labeled clockwise, and that each z_i is nondegenerate. Let the interval $J \subset \mathbb{S}$ be such that $B_\varepsilon(\tau_a(J)) \neq \mathbb{S}$ — otherwise we are done. Then J contains at most one of the critical points z_i and intersects at most one component of I_K . Recall that b_K denotes the length of the largest component of I_K . For any $K > 2$ and any $\varepsilon \geq Kb_K/4$,

$$\begin{aligned} m(B_\varepsilon(\tau_a(J))) &\geq 2\varepsilon + m(\tau_a(J \cap (I_K)^c)) \\ &\geq 2\varepsilon + \frac{K}{2}m(J \cap (I_K)^c) \\ &\geq 2\varepsilon + \frac{K}{2}(m(J) - b_K) \\ &\geq \frac{K}{2}m(J). \end{aligned}$$

Again, we are able to conclude that the interval J_i grows exponentially with i until it covers \mathbb{S} .

As a final remark, the value of K was arbitrary. We see that, for arbitrary L and for large L , it is enough to assume $\varepsilon > m(I_{N+1})/2$ and $\varepsilon > b_2/2$, respectively, for the theorem to hold. \square

3. LYAPUNOV EXPONENT

In this section we prove our main results, Theorems 2 and 5. Before that, we present a short proof of Proposition 4 on the existence of negative Lyapunov exponents for moderate size perturbations.

Proof of Proposition 4. Fix $i \in \{1, \dots, N\}$. If L is sufficiently large, the map τ_a has a critical point $z \in \mathbb{S}$ close to c_i . Now, tune $a_z \in \mathbb{S}$ so that $\tau_{a_z}(z) = z$. Taylor expanding at z ,

$$\tau_a^\omega(x) = \omega + a - a_z + \tau_{a_z}(x) = \omega + a - a_z + z + L \int_z^x (x-t)\psi''(t) dt.$$

We see that

$$\tau_a^\omega(B_\nu(z)) \subset B_\nu(z)$$

for any $\omega \in [-\varepsilon, \varepsilon]$, provided that

$$\varepsilon + |a - a_z| + \frac{L}{2} \sup |\psi''| \nu^2 \leq \nu.$$

Moreover,

$$|\tau_a(x) - \tau_a(y)| \leq ML\nu |x - y| \quad \forall x, y \in B_\nu(z), \quad (10)$$

where the constant M only depends on ψ . Let us now choose $\nu = \frac{1}{2ML}$ and $\varepsilon = \frac{\nu}{3}$. For any a with $|a - a_z| \leq \frac{\nu}{3}$ and a large enough L , any realization of the random map $\tau_a + Y$ maps the interval $B_\nu(z)$ inside itself. By the same argument as in the beginning of Section 2, the map $\tau_a|_{B_\nu(z)} + Y$ has an invariant measure μ . This is an invariant measure for $\tau_a + Y$ supported on a subset of $B_\nu(z)$. Since (10) implies that $|\tau'_a| \leq \frac{1}{2}$ on $B_\nu(z)$, we obtain directly from (1) the bound

$$\lambda_a((\omega_n)_{n=1}^\infty, x; L) \leq -\log 2$$

for $(\eta^{\mathbb{Z}^+} \times \mu)$ -a.e. $((\omega_n)_{n=1}^\infty, x)$. \square

To estimate the Lyapunov exponent from below, we first need to bound the invariant density ρ from above.

Notice that $p(x, \cdot)$ in (6) is a Borel probability measure and that it has the representation

$$p(x, A) = \int_A \phi(x, y) dm(y),$$

where the density $\phi(x, \cdot)$ is the Radon–Nikodym derivative

$$\phi(x, y) = \left. \frac{dp(x, \cdot)}{dm} \right|_y = \frac{1}{2\varepsilon} \mathbf{1}_{B_\varepsilon(\tau_a(x))}(y).$$

Iterating (7) once,

$$\mu(A) = \int_{\mathbb{S}} p(x, A) d\mu(x) = \iint_{\mathbb{S} \times \mathbb{S}} p(x, dy) p(y, A) d\mu(x). \quad (11)$$

Recall that μ is absolutely continuous with density ρ . Thus,

$$\lim_{\delta \rightarrow 0^+} \frac{\mu(B_\delta(x_0))}{m(B_\delta(x_0))} = \rho(x_0).$$

Applying the bounded convergence theorem to (11) with $A = B_\delta(x_0)$,

$$\rho(x_0) = \int_{\mathbb{S}} \phi(x, x_0) d\mu(x) = \iint_{\mathbb{S} \times \mathbb{S}} \phi(x, y) \phi(y, x_0) dm(y) d\mu(x).$$

The first equality immediately yields the bound

$$\rho(x_0) \leq \frac{1}{2\varepsilon}, \quad (12)$$

whereas the second one shows that

$$\rho(x_0) \leq \max_x \int_{\mathbb{S}} \phi(x, y) \phi(y, x_0) dm(y).$$

Here

$$\phi(x, y) \phi(y, x_0) = \frac{1}{4\varepsilon^2} 1_{B_\varepsilon(\tau_a(x))}(y) \cdot 1_{B_\varepsilon(\tau_a(y))}(x_0)$$

so that

$$\rho(x_0) \leq \frac{1}{4\varepsilon^2} \max_z m(B_\varepsilon(z) \cap \tau_a^{-1} B_\varepsilon(x_0)). \quad (13)$$

It turns out that we also need the following estimate.

Lemma 7. *There exists a constant $C > 0$ such that, for sufficiently large values of $L > 0$,*

$$\int_{I_1} \log |\tau'_a| dm \geq -CL^{-1}.$$

Proof. Recall that the critical points of ψ are nondegenerate and observe that the set I_1 is precisely $\{x \in \mathbb{S} : -2/L \leq \psi'(x) \leq 0\}$. Thus, for large L , I_1 consists of N disjoint intervals, none of which contains any zeros of ψ'' . Thus, $\inf_{I_1} |\psi''| > 0$. Moreover, I_1 is the union of $2N$ intervals $I_1^{(1)}, \dots, I_1^{(2N)}$ on the interior of each of which $|\tau'_a|$ is one-to-one and onto $(0, 1)$. Therefore, by the change of variables $t = |\tau'_a|$ and the fact that $\int_0^1 \log t dt = -1$,

$$\begin{aligned} \int_{I_1} \log |\tau'_a| dm &= \sum_{i=1}^{2N} \int_{I_1^{(i)}} \log |\tau'_a| dm \geq \sum_{i=1}^{2N} \frac{1}{\inf_{I_1^{(i)}} |\tau''_a|} \int_0^1 \log t dt \\ &\geq -\frac{2N}{\inf_{I_1} |\tau''_a|} \geq -\frac{2N}{L \inf_{I_1} |\psi''|}. \end{aligned}$$

Since ψ is independent of any parameters, the proof is complete. \square

Proof of Theorem 2. We first consider the case $\beta \in (0, 1)$, i.e., $\beta \neq 1$. Since $\varepsilon \geq CL^{\beta-1}$, the bound (12) on the density ρ of μ , together with (4), yields

$$\rho \leq CL^{1-\beta} \quad \text{and} \quad \mu(I_{L\hat{\beta}}) \leq \sup \rho \cdot m(I_{L\hat{\beta}}) \leq CL^{\hat{\beta}-\beta}$$

uniformly for $\hat{\beta} \in (0, \beta)$. The conditions of Theorem 1 are satisfied for large enough L . We can therefore use the formula in (2) for ergodic measures μ to bound the Lyapunov

exponent $\lambda_a(L)$:

$$\begin{aligned}
 \lambda_a(L) &\geq \int_{(I_{L^{\hat{\beta}}})^c} \log |\tau'_a| d\mu + \int_{I_1} \log |\tau'_a| d\mu \\
 &\geq \left(1 - \mu(I_{L^{\hat{\beta}}})\right) \log L^{\hat{\beta}} + \sup_{I_1} \rho \cdot \int_{I_1} \log |\tau'_a| dm \\
 &\geq (1 - CL^{\hat{\beta}-\beta}) \log L^{\hat{\beta}} - CL^{1-\beta} \cdot CL^{-1} \\
 &\geq \left((1 - o(1))\hat{\beta} - o(1)\right) \log L.
 \end{aligned}$$

Above, Lemma 7 was used to bound the last integral. Since $\hat{\beta}$ can be chosen arbitrarily close to β , the proof is complete for $\beta \neq 1$.

In order to analyze the case $\beta = 1$, we replace $I_{L^{\hat{\beta}}}$ by $I_{h(L)}$, where $h(L) = L/\log L$. Notice that $\varepsilon \geq C$ results in

$$\rho \leq C \quad \text{and} \quad \mu(I_{h(L)}) \leq C/\log L$$

by the same arguments as above. Therefore,

$$\begin{aligned}
 \lambda_a(L) &\geq \int_{(I_{h(L)})^c} \log |\tau'_a| d\mu + \int_{I_1} \log |\tau'_a| d\mu \\
 &\geq \left(1 - \mu(I_{h(L)})\right) \log(L/\log L) + \sup_{I_1} \rho \cdot \int_{I_1} \log |\tau'_a| dm \\
 &\geq (1 - C/\log L)(\log L - \log \log L) - CL^{-1} \\
 &\geq (1 - o(1)) \log L,
 \end{aligned}$$

which proves the theorem also for $\beta = 1$. \square

Proof of Theorem 5. Below, we will specify a set A_L , taking a from which a lower bound on $\lambda_a(L)$ can be deduced.

It will be helpful to keep in mind that, for large enough L , there are precisely N critical points of τ_a , all nondegenerate, which are $O(L^{-1})$ units apart from the critical points c_1, \dots, c_N of the map ψ .

Recall that ε depends on L . Let us first assume that there exists a non-increasing positive function $\varepsilon_0(L)$, and point out the existence of a constant $C > 0$, such that

$$C^{-1}L^{-1} \leq \varepsilon \leq \varepsilon_0(L) \quad \text{for any sufficiently large } L \quad \text{and} \quad \lim_{L \rightarrow \infty} \varepsilon_0(L) = 0.$$

For any pair $K_1, K_2 \geq 1$ and any $\varepsilon > 0$ we define

$$A_{L,\varepsilon}^{K_1,K_2} = \{a \in [0, 1) : B_\varepsilon(I_{K_2}) \cap \tau_a(I_{K_1}) = \emptyset\}. \quad (14)$$

Because of the monotonicity of the set $A_{L,\varepsilon}^{K_1,K_2}$ with respect to ε ,

$$A_L^{K_1,K_2} := A_{L,\varepsilon_0(L)}^{K_1,K_2} = \bigcap_{\varepsilon \leq \varepsilon_0(L)} A_{L,\varepsilon}^{K_1,K_2}.$$

For large enough L and any $\varepsilon \leq \varepsilon_0(L)$, $B_\varepsilon(I_{K_2})$ is the union of N disjoint intervals almost centered at the points c_1, \dots, c_N , which do not depend on the parameter a .

Moreover, $\tau_a(I_{K_1})$ is the union of at most N intervals in \mathbb{S} which can be obtained from $\tau_0(I_{K_1})$ by a rigid rotation. Moreover, using (4), we obtain

$$m(B_\varepsilon(I_K)) \leq CKL^{-1} + 2\varepsilon \quad \text{and} \quad m(\tau_a(I_K)) \leq CK^2L^{-1},$$

if K/L is small enough. These observations together yield

$$\begin{aligned} m(A_L^{K_1, K_2}) &\geq 1 - Nm(\tau_a(I_{K_1})) - Nm(B_{\varepsilon_0(L)}(I_{K_2})) \\ &\geq 1 - CK_1^2L^{-1} - CK_2L^{-1} - 2\varepsilon_0(L). \end{aligned} \tag{15}$$

Also note that the complement of $A_L^{K_1, K_2}$ is the union of at most N^2 intervals, meaning that the same is true of $A_L^{K_1, K_2}$ itself.

Since the conditions of Theorem 1 are assumed, we can use the formula in (2) for ergodic measures μ to bound the Lyapunov exponent $\lambda_a(L)$:

$$\begin{aligned} \lambda_a(L) &\geq \int_{(I_{K_2})^c} \log |\tau'_a| d\mu + \int_{I_1} \log |\tau'_a| d\mu \\ &\geq (1 - \mu(I_{K_2})) \log K_2 + \sup_{I_1} \rho \cdot \int_{I_1} \log |\tau'_a| dm \\ &\geq \left(1 - \sup_{I_{K_2}} \rho \cdot CK_2L^{-1}\right) \log K_2 - \sup_{I_1} \rho \cdot CL^{-1}, \end{aligned}$$

where on the last line Lemma 7 and (4) have been used.

To resume the above estimate, we use the upper bound (13) on the invariant density ρ on $I_{K_2} \supset I_1$. Let $x_0 \in I_{K_2}$. Note that $\tau_a^{-1}B_\varepsilon(x_0)$ consists of finitely many disjoint intervals, and thus so does its complementary set. We label all these intervals of \mathbb{S} clockwise by J_1, \dots, J_{2k} , $J_{2k+1} = J_1$ so that, for any $i \in \{1, \dots, k\}$,

$$J_{2i-1} \subset \tau_a^{-1}B_\varepsilon(x_0) \quad \text{while} \quad J_{2i} \subset (\tau_a^{-1}B_\varepsilon(x_0))^c.$$

For $a \in A_L^{K_1, K_2}$, we have

$$J_{2i-1} \cap I_{K_1} = \emptyset, \quad \text{i.e.,} \quad |\tau'_a|_{J_{2i-1}} \geq K_1,$$

such that

$$m(J_{2i-1}) \leq 2\varepsilon/K_1.$$

For any point $z \in \mathbb{S}$, the interval $B_\varepsilon(z)$ can overlap with no more than

$$M = 2\varepsilon \left(\min \left\{ m(J_i) + m(J_{i+1}) : 1 \leq i \leq 2k \ \& \ \tau'_a(x) \neq 0 \ \forall x \in J_i \cup J_{i+1} \right\} \right)^{-1} + 1 + N$$

of the intervals J_{2i-1} . Here N is the number of those intervals J_{2i} which contain a critical point of τ_a . On the other hand, if $\tau'_a \neq 0$ on J_{2i} , τ_a maps the interval J_{2i} onto $(B_\varepsilon(x_0))^c$. In this case, the bound $|\tau'_a(x)| \leq CL$ implies $m(J_{2i}) \geq C^{-1}L^{-1}(1 - 2\varepsilon)$. As $C^{-1}L^{-1} \leq \varepsilon \leq \frac{1}{3}$ holds for large L ,

$$M \leq CL\varepsilon$$

uniformly in z and L , for such L . Therefore, (13) shows that

$$\sup_{I_{K_2}} \rho \leq \frac{CL\varepsilon}{4\varepsilon^2} m(J_{2i-1}) \leq C \frac{L}{K_1},$$

which in combination with the earlier bound on $\lambda_a(L)$ results in

$$\begin{aligned} \lambda_a(L) &\geq \left(1 - C \frac{K_2}{K_1}\right) \log K_2 - C \frac{1}{K_1} \\ &\geq \left(1 - C \frac{K_2}{K_1}\right) \log K_2. \end{aligned} \tag{16}$$

Finally, define

$$\varepsilon_0(L) = L^{-1/2}, \quad K_1 = (L/\log L)^{1/2}, \quad \text{and} \quad K_2 = L^{1/2}/\log L.$$

Then the parameter set

$$A_L = A_L^{K_1, K_2}$$

has all desired properties, as can be checked using (15) and (16), so that the theorem has been verified in the special case in which $\varepsilon \leq \varepsilon_0(L)$ holds for all large L . Now, assume that $\varepsilon > \varepsilon_0(L)$ for an unbounded set of values of L and observe that

$$\liminf_{L \rightarrow \infty} \inf_{a \in A_L} \frac{\lambda_a(L)}{\log L} = \min \left(\liminf_{\substack{L \rightarrow \infty \\ L: \varepsilon \leq \varepsilon_0(L)}} \inf_{a \in A_L} \frac{\lambda_a(L)}{\log L}, \liminf_{\substack{L \rightarrow \infty \\ L: \varepsilon > \varepsilon_0(L)}} \inf_{a \in A_L} \frac{\lambda_a(L)}{\log L} \right).$$

The theorem follows by combining the previous special case with Theorem 2. \square

REFERENCES

- [1] Peter H. Baxendale, *Stability and equilibrium properties of stochastic flows of diffeomorphisms*, (Charlotte, NC, 1990), Progr. Probab., vol. 27, Birkhäuser Boston, Boston, MA, 1992, pp. 3–35. MR1187984 (93h:58167) \uparrow 2, 5
- [2] Michael Benedicks and Lennart Carleson, *On iterations of $1 - ax^2$ on $(-1, 1)$* , Ann. of Math. (2) **122** (1985), no. 1, 1–25, DOI 10.2307/1971367. MR799250 (87c:58058) \uparrow 1
- [3] Michael Benedicks and Marcelo Viana, *Random perturbations and statistical properties of Hénon-like maps*, Ann. Inst. H. Poincaré Anal. Non Linéaire **23** (2006), no. 5, 713–752, DOI 10.1016/j.anihpc.2004.10.013. MR2259614 (2007f:37042) \uparrow 2
- [4] Michael Benedicks and Lai-Sang Young, *Absolutely continuous invariant measures and random perturbations for certain one-dimensional maps*, Ergodic Theory Dynam. Systems **12** (1992), no. 1, 13–37, DOI 10.1017/S0143385700006556. MR1162396 (93d:58087) \uparrow 2
- [5] William Cowieson and Lai-Sang Young, *SRB measures as zero-noise limits*, Ergodic Theory Dynam. Systems **25** (2005), no. 4, 1115–1138, DOI 10.1017/S0143385704000604. MR2158399 (2006f:37030) \uparrow 2
- [6] Jacek Graczyk and Grzegorz Świątek, *Generic hyperbolicity in the logistic family*, Ann. of Math. (2) **146** (1997), no. 1, 1–52, DOI 10.2307/2951831. MR1469316 (99b:58079) \uparrow 1
- [7] Michael V. Jakobson, *Absolutely continuous invariant measures for one-parameter families of one-dimensional maps*, Comm. Math. Phys. **81** (1981), no. 1, 39–88. MR630331 (83j:58070) \uparrow 1
- [8] Anatole Katok and Yuri Kifer, *Random perturbations of transformations of an interval*, J. Analyse Math. **47** (1986), 193–237. MR874051 (88c:58053) \uparrow 2
- [9] Yuri Kifer, *Ergodic theory of random transformations*, Progress in Probability and Statistics, vol. 10, Birkhäuser Boston Inc., Boston, MA, 1986. MR884892 (89c:58069) \uparrow 6
- [10] ———, *Random perturbations of dynamical systems*, Progress in Probability and Statistics, vol. 16, Birkhäuser Boston Inc., Boston, MA, 1988. MR1015933 (91e:58159) \uparrow 2
- [11] Oleg Kozlovski, Weixiao Shen, and Sebastian van Strien, *Density of hyperbolicity in dimension one*, Ann. of Math. (2) **166** (2007), no. 1, 145–182, DOI 10.4007/annals.2007.166.145. MR2342693 (2008j:37081) \uparrow 1

- [12] François Ledrappier and Lai-Sang Young, *Entropy formula for random transformations*, Probab. Theory Related Fields **80** (1988), no. 2, 217–240, DOI 10.1007/BF00356103. MR968818 (90d:58079) ↑2
- [13] Yves Le Jan, *On isotropic Brownian motions*, Z. Wahrsch. Verw. Gebiete **70** (1985), no. 4, 609–620, DOI 10.1007/BF00531870. MR807340 (87a:60090) ↑2, 5
- [14] Kevin Lin and Lai-Sang Young, *Dynamics of periodically kicked oscillators*, J. Fixed Point Theory Appl. **7** (2010), no. 2, 291–312, DOI 10.1007/s11784-010-0025-9. ↑3
- [15] Mikhail Lyubich, *Regular and stochastic dynamics in the real quadratic family*, Proc. Natl. Acad. Sci. USA **95** (1998), no. 24, 14025–14027 (electronic), DOI 10.1073/pnas.95.24.14025. MR1661281 (2000c:37060) ↑1
- [16] ———, *Dynamics of quadratic polynomials. I, II*, Acta Math. **178** (1997), no. 2, 185–247, 247–297, DOI 10.1007/BF02392694. MR1459261 (98e:58145) ↑1
- [17] ———, *Almost every real quadratic map is either regular or stochastic*, Ann. of Math. (2) **156** (2002), no. 1, 1–78, DOI 10.2307/3597183. MR1935840 (2003i:37032) ↑1
- [18] William Ott and Mikko Stenlund, *From limit cycles to strange attractors*, Comm. Math. Phys. **296** (2010), no. 1, 215–249, DOI 10.1007/s00220-010-0994-y. MR2606633 ↑3
- [19] Lev Semenovich Pontryagin, Aleksandr Aleksandrovich Andronov, and Aleksand Adolfovich Vitt, *On statistical consideration of dynamical systems*, J. Experiment. Theor. Phys. **3** (1933), no. 3, 165–180. In Russian. ↑2
- [20] Marek Ryszard Rychlik, *Another proof of Jakobson’s theorem and related results*, Ergodic Theory Dynam. Systems **8** (1988), no. 1, 93–109, DOI 10.1017/S014338570000434X. MR939063 (90b:58149) ↑1
- [21] Yakov Grigorevich Sinai, *Kolmogorov’s work on ergodic theory*, Ann. Probab. **17** (1989), no. 3, 833–839. MR1009437 (91e:01017) ↑2
- [22] Masato Tsujii, *Positive Lyapunov exponents in families of one-dimensional dynamical systems*, Invent. Math. **111** (1993), no. 1, 113–137, DOI 10.1007/BF01231282. MR1193600 (93j:58081) ↑1
- [23] Qiudong Wang and Lai-Sang Young, *From invariant curves to strange attractors*, Comm. Math. Phys. **225** (2002), no. 2, 275–304, DOI 10.1007/s002200100582. MR1889226 (2003e:37045) ↑3
- [24] ———, *Strange attractors in periodically-kicked limit cycles and Hopf bifurcations*, Comm. Math. Phys. **240** (2003), no. 3, 509–529. MR2005855 (2004i:37070) ↑3
- [25] ———, *Nonuniformly expanding 1D maps*, Comm. Math. Phys. **264** (2006), no. 1, 255–282, DOI 10.1007/s00220-005-1485-4. MR2212223 (2006k:37103) ↑1, 3
- [26] Lai-Sang Young, *Stochastic stability of hyperbolic attractors*, Ergodic Theory Dynam. Systems **6** (1986), no. 2, 311–319, DOI 10.1017/S0143385700003473. MR857204 (88a:58160) ↑2
- [27] ———, *Chaotic phenomena in three settings: large, noisy and out of equilibrium*, Nonlinearity **21** (2008), no. 11, T245–T252, DOI 10.1088/0951-7715/21/11/T04. MR2448225 (2009m:37001) ↑2

(Zeng Lian) COURANT INSTITUTE OF MATHEMATICAL SCIENCES, NEW YORK, NY 10012, USA.
E-mail address: lian@cims.nyu.edu

(Mikko Stenlund) COURANT INSTITUTE OF MATHEMATICAL SCIENCES, NEW YORK, NY 10012, USA;
 DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O. Box 68, FIN-00014 UNIVERSITY OF HELSINKI, FINLAND.

E-mail address: mikko@cims.nyu.edu

URL: <http://www.math.helsinki.fi/mathphys/mikko.html>