

Non-equilibrium Steady States of Quantum Systems on Star Graphs

Mihail Mintchev

Istituto Nazionale di Fisica Nucleare and Dipartimento di Fisica, Università di Pisa, Largo Pontecorvo 3, 56127 Pisa, Italy

Abstract

Non-equilibrium steady states of quantum fields on star graphs are explicitly constructed. These states are parametrized by the temperature and the chemical potential, associated with each edge of the graph. Time reversal invariance is spontaneously broken. We study in this general framework the transport properties of the Schrödinger and the Dirac systems on a star graph, modeling a quantum wire junction. The interaction, which drives the system away from equilibrium, is localized in the vertex of the graph. All point-like vertex interactions, giving rise to self-adjoint Hamiltonians possibly involving the minimal coupling to a static electromagnetic field in the ambient space, are considered. In this context we compute the exact electric steady current and the non-equilibrium charge density. We investigate also the heat transport and derive the Casimir energy density away from equilibrium. The appearance of Friedel type oscillations of the charge and energy densities along the edges of the graph is established. We focus finally on the noise power and discuss the non-trivial impact of the point-like interactions on the noise.

1 Introduction

Transport properties of quantum wire junctions attracted in the last two decades much attention [1]-[32]. The experimental realizations of quantum wires include nowadays carbon nanotubes, semiconductor, metallic and polymer nanowires, and quantum Hall edges. While the equilibrium features of these devices has been extensively explored, there is recently a growing interest in non-equilibrium phenomena. A typical problem in this context is schematically represented in Fig. 1. A quantum wire junction is modeled by a star graph Γ with n edges (leads) E_i , each of them connected to a heat reservoir (bath) with (inverse) temperature β_i and chemical potential μ_i . Assume for simplicity that E_i are infinite and that the interaction, described by a scattering matrix $\mathbb{S}(k)$, is localized at the vertex V of Γ . The system is away from equilibrium if $\mathbb{S}(k)$ admits at least one non-trivial transmission coefficient among edges with different temperature and/or chemical potential. Two main questions arise at this point. The first one concerns the existence of a steady state describing the above situation. Provided that such a state exists, it is natural to ask about the general features of the system in this state. In the present paper we show that the first question has an affirmative answer, constructing explicitly a suitable steady state $\Omega_{\beta,\mu}$, which is parametrized by $\beta = (\beta_1, \dots, \beta_n)$ and $\mu = (\mu_1, \dots, \mu_n)$ and captures the non-equilibrium properties of the system. Afterwards we derive the expectation values of several basic physical observables (currents, charge and energy densities,...) in $\Omega_{\beta,\mu}$, which characterize the system and thus answer the second question.




Figure 1: A star graph Γ with scattering matrix $\mathbb{S}(k)$ at the vertex V and edges E_i connected at infinity to thermal reservoirs with inverse temperature β_i and chemical potential μ_i .

It is perhaps useful to recall that the systems admitting non-equilibrium steady states (NESS's) [33] represent an important subclass of the large family of non-equilibrium systems. Unlike in equilibrium, a system in a NESS admits steady currents. Nevertheless, all macroscopic properties are still time independent like in equilibrium.

The formulation of a suitable statistical mechanical framework for treating NESS's has not been yet completed and is intensively investigated [34]-[40]. We follow below a microscopic

approach and address the question using quantum field theory methods. Our construction of NESS generalizes that of a Gibbs state in finite temperature quantum field theory. We develop first a general algebraic framework, which applies for any dispersion relation, thus covering both the relativistic and non-relativistic cases. As already mentioned, we consider for simplicity fields which propagate freely in the bulk $\Gamma \setminus V$ of the graph, the interaction being localized exclusively in the vertex V . We discuss all possible point-like interactions for which the Hamiltonian of our system, being a Hermitian operator in the bulk $\Gamma \setminus V$, extends to a self-adjoint operator on the whole graph Γ . We use at this point of our construction some simple results [41]-[43] from the spectral theory of differential operators on graphs (known also as “quantum graphs”).

It is worth mentioning that non-equilibrium systems of the type shown in Fig. 1 have been investigated in the past by two different methods. The first one is the scattering approach initiated by Landauer [44] and completed by Büttiker [45]. The second one is based on the linear response theory [46], originally developed by Kubo for macroscopic samples and extended later by Baranger and Stone [47] to mesoscopic systems. The equivalence between these two approaches has been demonstrated in Ref. [48].

The construction of the NESS $\Omega_{\beta,\mu}$, developed in this paper, is formulated in purely algebraic terms and adopts a deformation of the algebra of canonical (anti)commutation relations. No approximations, like linear response theory, are involved. The states $\Omega_{\beta,\mu}$ allow for a unified description of systems with different dynamics. We illustrate this fact by treating below both the Schrödinger and Dirac equations in the same way. The abstract construction of $\Omega_{\beta,\mu}$ is first tested by reproducing the Landauer-Büttiker (L-B) steady current. Afterwards, taking advantage of the exact solvability of the point-like interactions in the state $\Omega_{\beta,\mu}$, we investigate several new features of the non-equilibrium dynamics, applying the general framework to other physical observables. We compute the charge density in $\Omega_{\beta,\mu}$ and show the presence of characteristic Friedel type oscillations along the leads. We also present a fully microscopic-based calculation of the Casimir effect in a quantum wire junction away from equilibrium. In this context we establish the non-equilibrium Stefan-Boltzmann law and derive the heat (energy) transport. We also compute the exact two-point current-current correlation function away from equilibrium and investigate the noise power. In particular, we discuss the impact of the point-like interaction in the junction on the behavior of both thermal and shot noises. Finally, we generalize all results in the presence of a static electromagnetic field in the ambient space.

The paper is organized as follows. In the next section we develop a simple algebraic framework for the construction of steady states of quantum wire junctions. Section 3 is devoted to the Schrödinger junction. In section 4 we change the dynamics, considering the Dirac equation and the relative steady states on the junction. The novelty here is the presence of antiparticles which contribution and role are considered in detail. In section 5 we discuss the results and comment on the possible further developments.

2 Algebraic construction of the states $\Omega_{\beta,\mu}$

Previous investigations [18]-[21], [49] have shown that a convenient coordinate system for expressing quantum fields on star graphs is provided by the deformed algebras \mathcal{A}_{\pm} of canonical

(anti)commutation relations, generated by $\{a_i(k), a_i^*(k) : k \in \mathbb{R}, i = 1, \dots, n\}$ which satisfy

$$[a_i(k), a_j(p)]_{\pm} = [a_i^*(k), a_j^*(p)]_{\pm} = 0, \quad (2.1)$$

$$[a_i(k), a_j^*(p)]_{\pm} = 2\pi[\delta(k-p)\delta_{ij} + \mathbb{S}_{ij}(k)\delta(k+p)], \quad (2.2)$$

and the constraints

$$a_i(k) = \sum_{j=1}^n \mathbb{S}_{ij}(k)a_j(-k), \quad a_i^*(k) = \sum_{j=1}^n a_j^*(-k)\mathbb{S}_{ji}(-k). \quad (2.3)$$

The index \pm in (2.1, 2.2) refers to Fermi/Bose statistics and $\mathbb{S}(k)$ is the $n \times n$ scattering matrix describing the point-like interaction in the vertex of the graph. We assume in what follows unitarity

$$\mathbb{S}(k)\mathbb{S}(k)^* = \mathbb{I} \quad (2.4)$$

and Hermitian analyticity [50]-[53]

$$\mathbb{S}(k)^* = \mathbb{S}(-k). \quad (2.5)$$

The latter implies that the $*$ -operation is a conjugation in \mathcal{A}_{\pm} . Combining (2.4) and (2.5) one concludes that $\mathbb{S}(k)\mathbb{S}(-k) = \mathbb{I}$, ensuring the consistency of the constraints (2.3).

Our main goal now is to construct $\Omega_{\beta,\mu}$ as a state, i.e. as a (continuous) linear functional over \mathcal{A}_{\pm} . We recall in this respect that \mathcal{A}_{\pm} is a simplified version of the so called reflection-transmission algebra [50]-[53], describing factorized scattering in integrable models with point-like defects in one dimension. The Fock and the (grand canonical) Gibbs state over \mathcal{A}_{\pm} describe equilibrium physics and have been largely explored [50]-[53]. The physical input for constructing the new states $\Omega_{\beta,\mu}$ is the observation that the sub-algebras $\mathcal{A}_{\pm}^{\text{in}}$ and $\mathcal{A}_{\pm}^{\text{out}}$, generated by $\{a_i(k), a_i^*(k) : k < 0\}$ and $\{a_i(k), a_i^*(k) : k > 0\}$ respectively, parametrize the asymptotic incoming and outgoing fields. Accordingly, both $\mathcal{A}_{\pm}^{\text{in}}$ and $\mathcal{A}_{\pm}^{\text{out}}$ are conventional canonical (anti)commutation relation algebras. Indeed, the $\delta(k+p)$ term in (2.2) vanishes if both momenta are negative or positive. Notice also that the constraints (2.3) relate $\mathcal{A}_{\pm}^{\text{in}}$ with $\mathcal{A}_{\pm}^{\text{out}}$. It is crucial for what follows that the whole reflection-transmission algebra \mathcal{A}_{\pm} can be generated either by $\mathcal{A}_{\pm}^{\text{in}}$ or by $\mathcal{A}_{\pm}^{\text{out}}$ via (2.3). Our strategy for constructing the NESS $\Omega_{\beta,\mu}$ is based on this kind of asymptotic completeness property. In fact, we will start with an equilibrium state on $\mathcal{A}_{\pm}^{\text{in}}$ and extend it by means of (2.3) to a non-equilibrium state on \mathcal{A}_{\pm} .

The first step is to describe the *asymptotic* dynamics and symmetries at $t = -\infty$ (i.e. before the interaction) in terms of $\mathcal{A}_{\pm}^{\text{in}}$. Since the asymptotic fields are free, it is natural to introduce the edge Hamiltonians

$$h_i = \int_{-\infty}^0 \frac{dk}{2\pi} \omega_i(k) a_i^*(k) a_i(k) \quad (2.6)$$

and edge charges

$$q_i = \int_{-\infty}^0 \frac{dk}{2\pi} a_i^*(k) a_i(k), \quad (2.7)$$

where $\omega_i(k) \geq 0$ is the dispersion relation in the edge E_i . At this point we define

$$K = \sum_{i=1}^n \beta_i (h_i - \mu_i q_i). \quad (2.8)$$

and introduce the equilibrium Gibbs state over $\mathcal{A}_{\pm}^{\text{in}}$ in the standard way [54]. For any polynomial \mathcal{P} over $\mathcal{A}_{\pm}^{\text{in}}$ we set

$$(\Omega_{\beta,\mu}, \mathcal{P}(a_i^*(k_i), a_j(p_j))\Omega_{\beta,\mu}) \equiv \langle \mathcal{P}(a_i^*(k_i), a_j(p_j)) \rangle_{\beta,\mu} = \frac{1}{Z} \text{Tr} [e^{-K} \mathcal{P}(a_i^*(k_i), a_j(p_j))] , \quad (2.9)$$

where $k_i < 0$, $p_j < 0$ and $Z = \text{Tr} (e^{-K})$. It is well known [54] that one can compute the expectation values (2.9) by purely algebraic manipulations and that all these expectation values can be expressed in terms of the two-point functions, which are written in terms of the familiar Fermi/Bose distributions in the following way

$$\langle a_j^*(p) a_i(k) \rangle_{\beta,\mu} = \frac{e^{-\beta_i[\omega_i(k)-\mu_i]}}{1 \pm e^{-\beta_i[\omega_i(k)-\mu_i]}} \delta_{ij} 2\pi \delta(k-p) , \quad (2.10)$$

$$\langle a_i(k) a_j^*(p) \rangle_{\beta,\mu} = \frac{1}{1 \pm e^{-\beta_i[\omega_i(k)-\mu_i]}} \delta_{ij} 2\pi \delta(k-p) . \quad (2.11)$$

We stress that (2.10,2.11) hold on $\mathcal{A}_{\pm}^{\text{in}}$, i.e. only for *negative* momenta.

The second step is to extend (2.9-2.11) to the whole algebra \mathcal{A}_{\pm} , namely to *positive* momenta. For this purpose we use the relations (2.3). One finds in this way

$$\begin{aligned} \langle a_j^*(p) a_i(k) \rangle_{\beta,\mu} = 2\pi \left\{ \left[\theta(-k) d_i^{\pm}(k) \delta_{ij} + \theta(k) \sum_{l=1}^n \mathbb{S}_{il}(k) d_l^{\pm}(-k) \mathbb{S}_{lj}(-k) \right] \delta(k-p) \right. \\ \left. + \left[\theta(-k) d_i^{\pm}(k) \mathbb{S}_{ij}(k) + \theta(k) \mathbb{S}_{ij}(k) d_j^{\pm}(-k) \right] \delta(k+p) \right\} , \end{aligned} \quad (2.12)$$

where for simplifying the notation we introduced

$$d_i^{\pm}(k) = \frac{e^{-\beta_i[\omega_i(k)-\mu_i]}}{1 \pm e^{-\beta_i[\omega_i(k)-\mu_i]}} . \quad (2.13)$$

The explicit expression of $\langle a_i(k) a_j^*(p) \rangle_{\beta,\mu}$ is obtained from (2.12) by the substitution

$$d_i^{\pm}(k) \mapsto c_i^{\pm}(k) = \frac{1}{1 \pm e^{-\beta_i[\omega_i(k)-\mu_i]}} . \quad (2.14)$$

The final step is to compute a generic correlation function. Employing the commutation relations (2.1,2.2), one can reduce the problem to the evaluation of correlators of the form

$$\left\langle \prod_{m=1}^M a_{i_m}(k_{i_m}) \prod_{n=1}^N a_{j_n}^*(p_{j_n}) \right\rangle_{\beta,\mu} , \quad (2.15)$$

which can be computed in turn by iteration via

$$\left\langle \prod_{m=1}^M a_{i_m}(k_{i_m}) \prod_{n=1}^N a_{j_n}^*(p_{j_n}) \right\rangle_{\beta,\mu} = \delta_{MN} \sum_{m=1}^M \langle a_{i_1}(k_{i_1}) a_{j_m}^*(p_{j_m}) \rangle_{\beta,\mu} \left\langle \prod_{m=2}^M a_{i_m}(k_{i_m}) \prod_{\substack{n=1 \\ n \neq m}}^N a_{j_n}^*(p_{j_n}) \right\rangle_{\beta,\mu} . \quad (2.16)$$

In conclusion, we emphasize once more that the use of the deformed algebras \mathcal{A}_\pm in the construction of $\Omega_{\beta,\mu}$ represents only a convenient choice of coordinates, which has the advantage to be universal and to apply to a variety of systems characterized by a scattering matrix $\mathbb{S}(k)$. In support of this statement, we consider below the Schrödinger and the Dirac equations on the star graph Γ .

3 The Schrödinger junction

3.1 Preliminaries

In this section we apply the general algebraic construction of the state $\Omega_{\beta,\mu}$ to the Schrödinger system on a star graph Γ with point-like interactions in the vertex V of Γ . We will consider Fermi statistics and for simplifying the notation will omit the apex $+$ in the Dirac distributions $d_i^+(k)$ and $c_i^+(k)$. As observed in section 3.5 below, most of the results can be easily extended to Bose statistics.

We start by summarizing the main features [18] of the Schrödinger equation on Γ , recalling the description of all point-like interactions leading to a self-adjoint Hamiltonian. Each point P in the bulk of Γ is parametrized by (x, i) , where $x > 0$ is the distance of P from the vertex V and i labels the edge. In the bulk $\Gamma \setminus V$ of the graph the Schrödinger field $\psi(t, x, i)$ with Fermi statistics satisfies

$$\left(i\partial_t + \frac{1}{2m}\partial_x^2 \right) \psi(t, x, i) = 0, \quad (3.17)$$

with standard equal-time canonical anticommutation relations

$$[\psi(0, x_1, i_1), \psi(0, x_2, i_2)]_+ = [\psi^*(0, x_1, i_1), \psi^*(0, x_2, i_2)]_+ = 0, \quad (3.18)$$

$$[\psi(0, x_1, i_1), \psi^*(0, x_2, i_2)]_+ = \delta_{i_1 i_2} \delta(x_1 - x_2). \quad (3.19)$$

The interaction in the vertex is fixed by requiring that the bulk Hamiltonian defined by (3.17) (essentially the operator $-\partial_x^2$) admits a *self-adjoint extension* on the whole graph. According to some elementary results from the spectral theory [41]-[43] of differential operators on graphs, all such interactions are described by the boundary conditions

$$\lim_{x \rightarrow 0^+} \sum_{j=1}^n [\lambda(\mathbb{I} - \mathbb{U})_{ij} - i(\mathbb{I} + \mathbb{U})_{ij}\partial_x] \psi(t, x, j) = 0, \quad (3.20)$$

where \mathbb{U} is an arbitrary $n \times n$ unitary matrix and $\lambda \in \mathbb{R}$ is a parameter with dimension of mass. Eq. (3.20) guaranties *unitary time evolution* of the system and generalizes to the graph Γ the familiar mixed (Robin) boundary condition on the half-line \mathbb{R}_+ . The matrices $\mathbb{U} = \mathbb{I}$ and $\mathbb{U} = -\mathbb{I}$ define the Neumann and Dirichlet boundary conditions respectively.

The explicit form of the scattering matrix, expressed in terms of \mathbb{U} and λ , is [41]-[43]

$$\mathbb{S}(k) = -\frac{[\lambda(\mathbb{I} - \mathbb{U}) - k(\mathbb{I} + \mathbb{U})]}{[\lambda(\mathbb{I} - \mathbb{U}) + k(\mathbb{I} + \mathbb{U})]} \quad (3.21)$$

and has a simple physical interpretation: the diagonal element $\mathbb{S}_{ii}(k)$ represents the reflection amplitude from the vertex on the edge E_i , whereas $\mathbb{S}_{ij}(k)$ with $i \neq j$ equals the transmission amplitude from E_i to E_j . One easily verifies that (3.21) satisfies (2.4, 2.5) and therefore defines an algebra \mathcal{A}_+ of the type introduced in the previous section. Notice that

$$\mathbb{S}(\lambda) = \mathbb{U}, \quad \mathbb{S}(-\lambda) = \mathbb{U}^{-1}, \quad (3.22)$$

showing that the unitary matrix \mathbb{U} entering the boundary conditions (3.20) is actually the scattering matrix at scale λ .

A remarkable property of (3.21) is that it can be diagonalized for any k by a k -independent unitary matrix. In fact, let \mathcal{U} be the unitary matrix diagonalizing \mathbb{U} , namely

$$\mathcal{U}^{-1} \mathbb{U} \mathcal{U} = \mathbb{U}_d = \text{diag} (e^{2i\alpha_1}, e^{2i\alpha_2}, \dots, e^{2i\alpha_n}), \quad \alpha_i \in \mathbb{R}. \quad (3.23)$$

By means of (3.21) one concludes that \mathcal{U} diagonalizes $\mathbb{S}(k)$ for any k as well, and that

$$\mathbb{S}_d(k) = \mathcal{U}^* \mathbb{S}(k) \mathcal{U} = \text{diag} \left(\frac{k + i\eta_1}{k - i\eta_1}, \frac{k + i\eta_2}{k - i\eta_2}, \dots, \frac{k + i\eta_n}{k - i\eta_n} \right), \quad (3.24)$$

where

$$\eta_i = \lambda \tan(\alpha_i), \quad -\frac{\pi}{2} \leq \alpha_i \leq \frac{\pi}{2}. \quad (3.25)$$

Illustrating various aspects of the Schrödinger junction, we will use below the most general 2×2 \mathbb{S} -matrix

$$\mathbb{S}(k) = \begin{pmatrix} \frac{k^2 + ik(\eta_1 - \eta_2) \cos(\theta) + \eta_1 \eta_2}{(k - i\eta_1)(k - i\eta_2)} & \frac{-ie^{i\varphi} k(\eta_1 - \eta_2) \sin(\theta)}{(k - i\eta_1)(k - i\eta_2)} \\ \frac{-ie^{-i\varphi} k(\eta_1 - \eta_2) \sin(\theta)}{(k - i\eta_1)(k - i\eta_2)} & \frac{k^2 - ik(\eta_1 - \eta_2) \cos(\theta) + \eta_1 \eta_2}{(k - i\eta_1)(k - i\eta_2)} \end{pmatrix}, \quad (3.26)$$

where φ and θ are arbitrary angles.

The general representation (3.24) implies that $\mathbb{S}(k)$ is a meromorphic function in the complex k -plane with finite number of simple poles on the imaginary axis. For simplicity we consider in this paper the case without bound states (poles in the upper half plane), referring for the general case to [53], [28], [31] and the comments in section 3.5. In other words, we assume that

$$\int_{-\infty}^{\infty} \frac{dk}{2\pi} e^{ikx} \mathbb{S}_{ij}(k) = 0, \quad x > 0. \quad (3.27)$$

It has been shown in Ref. [18] that in this case the solution of equation (3.17) is fixed uniquely by (3.18-3.20) and takes the following simple form

$$\psi(t, x, i) = \int_{-\infty}^{\infty} \frac{dk}{2\pi} a_i(k) e^{-i\omega(k)t + ikx}, \quad \omega(k) = \frac{k^2}{2m}. \quad (3.28)$$

The equation of motion (3.17) is invariant under the time reversal operation

$$T\psi(t, x, i)T^{-1} = -\eta_T \psi(-t, x, i), \quad |\eta_T| = 1, \quad (3.29)$$

T being an anti-unitary operator. We stress however that the boundary condition (3.20) preserves the time reversal symmetry only if \mathbb{U} is symmetric [30], namely

$$\mathbb{U}^t = \mathbb{U}. \quad (3.30)$$

The electric current

$$j_x(t, x, i) = i \frac{e}{2m} [\psi^* (\partial_x \psi) - (\partial_x \psi^*) \psi] (t, x, i), \quad (3.31)$$

e being the electric charge, and the energy current

$$\theta_{xt}(t, x, i) = \frac{1}{4m} [(\partial_t \psi^*) (\partial_x \psi) + (\partial_x \psi^*) (\partial_t \psi) - (\partial_t \partial_x \psi^*) \psi - \psi^* (\partial_t \partial_x \psi)] (t, x, i), \quad (3.32)$$

are among the basic physical observables. The time components of these currents are

$$j_t(t, x, i) = e (\psi^* \psi) (t, x, i), \quad (3.33)$$

$$\theta_{tt}(t, x, i) = -\frac{1}{4m} [\psi^* (\partial_x^2 \psi) + (\partial_x^2 \psi^*) \psi] (t, x, i), \quad (3.34)$$

respectively and, as a consequence of (3.17), satisfy the local conservation laws

$$(\partial_t j_t - \partial_x j_x) (t, x, i) = (\partial_t \theta_{tt} - \partial_x \theta_{xt}) (t, x, i) = 0. \quad (3.35)$$

Equations (3.35), combined with the Kirchhoff rules

$$\sum_{i=1}^n j_x(t, 0, i) = 0, \quad \sum_{i=1}^n \theta_{xt}(t, 0, i) = 0, \quad (3.36)$$

ensure the charge and energy conservation in the system. Since the proof of (3.36) at the quantum level is a quite subtle, we provide the main steps, focussing for instance on the electric current. Using the basic definitions, one easily derives the representation

$$\begin{aligned} j_x(t, 0, i) &= i \frac{e}{2m} \int_{-\infty}^0 \frac{dk}{2\pi} \int_{-\infty}^0 \frac{dp}{2\pi} e^{it[\omega(k) - \omega(p)]} \\ &\times \sum_{j,l=1}^n a_j^*(k) \left\{ \chi_{ji}^*(k; 0) [\partial_x \chi_{il}] (p; 0) - [\partial_x \chi_{ji}^*] (k; 0) \chi_{il}(p; 0) \right\} a_l(p), \end{aligned} \quad (3.37)$$

where

$$\chi(k; x) = e^{ikx} \mathbb{I} + e^{-ikx} \mathbb{S}^*(k), \quad \chi^*(k; x) = e^{-ikx} \mathbb{I} + e^{ikx} \mathbb{S}(k). \quad (3.38)$$

The trick now is represent the right hand side of (3.38) as a boundary term of an integral over the half line, namely

$$\begin{aligned} j_x(t, 0, i) &= -i \frac{e}{2m} \int_{-\infty}^0 \frac{dk}{2\pi} \int_{-\infty}^0 \frac{dp}{2\pi} \int_0^\infty dx e^{it[\omega(k) - \omega(p)]} \\ &\times \sum_{j,l=1}^n a_j^*(k) \left\{ \chi_{ji}^*(k; x) [\partial_x^2 \chi_{il}] (p; x) - [\partial_x^2 \chi_{ji}^*] (k; x) \chi_{il}(p; x) \right\} a_l(p) = \\ &-i \frac{e}{2m} \int_{-\infty}^0 \frac{dk}{2\pi} \int_{-\infty}^0 \frac{dp}{2\pi} e^{it[\omega(k) - \omega(p)]} (k^2 - p^2) \int_0^\infty dx \sum_{j,l=1}^n a_j^*(k) [\chi_{ji}^*(k; x) \chi_{il}(p; x)] a_l(p), \end{aligned} \quad (3.39)$$

The final step is to apply the orthogonality relations

$$\sum_{i=1}^n \int_0^\infty dx \chi_{ji}^*(k; x) \chi_{il}(p; x) = 2\pi \delta_{jl} \delta(k - p), \quad (3.40)$$

which hold [18] for \mathbb{S} given by (3.21).

At this stage we are ready to investigate the properties of the system in the state $\Omega_{\beta, \mu}$. Being expressed in terms of the algebra \mathcal{A}_+ , the solution (3.28) and the observables (3.31-3.34) apply for any representation of this algebra. This fundamental universality property has been already largely explored in the Fock and the Gibbs representations of \mathcal{A}_+ , which describe equilibrium physics. In order to study the non-equilibrium properties of the Schrödinger system in Fig. 1, we apply below the representation generated by the state $\Omega_{\beta, \mu}$. Since antiparticle excitations are absent, we assume without loss of generality that $\mu_i \geq 0$.

There are two non-trivial two-point correlation functions. Using (2.12), one finds

$$\langle \psi^*(t_1, x_1, i) \psi(t_2, x_2, j) \rangle_{\beta, \mu} = \int_0^\infty \frac{dk}{2\pi} e^{i\omega(k)t_{12}} \left[\delta_{ji} d_i(k) e^{-ikx_{12}} + \mathbb{S}_{ji}(k) d_j(k) e^{ik\tilde{x}_{12}} + \mathbb{S}_{ji}^*(k) d_i(k) e^{-ik\tilde{x}_{12}} + \sum_{l=1}^n \mathbb{S}_{jl}^*(k) d_l(k) \mathbb{S}_{li}(k) e^{ikx_{12}} \right], \quad (3.41)$$

$$\langle \psi(t_1, x_1, i) \psi^*(t_2, x_2, j) \rangle_{\beta, \mu} = \int_0^\infty \frac{dk}{2\pi} e^{-i\omega(k)t_{12}} \left[\delta_{ij} c_i(k) e^{ikx_{12}} + \mathbb{S}_{ij}(k) c_i(k) e^{ik\tilde{x}_{12}} + \mathbb{S}_{ij}^*(k) c_j(k) e^{-ik\tilde{x}_{12}} + \sum_{l=1}^n \mathbb{S}_{il}^*(k) c_l(k) \mathbb{S}_{lj}(k) e^{-ikx_{12}} \right], \quad (3.42)$$

where $t_{12} = t_1 - t_2$, $x_{12} = x_1 - x_2$ and $\tilde{x}_{12} = x_1 + x_2$. The invariance of (3.41, 3.42) under time translation implies energy conservation. For systems away from equilibrium one expects that the time reversal (3.29) symmetry is instead broken. In fact,

$$\langle \psi^*(t_1, x_1, i) \psi(t_2, x_2, j) \rangle_{\beta, \mu} \neq \langle \psi^*(-t_2, x_2, j) \psi(-t_1, x_1, i) \rangle_{\beta, \mu}. \quad (3.43)$$

showing that $T\Omega_{\beta, \mu} \neq \Omega_{\beta, \mu}$, i.e. that $\Omega_{\beta, \mu}$ breaks down spontaneously time reversal invariance even if (3.30) holds.

3.2 Charge transport and density

Using the explicit form of the correlation function (3.41) and applying standard point-splitting technique, one obtains

$$J_i(\beta, \mu) \equiv \langle j_x(t, x, i) \rangle_{\beta, \mu} = \lim_{\substack{t_1 \rightarrow t_2 = t \\ x_1 \rightarrow x_2 = x}} i \frac{e}{2m} [\langle \psi^*(t_1, x_1, i) \partial_{x_2} \psi(t_2, x_2, i) \rangle_{\beta, \mu} - \langle \partial_{x_1} \psi^*(t_1, x_1, i) \psi(t_2, x_2, i) \rangle_{\beta, \mu}] = \frac{e}{m} \int_0^\infty \frac{dk}{2\pi} k \sum_{j=1}^n [\delta_{ij} - |\mathbb{S}_{ij}(k)|^2] d_j(k), \quad (3.44)$$

where $\mathbb{S}(k)$ is given by (3.21) and covers all point-like interactions leading to a self-adjoint Schrödinger Hamiltonian on the star graph. The current $J_i(\beta, \mu)$ is t -independent, implying that $\Omega_{\beta, \mu}$ are indeed steady states. We emphasize that (3.44) is precisely the Büttiker multi-channel generalization [45] of the Landauer expression [44] for the steady current. It is also worth stressing that in our context (3.44) is an exact formula, which does not rely on linear response theory. These remarkable features of the states $\Omega_{\beta, \mu}$ confirm their physical relevance and suggest to call them L-B states.

Let us summarize some of the basic properties of the steady current $J_i(\beta, \mu)$. First of all $J_i(\beta, \mu)$ is homogeneous (x -independent) and, because of unitarity (2.4), satisfies

$$\sum_{i=1}^n J_i(\beta, \mu) = 0. \quad (3.45)$$

This is the manifestation of the operator Kirchhoff rule (3.36) and represents a non-trivial check on (3.44). Moreover, there are two particular cases in which the system is in equilibrium and the current (3.44) must therefore vanish. The first one is when all thermal reservoirs are equivalent ($\beta_1 = \beta_2 = \dots = \beta_n$ and $\mu_1 = \mu_2 = \dots = \mu_n$). In fact, (2.4) implies in this case

$$J_i(\beta, \mu) = 0. \quad (3.46)$$

Another possibility to be in equilibrium is when all transmission coefficients vanish and the leads are therefore isolated. In this case $\mathbb{S}(k)$ is diagonal,

$$\mathbb{S}_{ij}(k) = \delta_{ij} e^{i\phi_j(k)}, \quad \phi_j(k) \in \mathbb{R} \quad (3.47)$$

which implies (3.46) as well.

In order to illustrate the role of the \mathbb{S} -matrices (3.21), it is instructive to consider (3.44) for $n = 2$. Using the general expression (3.26), one finds

$$J_1(\beta, \mu) = -J_2(\beta, \mu) = \frac{e}{m} [(\eta_1 - \eta_2) \sin(\theta)]^2 \int_0^\infty \frac{dk}{2\pi} \frac{k^3}{(k^2 + \eta_1^2)(k^2 + \eta_2^2)} [d_1(k) - d_2(k)], \quad (3.48)$$

where the sign-difference between J_1 and J_2 reflects the orientation of the leads. The k -integration in (3.48) can not be performed in a closed analytic form, but the integral is well-defined and can be computed numerically. The contour plots of J_1 for fixed e, θ and m , displayed in Fig. 2, give an idea about the behavior of (3.48) in the variables (β_1, β_2) , (μ_1, μ_2) and (η_1, η_2) . As usual, higher regions are shown in lighter shades. The plot on the left concerns J_1 in the plane (β_1, β_2) for fixed (μ_1, μ_2) and (η_1, η_2) . The plot in the middle illustrates the behavior of J_1 as a function of (μ_1, μ_2) , the variables (β_1, β_2) and (η_1, η_2) being fixed. Finally, the plot on the right shows the dependence on the \mathbb{S} -matrix variables (η_1, η_2) at fixed temperatures and chemical potentials.

The plots in Fig. 3 are obtained from those in Fig. 2 by fixing β_1, μ_1 and η_1 respectively. The sign change of J_1 in Fig. 3 indicates that varying β_2, μ_2 and η_2 one can invert the direction of the current.




Figure 2: Contour plots of the current J_1 in the plane (β_1, β_2) , (μ_1, μ_2) and (η_1, η_2) respectively.




Figure 3: Plots of J_1 as a function of β_2 , μ_2 and η_2 respectively, with all other variables fixed.

The expression (3.44) significantly simplifies at criticality, i.e. for *scale invariant* point-like interactions in the vertex of Γ . In this case the scattering matrix takes the form

$$\mathbb{S}(k) = \theta(k)\mathbb{U} + \theta(-k)\mathbb{U}^{-1}, \quad (3.49)$$

θ being the Heaviside step function. Plugging (3.49) in (3.44), one gets

$$J_i(\beta, \mu) = \frac{e}{2\pi} \sum_{j=1}^n (\delta_{ij} - |\mathbb{U}_{ij}|^2) \frac{1}{\beta_j} \ln(1 + e^{\beta_j \mu_j}). \quad (3.50)$$

The high and zero temperature limits of (3.50) are

$$J_i(0, \mu) \equiv \lim_{\beta_k = \beta \rightarrow 0} J_i(\beta, \mu) = \frac{e}{4\pi} \sum_{j=1}^n (\delta_{ij} - |\mathbb{U}_{ij}|^2) \mu_j, \quad (3.51)$$

$$J_i(\infty, \mu) \equiv \lim_{\beta_k = \beta \rightarrow \infty} J_i(\beta, \mu) = \frac{e}{2\pi} \sum_{j=1}^n (\delta_{ij} - |\mathbb{U}_{ij}|^2) \mu_j, \quad (3.52)$$

respectively and are related by

$$J_i(0, \mu) = \frac{1}{2} J_i(\infty, \mu). \quad (3.53)$$

By means of (3.44) one can derive the conductance tensor $\mathbb{G}_{ij}(\beta, \mu)$ defined by

$$J_i(\beta, \mu) = \sum_{j=1}^n \mathbb{G}_{ij}(\beta, \mu) V_j, \quad (3.54)$$

where V_j is the voltage applied at the edge E_j . It is well known [56] that $\mathbb{G}_{ij}(\beta, \mu)$ depends on the point on E_j where the voltage is applied. Assuming that this point is deeply in the reservoir with chemical potential μ_j , one has [56]

$$V_j = \frac{\mu_j}{e}. \quad (3.55)$$

Combining (3.44), (3.54) and (3.55) one obtains

$$\mathbb{G}_{ij}(\beta, \mu) = \frac{e^2}{m} \int_0^\infty \frac{dk}{2\pi} \frac{k}{\mu_j} [\delta_{ij} - |\mathbb{S}_{ij}(k)|^2] d_j(k), \quad (3.56)$$

which satisfies Kirchhoff's rule

$$\sum_{i=1}^n \mathbb{G}_{ij}(\beta, \mu) = 0 \quad (3.57)$$

as it should be.

Let us focus now on the charge density distribution $\langle j_t(t, x, i) \rangle_{\beta, \mu}$ in the state $\Omega_{\beta, \mu}$, which can be computed following the above procedure as well. As expected from the current conservation (3.35), the charge density

$$\varrho_i(\beta, \mu, x) \equiv \langle j_t(t, x, i) \rangle_{\beta, \mu} = e \int_0^\infty \frac{dk}{2\pi} \left\{ [\mathbb{S}_{ii}(k) e^{-2ikx} + \bar{\mathbb{S}}_{ii}(k) e^{2ikx} + 1] d_i(k) + \sum_{j=1}^n |\mathbb{S}_{ij}(k)|^2 d_j(k) \right\}, \quad (3.58)$$

is time independent as well. There are however two essential novelties with respect to the current:

- (a) ϱ_i does not vanish at equilibrium;
- (b) ϱ_i depends on the position x .

Concerning point (a), we observe that at equilibrium (n isolated leads with $\mathbb{S}(k)$ defined by (3.47)) one has

$$\varrho_i^{\text{eq}}(\beta, \mu, x) = e \int_0^\infty \frac{dk}{2\pi} \{ [\mathbb{S}_{ii}(k) e^{-2ikx} + \bar{\mathbb{S}}_{ii}(k) e^{2ikx} + 2] d_i(k) \}. \quad (3.59)$$

Therefore, the non-equilibrium charge distribution is x -independent and is given by

$$\varrho_i^{\text{neq}}(\beta, \mu) \equiv \varrho_i^{\text{eq}}(\beta, \mu, x) - \varrho_i(\beta, \mu, x) = e \int_0^\infty \frac{dk}{2\pi} \sum_{j=1}^n [\delta_{ij} - |\mathbb{S}_{ij}(k)|^2] d_j(k). \quad (3.60)$$

The x -dependence, mentioned in point (b), is carried by

$$\varrho_i^{\text{osc}}(\beta, \mu; x) = e \int_0^\infty \frac{dk}{2\pi} [\mathbb{S}_{ii}(k)e^{-2ikx} + \bar{\mathbb{S}}_{ii}(k)e^{2ikx}] d_i(k) \quad (3.61)$$

which oscillates with the distance x from the vertex. The appearance of such Friedel-type oscillations [55] confirms once more that the junction behaves indeed as a point-like defect. Since the integration in (3.61) can not be performed in closed form, in order to get an idea about the oscillations, it is useful to consider the zero-temperature limit

$$\varrho_i^{\text{osc}}(\infty, \mu; x) \equiv \lim_{\beta_k = \beta \rightarrow \infty} \varrho_i^{\text{osc}}(\beta, \mu; x) = e \int_0^{\sqrt{2m\mu_i}} \frac{dk}{2\pi} [\mathbb{S}_{ii}(k)e^{-2ikx} + \bar{\mathbb{S}}_{ii}(k)e^{2ikx}]. \quad (3.62)$$

At criticality (3.49) and setting $\mathbb{U}_{ii} = \bar{\mathbb{U}}_{ii}$ for simplicity, one finds

$$\varrho_i^{\text{osc}}(\infty, \mu; x) = \frac{e\mathbb{U}_{ii}}{\pi x} \sin(2x\sqrt{2m\mu_i}), \quad (3.63)$$

which shows that the amplitude of the oscillations on the graph decays with the distance from the vertex like x^{-1} , which is a typical behavior in one space dimension.

3.3 Casimir effect away from equilibrium and heat flow

The structure of the energy density in the state $\Omega_{\beta, \mu}$ resembles very much that of the charge density (3.58). One has

$$\begin{aligned} \mathcal{E}_i(x; \beta, \mu) &\equiv \langle \theta_{tt}(t, x, i) \rangle_{\beta, \mu} = \\ &\frac{1}{2} \int_0^\infty \frac{dk}{2\pi} \omega(k) \left\{ [\mathbb{S}_{ii}(k)e^{-2ikx} + \bar{\mathbb{S}}_{ii}(k)e^{2ikx} + 1] d_i(k) + \sum_{j=1}^n |\mathbb{S}_{ij}(k)|^2 d_j(k) \right\}, \end{aligned} \quad (3.64)$$

This result confirms the presence of Friedel oscillations in the energy density as well. It is instructive to compare (3.64) to the equilibrium energy density

$$\mathcal{E}_i^{\text{eq}}(x; \beta, \mu) = \int_0^\infty \frac{dk}{2\pi} \omega(k) [\mathbb{S}_{ii}(k)e^{-2ikx} + \bar{\mathbb{S}}_{ii}(k)e^{2ikx} + 2] d_i(k), \quad (3.65)$$

associated to (3.47)). One finds

$$\mathcal{E}_i^{\text{eq}}(x; \beta, \mu) - \mathcal{E}_i(x; \beta, \mu) = \frac{1}{2} \int_0^\infty \frac{dk}{2\pi} \omega(k) \sum_{j=1}^n [\delta_{ij} - |\mathbb{S}_{ij}(k)|^2] d_j(k), \quad (3.66)$$

which gives the genuine non-equilibrium part of the energy density (3.64). The x -independent contribution to (3.64), namely

$$\varepsilon_i(\beta, \mu) = \frac{1}{2} \sum_{j=1}^n \int_0^\infty \frac{dk}{2\pi} \omega(k) [\delta_{ij} + |\mathbb{S}_{ij}(k)|^2] d_j(k) \quad (3.67)$$

is the Stefan-Boltzmann law in the present context. At criticality

$$\varepsilon_i(\beta, \mu) = -\frac{1}{4} \sqrt{\frac{m}{2\pi}} \sum_{j=1}^n (\delta_{ij} + |\mathbb{U}_{ij}|^2) \frac{1}{\beta_j^{\frac{3}{2}}} \text{Li}_{\frac{3}{2}}(-e^{\beta_j \mu_j}), \quad (3.68)$$

where Li_s is the polylogarithm function.

The counterpart of the L-B formula for the heat (energy) flow is

$$\langle \theta_{xt}(t, x, i) \rangle_{\beta, \mu} = \frac{1}{m} \int_0^\infty \frac{dk}{2\pi} k \omega(k) \sum_{j=1}^n [\delta_{ij} - |\mathbb{S}_{ij}(k)|^2] d_j(k) \equiv \mathcal{T}_i(\beta, \mu). \quad (3.69)$$

Apart from the additional $\omega(k)$ factor in the integrand of (3.69), the charge (3.44) and energy (3.69) flows have the same structure. For this reason $\mathcal{T}_i(\beta, \mu)$ shares with $J_i(\beta, \mu)$ the general properties listed after equation (3.44). In the scale invariant case the energy flow is

$$\begin{aligned} \mathcal{T}_i(\beta, \mu) &= \frac{1}{2m^2} \sum_{j=1}^n (\delta_{ij} - |\mathbb{U}_{ij}|^2) \int_0^\infty \frac{dk}{2\pi} \frac{k^3 e^{-\beta_j[\omega(k) - \mu_j]}}{1 + e^{-\beta_j[\omega(k) - \mu_j]}} = \\ &= \frac{1}{2\pi} \sum_{j=1}^n (|\mathbb{U}_{ij}|^2 - \delta_{ij}) \frac{1}{\beta_j^2} \text{Li}_2(-e^{\beta_j \mu_j}), \end{aligned} \quad (3.70)$$

which gives in the zero temperature limit

$$\mathcal{T}_i(\infty, \mu) \equiv \lim_{\beta_k = \beta \rightarrow \infty} = \frac{1}{4\pi} \sum_{j=1}^n (\delta_{ij} - |\mathbb{U}_{ij}|^2) \mu_j^2. \quad (3.71)$$

3.4 Noise

In this section we derive the *noise power* generated by the point like interactions in the Schrödinger junction. For this purpose we need [57]-[59] the two-point *connected* current-current correlator. After some algebra one finds

$$\begin{aligned} \langle j_x(t_1, x_1, i) j_x(t_2, x_2, j) \rangle_{\beta, \mu}^{\text{conn}} &\equiv \langle j_x(t_1, x_1, i) j_x(t_2, x_2, j) \rangle_{\beta, \mu} - \langle j_x(t_1, x_1, i) \rangle_{\beta, \mu} \langle j_x(t_2, x_2, j) \rangle_{\beta, \mu} = \\ &= -\frac{e^2}{4m^2} \int_{-\infty}^0 \frac{dk_1}{2\pi} \int_{-\infty}^0 \frac{dk_2}{2\pi} e^{it_{12}[\omega(k_1) - \omega(k_2)]} \sum_{l, m=1}^n d_l(k_1) c_m(k_2) \\ &\times \left\{ \chi_{li}^*(k_1; x_1) [\partial_x \chi_{im}](k_2; x_1) - [\partial_x \chi_{li}^*](k_1; x_1) \chi_{im}(k_2; x_1) \right\} \\ &\times \left\{ \chi_{mj}^*(k_2; x_2) [\partial_x \chi_{jl}](k_1; x_2) - [\partial_x \chi_{mj}^*](k_2; x_2) \chi_{jl}(k_1; x_2) \right\}, \end{aligned} \quad (3.72)$$

where χ and χ^* are given by (3.38). Using the time translation invariance of (3.72), the noise power is defined [59] by

$$P_{ij}(\beta, \mu; x_1, x_2; \omega) \equiv \int_{-\infty}^{\infty} dt e^{i\omega t} \langle j_x(t, x_1, i) j_x(0, x_2, j) \rangle_{\beta, \mu}^{\text{conn}} \quad (3.73)$$

The zero-frequency limit (*zero-frequency noise power*)

$$P_{ij}(\beta, \mu) \equiv \lim_{\omega \rightarrow 0^+} P_{ij}(\beta, \mu; x_1, x_2; \omega) \quad (3.74)$$

turns out to be $x_{1,2}$ -independent and is given by:

$$P_{ij}(\beta, \mu) = \frac{e^2}{m} \int_0^{\infty} \frac{dk}{2\pi} k \left[\delta_{ij} d_i(k) c_i(k) - |\mathbb{S}_{ij}(k)|^2 d_j(k) c_j(k) - |\mathbb{S}_{ji}(k)|^2 d_i(k) c_i(k) + \sum_{l,m=1}^n \mathbb{S}_{il}(k) c_l(k) \bar{\mathbb{S}}_{jl}(k) \mathbb{S}_{jm}(k) d_m(k) \bar{\mathbb{S}}_{im}(k) \right]. \quad (3.75)$$

It is instructive to summarize at this point the general features of (3.75):

(i) $P_{ij}(\beta, \mu)$ is symmetric in i and j . The first three terms of the integrand are manifestly symmetric. Concerning the last term, using the identity $c_i(k) = 1 - d_i(k)$ one gets

$$\begin{aligned} & \sum_{l,m=1}^n \mathbb{S}_{il}(k) c_l(k) \bar{\mathbb{S}}_{jl}(k) \mathbb{S}_{jm}(k) d_m(k) \bar{\mathbb{S}}_{im}(k) = \\ & \delta_{ij} \sum_{m=1}^n \mathbb{S}_{im}(k) d_m(k) \bar{\mathbb{S}}_{im}(k) \mp \sum_{l,m=1}^n \mathbb{S}_{il}(k) d_l(k) \bar{\mathbb{S}}_{jl}(k) \mathbb{S}_{jm}(k) d_m(k) \bar{\mathbb{S}}_{im}(k), \end{aligned} \quad (3.76)$$

which is symmetric as well. One can therefore rewrite $P_{ij}(\beta, \mu)$ in the following manifestly symmetric form:

$$P_{ij}(\beta, \mu) = \frac{e^2}{m} \int_0^{\infty} \frac{dk}{2\pi} k \left\{ \delta_{ij} d_i(k) c_i(k) - |\mathbb{S}_{ij}(k)|^2 d_j(k) c_j(k) - |\mathbb{S}_{ji}(k)|^2 d_i(k) c_i(k) + \frac{1}{2} \sum_{l,m=1}^n \mathbb{S}_{il}(k) \bar{\mathbb{S}}_{jl}(k) \mathbb{S}_{jm}(k) \bar{\mathbb{S}}_{im}(k) [c_l(k) d_m(k) + c_m(k) d_l(k)] \right\}. \quad (3.77)$$

(ii) The last identity implies also that $P_{ij}(\beta, \mu)$ is real;

(iii) As expected, $P_{ij}(\beta, \mu)$ satisfies the Kirchhoff rule

$$\sum_{i=1}^n P_{ij}(\beta, \mu) = \sum_{j=1}^n P_{ij}(\beta, \mu) = 0, \quad (3.78)$$

which provides an useful check. One has actually

$$\sum_{i=1}^n P_{ij}(\beta, \mu; 0, x_2; \omega) = \sum_{j=1}^n P_{ij}(\beta, \mu; x_1, 0; \omega) = 0, \quad (3.79)$$

at any frequency ω .

(iv) All noise components $P_{ij}(\beta, \mu)$ vanish for isolated leads (3.47).

Let us discuss now the behavior of the noise, starting with the case $n = 2$. Combining (3.26) with (3.77), we find

$$P_{11}(\beta, \mu) = [e(\eta_1 - \eta_2) \sin(\theta)]^2 \frac{1}{m} \int_0^\infty \frac{dk}{2\pi} \frac{k^3}{(k^2 + \eta_1^2)(k^2 + \eta_2^2)} \left\{ d_1(k) + d_2(k) - 2d_1(k)d_2(k) - \frac{k^2[(\eta_1 - \eta_2) \sin(\theta)]^2}{(k^2 + \eta_1^2)(k^2 + \eta_2^2)} [d_1(k) - d_2(k)]^2 \right\}. \quad (3.80)$$

For $\eta_1 = \eta_2$ and/or $\theta = 0$ the leads are isolated (see (3.26)) and the noise vanishes according to point (iv) above. Like for the steady current, we report some contour plots, showing the complicated dependence of the noise on the parameters (β_1, β_2) , (μ_1, μ_2) and (η_1, η_2) for fixed e, θ and m . Fig. 4 illustrates the behavior of $P_{11}(\beta, \mu)$ in each pair of these variables, the remaining two being fixed. The left plot is the noise in the plane (β_1, β_2) . In the middle we display the noise as a function of the chemical potentials (μ_1, μ_2) . Finally, the right plot shows the dependence on the S-matrix variables (η_1, η_2) .




Figure 4: Contour plots of the noise P_{11} in the plane (β_1, β_2) , (μ_1, μ_2) and (η_1, η_2) respectively.

The plots in Fig. 5, obtained from those in Fig. 4 by fixing β_1 , μ_1 and η_1 respectively, confirm that the noise $P_{ij}(\beta, \mu)$ depends in a complicated way on β , μ and the S-matrix parameters. At criticality however, as expected on general grounds, the situation simplifies and one can push




Figure 5: Plots of P_{11} as a function of β_2 , μ_2 and η_2 respectively, with all other variables fixed.

further the analytic computation. In fact, inserting (3.49) in (3.77) one gets,

$$P_{ij}(\beta, \mu) = \frac{e^2}{m} \left\{ \delta_{ij} I_{ii}(\beta) - |\mathbb{U}_{ij}|^2 I_{jj}(\beta) - |\mathbb{U}_{ji}|^2 I_{ii}(\beta) + \right. \\ \left. + \frac{1}{2} \sum_{l,m=1}^n \bar{\mathbb{U}}_{il} \mathbb{U}_{jl} \bar{\mathbb{U}}_{jm} \mathbb{U}_{im} [I_{lm}(\beta) + I_{ml}(\beta)] \right\}, \quad (3.81)$$

where

$$I_{ij}(\beta) \equiv \int_0^\infty \frac{dk}{2\pi} k d_i(k) c_j(k). \quad (3.82)$$

For equal temperatures $\beta_i = \beta_j = \beta$ the integration in (3.82) can be performed explicitly and one finds

$$I_{ij}(\beta) = \begin{cases} \frac{m}{2\pi\beta} \frac{e^{\beta\mu}}{1+e^{\beta\mu}} & \text{if } \mu_i = \mu_j \equiv \mu, \\ \frac{m}{2\pi\beta} \frac{e^{\beta\mu_i}}{e^{\beta\mu_j} - e^{\beta\mu_i}} \ln \left(\frac{1+e^{\beta\mu_j}}{1+e^{\beta\mu_i}} \right) & \text{if } \mu_i \neq \mu_j. \end{cases} \quad (3.83)$$

Therefore, in the case $\beta_1 = \beta_2 = \dots = \beta_n = \beta$ with generic chemical potentials $\mu_i \geq 0$ one obtains

$$P_{ij}(\beta, \mu) = \frac{e^2}{2\pi\beta} \left[\delta_{ij} \frac{e^{\beta\mu_i}}{1+e^{\beta\mu_i}} - |\mathbb{U}_{ij}|^2 \frac{e^{\beta\mu_j}}{1+e^{\beta\mu_j}} - |\mathbb{U}_{ji}|^2 \frac{e^{\beta\mu_i}}{1+e^{\beta\mu_i}} + \right. \\ \left. + \sum_{l=1}^n |\mathbb{U}_{il}|^2 |\mathbb{U}_{jl}|^2 \frac{e^{\beta\mu_l}}{1+e^{\beta\mu_l}} + \frac{1}{2} \sum_{\substack{l,m=1 \\ l \neq m}}^n \bar{\mathbb{U}}_{il} \mathbb{U}_{jl} \bar{\mathbb{U}}_{jm} \mathbb{U}_{im} \frac{e^{\beta\mu_l} + e^{\beta\mu_m}}{e^{\beta\mu_l} - e^{\beta\mu_m}} \ln \left(\frac{1+e^{\beta\mu_l}}{1+e^{\beta\mu_m}} \right) \right]. \quad (3.84)$$

It is instructive to consider at this stage the two limits leading to the shot and thermal noise. For deriving the shot noise, we need the $\beta \rightarrow \infty$ limit of the integrals (3.83), which are

$$\lim_{\beta \rightarrow \infty} I_{ij}(\beta) = \begin{cases} 0 & \text{if } \mu_i \leq \mu_j, \\ \frac{m}{2\pi} (\mu_i - \mu_j) & \text{if } \mu_i > \mu_j. \end{cases} \quad (3.85)$$

Therefore, in the scale invariant case (3.49) the shot noise is

$$P_{ij}(\mu) \equiv \lim_{\beta \rightarrow \infty} P_{ij}(\beta, \mu) = \frac{e^2}{4\pi} \sum_{\substack{l,m=1 \\ l \neq m}}^n \bar{\mathbb{U}}_{il} \mathbb{U}_{jl} \bar{\mathbb{U}}_{jm} \mathbb{U}_{im} |\mu_l - \mu_m|, \quad (3.86)$$

which exhibits the standard behavior [57]-[59] in terms of $|\mu_i - \mu_j|$.

In order to compute the thermal noise, we consider (3.84) for $\mu_1 = \mu_2 = \dots = \mu_n = \mu$. One has

$$P_{ij}(\beta, \mu) = \frac{e^2}{2\pi\beta} \frac{e^{\beta\mu}}{1 + e^{\beta\mu}} [2\delta_{ij} - |\mathbb{U}_{ij}|^2 - |\mathbb{U}_{ji}|^2]. \quad (3.87)$$

In the purely thermal case ($\mu \rightarrow 0$) one finds

$$P_{ij}(\beta) = \frac{e^2}{2\pi\beta} [2\delta_{ij} - |\mathbb{U}_{ij}|^2 - |\mathbb{U}_{ji}|^2] \sim T, \quad (3.88)$$

which is the well-known Johnson-Nyquist formula.

A remarkable feature of the thermal noise is that away from criticality the point-like interactions at the vertex can modify the linear behavior (3.88) for large T . Let us consider indeed (3.80) for $\beta_1 = \beta_2 = \beta$ and $\mu_1 = \mu_2 = 0$, namely

$$P_{11}(\beta) = 2 [e(\eta_1 - \eta_2) \sin(\theta)]^2 \frac{1}{m} \int_0^\infty \frac{dk}{2\pi} \frac{k^3 e^{-\beta \frac{k^2}{2m}}}{(k^2 + \eta_1^2)(k^2 + \eta_2^2) \left(1 + e^{-\beta \frac{k^2}{2m}}\right)^2}. \quad (3.89)$$

Assuming for simplicity $\eta_2 = 0$ and introducing the variables $\eta = \eta_1$ and $\xi = e^{-\beta \frac{k^2}{2m}}$, one gets

$$P_{11}(\beta) = [e\eta \sin(\theta)]^2 \frac{1}{2\pi m} \int_0^1 d\xi \frac{1}{(1 + \xi)^2 \left[\frac{\beta\eta^2}{2m} - \ln(\xi)\right]}. \quad (3.90)$$

Since the k -integration in (3.26) can not be performed exactly, in order to estimate the temperature dependence one can use the inequalities

$$[e\eta \sin(\theta)]^2 \frac{1}{4\pi m} I\left(\frac{\beta\eta^2}{2m}\right) \leq P_{11}(\beta) \leq [e\eta \sin(\theta)]^2 \frac{1}{2\pi m} I\left(\frac{\beta\eta^2}{2m}\right), \quad (3.91)$$

where [60]

$$I(a) = \int_0^1 d\xi \frac{1}{[a - \ln(\xi)]} = -e^a \text{Ei}(-a), \quad (3.92)$$

Ei being the exponential integral function. In this way one finds

$$P_{11}(\beta) \sim \begin{cases} T & \text{for } T \rightarrow 0, \\ \ln(T) & \text{for } T \rightarrow \infty, \end{cases} \quad (3.93)$$

which shows that the k -dependence of the \mathbb{S} -matrix indeed modifies the Johnson-Nyquist behavior at high temperatures. The milder logarithmic divergence for large T provides an attractive experimental signature.

3.5 External electromagnetic field

In the above considerations the interaction was localized in the junction. We extend here the framework to the more realistic physical situation of a junction in a three-dimensional ambient space with a classical static magnetic field, interacting with the Schrödinger excitations along the leads. The graph Γ , modeling the junction, is embedded in \mathbb{R}^3 , equipped with a Cartesian coordinate system whose origin O coincides with the vertex V of Γ . The direction of each edge $E_i \subset \mathbb{R}^3$ is determined by the unit vector $\mathbf{e}^{(i)}$. At any point $P \in \mathbb{R}^3$ the magnetic field $\mathbf{B}(P) = \text{rot}[\mathbf{A}(P)]$ is generated by the potential $\mathbf{A}(P)$. The minimal coupling of the Schrödinger field φ with \mathbf{A} gives the following equation of motion

$$\left[i\partial_t - \frac{1}{2m} (i\partial_x - eA_x(x, i)) (i\partial_x - eA_x(x, i)) \right] \varphi(t, x, i) = 0, \quad (3.94)$$

$A_x(x, i)$ being the projection

$$A_x(x, i) = \mathbf{e}^{(i)} \cdot \mathbf{A}(P), \quad P \equiv (x, i) \in \Gamma \subset \mathbb{R}^3, \quad (3.95)$$

of the potential \mathbf{A} along the edge E_i . All self-adjoint extensions of the relative Hamiltonian are now parametrized by the boundary conditions [61]

$$\lim_{x \rightarrow 0^+} \sum_{j=1}^n [\lambda(\mathbb{I} - \mathbb{U})_{ij} - (\mathbb{I} + \mathbb{U})_{ij} (i\partial_x - A_x(x, j))] \varphi(t, x, j) = 0. \quad (3.96)$$

The conserved electric current is

$$j_x(t, x, i) = i \frac{e}{2m} [\varphi^* (\partial_x \varphi) - (\partial_x \varphi^*) \varphi] (t, x, i) - \frac{e}{m} A_x(x, i) (\varphi^* \varphi) (t, x, i). \quad (3.97)$$

It is easy to show now that the solution of the problem (3.94,3.96) can be reduced to that described in section 3.1. Indeed, let us introduce

$$\psi(t, x, i) = e^{-ie\alpha(x, i)} \varphi(t, x, i), \quad \alpha(x, i) = \int_x^\infty dy A_y(y, i), \quad (3.98)$$

where we assumed that $A_x(x, i)$ are integrable on the half line. Notice that ψ and φ have the same behavior for $x \rightarrow \infty$. Moreover, ψ satisfies (3.17) and in terms of ψ the current (3.97) takes precisely the form (3.31). The interaction is totally absorbed in the boundary condition for ψ following from (3.96,3.98). One has

$$\lim_{x \rightarrow 0^+} \sum_{j=1}^n [\lambda(\mathbb{I} - \mathbb{U}(\mathbf{A}))_{ij} - i(\mathbb{I} + \mathbb{U}(\mathbf{A}))_{ij} \partial_x] \psi(t, x, j) = 0, \quad (3.99)$$

where

$$\mathbb{U}_{ij}(\mathbf{A}) = e^{-ie\alpha_i} \mathbb{U}_{ij} e^{ie\alpha_j}, \quad \alpha_i = \alpha(0, i). \quad (3.100)$$

Combining (3.21) and (3.100) one concludes that the substitution

$$\mathbb{S}_{ij}(k) \longmapsto e^{-ie\alpha_i} \mathbb{S}_{ij}(k) e^{ie\alpha_j} \quad (3.101)$$

extends all the results of this section to the case of a junction minimally coupled to a time-independent ambient magnetic field. We stress that the correlation functions of fields localized in different edges (see e.g. (3.41,3.42,3.72)) are sensitive to the transformation (3.101). In particular, the field \mathbf{A} has a non-trivial impact on the noise power at frequency $\omega \neq 0$. We will analyze this issue in more details elsewhere.

3.6 Remarks

Let us discuss first the role of possible bound states of $\mathbb{S}(k)$, which have been excluded in the above considerations by assuming (3.27). It has been shown in previous work ([53], [28], [31]) that the bound states generate new quantum degrees of freedom, which have a non-trivial contribution to the correlation function (3.41). The key point is that this contribution depends on the space-time coordinates only through the combinations t_{12} and \tilde{x}_{12} . According to (3.31, 3.32), the charge (3.44) and energy (3.70) flows are therefore not affected by the presence of bound states. The relative densities however get [53] nontrivial bound state contributions.

One can investigate along the above lines also the Schrödinger equation (3.17) with Bose statistics. The final results in this case obviously follow from equations (3.41,3.44,3.58-3.70,3.77) by substituting the Fermi distribution with the Bose distribution $d_i^-(k)$. Concerning the noise, in the scale invariant case one gets for bosons

$$P_{ij}^-(\beta, \mu) = \frac{e^2}{2\pi\beta} \left[\delta_{ij} \frac{e^{\beta\mu_i}}{1 - e^{\beta\mu_i}} - |\mathbb{U}_{ij}|^2 \frac{e^{\beta\mu_j}}{1 - e^{\beta\mu_j}} - |\mathbb{U}_{ji}|^2 \frac{e^{\beta\mu_i}}{1 - e^{\beta\mu_i}} + \sum_{l=1}^n |\mathbb{U}_{il}|^2 |\mathbb{U}_{jl}|^2 \frac{e^{\beta\mu_l}}{1 - e^{\beta\mu_l}} + \frac{1}{2} \sum_{\substack{l,m=1 \\ l \neq m}}^n \overline{\mathbb{U}}_{il} \mathbb{U}_{jl} \overline{\mathbb{U}}_{jm} \mathbb{U}_{im} \frac{e^{\beta\mu_l} + e^{\beta\mu_m}}{e^{\beta\mu_l} - e^{\beta\mu_m}} \ln \left(\frac{1 - e^{\beta\mu_m}}{1 - e^{\beta\mu_l}} \right) \right], \quad (3.102)$$

which, compared to (3.84), shows how the zero frequency noise power depends on the statistics. For instance, in the shot noise limit one obtains

$$P_{ij}^-(\mu) \equiv \lim_{\beta \rightarrow \infty} P_{ij}^-(\beta, \mu) = -\frac{e^2}{4\pi} \sum_{\substack{l,m=1 \\ l \neq m}}^n \overline{\mathbb{U}}_{il} \mathbb{U}_{jl} \overline{\mathbb{U}}_{jm} \mathbb{U}_{im} |\mu_l - \mu_m|, \quad (3.103)$$

which has the magnitude of the fermionic shot noise (3.86) but the opposite sign [59].

4 The Dirac junction

4.1 Preliminaries

The massless Dirac equation on the star graph Γ is

$$(\gamma_t \partial_t - \gamma_x \partial_x) \psi(t, x, i) = 0, \quad x > 0, \quad (4.104)$$

where

$$\psi(t, x, i) = \begin{pmatrix} \psi_1(t, x, i) \\ \psi_2(t, x, i) \end{pmatrix}, \quad \gamma_t = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \gamma_x = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (4.105)$$

We assume that ψ_α satisfy the conventional equal-time anti-commutation relations. The boundary conditions which define all self-adjoint extensions of the bulk Hamiltonian $i\gamma_t\gamma_x\partial_x$ are [62], [63]

$$\psi_1(t, 0, i) = \sum_{j=1}^n \mathbb{U}_{ij} \psi_2(t, 0, j), \quad (4.106)$$

where \mathbb{U} is any unitary $n \times n$ matrix. In physical terms \mathbb{U} parametrizes all point-like interactions for which $i\gamma_t\gamma_x\partial_x$ extends to a self-adjoint Hamiltonian to the whole Γ . Observing that both the equation of motion (4.104) and the boundary condition (4.106) preserve scale invariance, it is not surprising that the scattering matrix corresponding to these interactions is simply (3.49).

The Dirac field ψ is complex, has a relativistic dispersion relation

$$\omega(k) = |k|, \quad (4.107)$$

and describes therefore both particle and antiparticle excitations. For quantizing (4.104, 4.106) we need for this reason two copies of reflection-transmission algebras [63]. The first one \mathcal{A}_+ is generated by $\{a_i(k), a_i^*(k) : k \in \mathbb{R}\}$ and $\mathbb{S}(k)$ given by (3.49). We denote the second one by \mathcal{A}_+^t because its generators $\{b_i(k), b_i^*(k) : k \in \mathbb{R}\}$ obey the anti-commutation relations (2.1, 2.2) with the *transpose* scattering matrix $\mathbb{S}^t(k)$. Besides (2.3), one has therefore

$$b_i(k) = \sum_{j=1}^n b_j(-k) \mathbb{S}_{ji}(k), \quad b_i^*(k) = \sum_{j=1}^n \mathbb{S}_{ij}(-k) b_j^*(-k). \quad (4.108)$$

In what follows we use the convention according to which $\{a_i(k), a_i^*(k)\}$ and $\{b_i(k), b_i^*(k)\}$ annihilate/create respectively antiparticles and particles. The solution of (4.104, 4.106) in this basis is

$$\psi_1(t, x, i) = \int_0^\infty \frac{dk}{2\pi} [a_i(k) e^{-ik(t-x)} + b_i^*(-k) e^{ik(t-x)}], \quad (4.109)$$

$$\psi_2(t, x, i) = \int_0^\infty \frac{dk}{2\pi} [a_i(-k) e^{-ik(t+x)} + b_i^*(k) e^{ik(t+x)}]. \quad (4.110)$$

In the Dirac junction the anti-unitary operator T of time reversal and the unitary operator C of charge conjugation act as follows:

$$T\psi_1(t, x, i)T^{-1} = \eta_T \psi_2(-t, x, i), \quad T\psi_2(t, x, i)T^{-1} = \eta_T \psi_1(-t, x, i), \quad |\eta_T| = 1, \quad (4.111)$$

$$C\psi_1(t, x, i)C^{-1} = -\eta_C\psi_1^*(t, x, i), \quad C\psi_2(t, x, i)C^{-1} = \eta_C\psi_2^*(t, x, i), \quad |\eta_C| = 1. \quad (4.112)$$

Like in the Schrödinger case the boundary condition (4.106) is invariant under time reversal only if \mathbb{U} is symmetric (3.30). The condition for charge conjugation invariance is instead

$$\bar{\mathbb{U}} = -\mathbb{U}. \quad (4.113)$$

The violation of (3.30) and/or (4.113) leads to the breakdown of the corresponding symmetry by means of the boundary condition (4.106).

The electric current and energy-momentum tensor are

$$j_t(t, x, i) = -e : \psi^* \psi : (t, x, i), \quad (4.114)$$

$$j_x(t, x, i) = -e : \psi^* \sigma \psi : (t, x, i), \quad (4.115)$$

$$\theta_{tt}(t, x, i) = \frac{i}{2} : [\psi^* (\partial_t \psi) - (\partial_t \psi^*) \psi] : (t, x, i), \quad (4.116)$$

$$\theta_{xt}(t, x, i) = \frac{i}{2} : [\psi^* \sigma (\partial_t \psi) - (\partial_t \psi^*) \sigma \psi] : (t, x, i), \quad (4.117)$$

where $:\dots:$ denotes the normal product in \mathcal{A}_+ and A_+^t and

$$\sigma = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (4.118)$$

According to our convention for particles and antiparticles, the incoming asymptotic sub-algebra $\mathcal{D}_+^{\text{in}}$ is generated by $\{a_i(k), a_i^*(k), b_i(-k), b_i^*(-k) : k > 0\}$. The edge Hamiltonians and the asymptotic charge operators of particles and antiparticles are

$$h_i = \int_{-\infty}^0 \frac{dk}{2\pi} |k| [a_i^*(-k)a_i(-k) + b_i^*(k)b_i(k)], \quad (4.119)$$

$$q_i = \int_{-\infty}^0 \frac{dk}{2\pi} b_i^*(k)b_i(k), \quad \tilde{q}_i = - \int_{-\infty}^0 \frac{dk}{2\pi} a_i^*(-k)a_i(-k), \quad (4.120)$$

respectively. We associate with (4.120) the chemical potentials μ_i and $\tilde{\mu}_i$. Now, following the general strategy explained in section 2, we set

$$K = \sum_{i=1}^n \beta_i (h_i - \mu_i q_i - \tilde{\mu}_i \tilde{q}_i) \quad (4.121)$$

and define first the steady state $\Omega_{\beta, \mu, \tilde{\mu}}$ on the sub-algebra $\mathcal{D}_+^{\text{in}}$ by means of (2.9). Employing (2.3, 4.108), we extend after that the state $\Omega_{\beta, \mu, \tilde{\mu}}$ to the whole algebra generated by \mathcal{A}_+ and \mathcal{A}_+^t . In this way one gets

$$\begin{aligned} \langle a_j^*(p)a_i(k) \rangle_{\beta, \mu, \tilde{\mu}} = 2\pi \left\{ \left[\theta(k) \tilde{f}_i(k) \delta_{ij} + \theta(-k) \sum_{l=1}^n \mathbb{U}_{il}^* \tilde{f}_l(-k) \mathbb{U}_{lj} \right] \delta(k-p) \right. \\ \left. + \left[\theta(k) \tilde{f}_i(k) \mathbb{U}_{ij} + \theta(-k) \mathbb{U}_{ij}^* \tilde{f}_j(-k) \right] \delta(k+p) \right\}, \quad (4.122) \end{aligned}$$

$$\begin{aligned} \langle b_j^*(p)b_i(k) \rangle_{\beta, \mu, \tilde{\mu}} &= 2\pi \left\{ \left[\theta(-k)f_i(k)\delta_{ij} + \theta(k) \sum_{l=1}^n \mathbb{U}_{jl}^* f_l(-k) \mathbb{U}_{li} \right] \delta(k-p) \right. \\ &\quad \left. + \left[\theta(-k)\mathbb{U}_{ji}^* f_i(k) + \theta(k)f_j(-k)\mathbb{U}_{ji} \right] \delta(k+p) \right\}, \end{aligned} \quad (4.123)$$

where

$$\tilde{f}_i(k) = \frac{e^{-\beta_i(|k|+\tilde{\mu}_i)}}{1 + e^{-\beta_i(|k|+\tilde{\mu}_i)}}, \quad f_i(k) = \frac{e^{-\beta_i(|k|-\mu_i)}}{1 + e^{-\beta_i(|k|-\mu_i)}} \quad (4.124)$$

are the Dirac distributions for antiparticles and particles respectively.

Let us discuss finally the behavior of $\Omega_{\beta, \mu, \tilde{\mu}}$ under charge conjugation. One easily verifies that $\Omega_{\beta, \mu, \tilde{\mu}}$ is invariant under charge conjugation, namely $C\Omega_{\beta, \mu, \tilde{\mu}} = \Omega_{\beta, \mu, \tilde{\mu}}$, provided that both conditions

$$\mu_i = -\tilde{\mu}_i \quad (4.125)$$

and (4.113) hold.

4.2 Transport properties

We are ready at this point to derive the steady currents in the state $\Omega_{\beta, \mu, \tilde{\mu}}$. A computation, analogous to that performed in section 3, gives

$$\begin{aligned} J_i(\beta, \mu, \tilde{\mu}) &\equiv \langle j_x(t, x, i) \rangle_{\beta, \mu, \tilde{\mu}} = e \sum_{j=1}^n (\delta_{ij} - |\mathbb{U}_{ij}|^2) \int_0^\infty \frac{dk}{2\pi} \left[f_j(-k) - \tilde{f}_j(k) \right] = \\ &\quad \frac{e}{2\pi} \sum_{j=1}^n (\delta_{ij} - |\mathbb{U}_{ij}|^2) \frac{1}{\beta_j} \ln \left(\frac{1 + e^{\beta_j \mu_j}}{1 + e^{-\beta_j \tilde{\mu}_j}} \right), \end{aligned} \quad (4.126)$$

which satisfies Kirchhoff's rule and vanishes at equilibrium exactly like the Schrödinger steady current. A new feature of the Dirac steady current is the presence of particle and antiparticle contributions, captured respectively by the numerator and the denominator in the fraction under the logarithm in (4.126).

Concerning the dependence of (4.126) on the chemical potentials, some particular cases are worth mentioning. We first observe that if charge conjugation is preserved, the steady current (4.126) vanishes because of (4.125). This is due to a cancellation between the particle and antiparticle contributions. If instead all $\tilde{\mu}_i = \mu_i$, the current (4.126) is temperature independent (in spite of the fact that the junction is in contact with heat reservoirs with different temperatures β_j) and takes the simple form

$$J_i(\beta, \mu, \mu) = \frac{e}{2\pi} \sum_{j=1}^n (\delta_{ij} - |\mathbb{U}_{ij}|^2) \mu_j. \quad (4.127)$$

In the case $\tilde{\mu}_i = 0$ the steady current (4.126) coincides with that of the Schrödinger junction (3.50) with scale invariant boundary conditions.

Let us focus now on the temperature dependence of (4.126). At high temperatures one has

$$\lim_{\beta_k=\beta\rightarrow 0} J_i(\beta, \mu, \tilde{\mu}) = \frac{e}{4\pi} \sum_{j=1}^n (\delta_{ij} - |\mathbb{U}_{ij}|^2) (\mu_j + \tilde{\mu}_j), \quad (4.128)$$

whereas at zero temperature

$$\lim_{\beta_k=\beta\rightarrow +\infty} J_i(\beta, \mu, \tilde{\mu}) = \frac{e}{2\pi} \sum_{j=1}^n (\delta_{ij} - |\mathbb{U}_{ij}|^2) [\mu_j \theta(\mu_j) + \tilde{\mu}_j \theta(-\tilde{\mu}_j)]. \quad (4.129)$$

The conductance tensor corresponding to (4.126) is

$$\mathbb{G}_{ij}(\beta, \mu) = \frac{e^2}{2\pi} (\delta_{ij} - |\mathbb{U}_{ij}|^2) \frac{1}{\beta_j \mu_j} \ln \left(\frac{1 + e^{\beta_j \mu_j}}{1 + e^{-\beta_j \tilde{\mu}_j}} \right) \quad (4.130)$$

and is not symmetric in general.

For the energy flow one obtains

$$\begin{aligned} \mathcal{T}_i(\beta, \mu, \tilde{\mu}) &= \langle \theta_{xt}(t, x, i) \rangle_{\beta, \mu, \tilde{\mu}} = \sum_{j=1}^n (\delta_{ij} - |\mathbb{U}_{ij}|^2) \int_0^\infty \frac{dk}{2\pi} [\tilde{f}_j(k) + f_j(-k)] = \\ &= \frac{1}{2\pi} \sum_{j=1}^n (|\mathbb{U}_{ij}|^2 - \delta_{ij}) \frac{1}{\beta_j^2} [\text{Li}_2(-e^{\beta_j \mu_j}) + \text{Li}_2(-e^{-\beta_j \tilde{\mu}_j})]. \end{aligned} \quad (4.131)$$

One can easily verify that the charge and energy densities are obtained by the replacement $(|\mathbb{U}_{ij}|^2 - \delta_{ij}) \mapsto (|\mathbb{U}_{ij}|^2 + \delta_{ij})$ in (4.126) and (4.131) respectively. Notice that these quantities are x -independent and therefore do not present Friedel oscillations. The reason is that both the dynamics (4.104) and the boundary conditions (4.106) are scale invariant for massless fermions.

4.3 Noise

For conciseness we report directly the zero-frequency noise power in terms of the matrix \mathbb{U} appearing in the boundary condition (4.106) and the distributions (4.124). One has

$$\begin{aligned} P_{ij}(\beta, \mu, \tilde{\mu}) &= e^2 \int_0^\infty \frac{dk}{2\pi} \left\{ \delta_{ij} F_{ii}(k) - |\mathbb{U}_{ij}|^2 F_{ii}(k) - |\mathbb{U}_{ji}|^2 F_{jj}(k) + \right. \\ &\quad \left. + \frac{1}{2} \sum_{l,m=1}^n \mathbb{U}_{li} \bar{\mathbb{U}}_{lj} \mathbb{U}_{mj} \bar{\mathbb{U}}_{mi} [F_{lm}(k) + F_{ml}(k)] \right\}, \end{aligned} \quad (4.132)$$

with

$$F_{ij}(k) = f_i(k)[1 - f_j(k)] + \tilde{f}_i(k)[1 - \tilde{f}_j(k)]. \quad (4.133)$$

If all the temperatures are equal ($\beta_i = \beta$), the k -integration in the right hand side of (4.132) can be performed exactly and gives

$$P_{ij}(\beta, \mu, \tilde{\mu}) = \frac{e^2}{2\pi\beta} \left\{ (\delta_{ij} - |\mathbb{U}_{ij}|^2) \left[\frac{e^{\beta\mu_i}}{1 + e^{\beta\mu_i}} + \frac{e^{-\beta\tilde{\mu}_i}}{1 + e^{-\beta\tilde{\mu}_i}} \right] - |\mathbb{U}_{ji}|^2 \left[\frac{e^{\beta\mu_j}}{1 + e^{\beta\mu_j}} + \frac{e^{-\beta\tilde{\mu}_j}}{1 + e^{-\beta\tilde{\mu}_j}} \right] + \frac{1}{2} \sum_{l,m=1}^n \mathbb{U}_{li} \bar{\mathbb{U}}_{lj} \mathbb{U}_{mj} \bar{\mathbb{U}}_{mi} \left[\frac{e^{\beta\mu_l} + e^{\beta\mu_m}}{e^{\beta\mu_l} - e^{\beta\mu_m}} \ln \left(\frac{1 + e^{\beta\mu_l}}{1 + e^{\beta\mu_m}} \right) + \frac{e^{-\beta\tilde{\mu}_l} + e^{-\beta\tilde{\mu}_m}}{e^{-\beta\tilde{\mu}_l} - e^{-\beta\tilde{\mu}_m}} \ln \left(\frac{1 + e^{-\beta\tilde{\mu}_l}}{1 + e^{-\beta\tilde{\mu}_m}} \right) \right] \right\}. \quad (4.134)$$

For the purely thermal noise one gets therefore

$$P_{ij}(\beta, 0, 0) = \frac{e^2}{2\pi\beta} (2\delta_{ij} - |\mathbb{U}_{ij}|^2 - |\mathbb{U}_{ji}|^2), \quad (4.135)$$

which coincides precisely with the result (3.88) for the Schrödinger junction at criticality.

4.4 Remarks

In spite of the different dispersion relations, the general structure of the steady currents and the noise in the Schrödinger and Dirac cases are quite similar. A characteristic feature of the Dirac case is the possibility to introduce the independent chemical potentials μ_i and $\tilde{\mu}_i$, associated with particles and antiparticles. This fact has elementary but important consequences. If $\mu_i = -\tilde{\mu}_i$ the particle and antiparticle contributions cancel each other in the electric steady current (4.126), but sum up in the heat current (4.131) and in the zero frequency noise (4.135).

Following the argument in section 3.5, the results about the Dirac junction have a straightforward generalization to the case when ψ is minimally coupled to a static classical electromagnetic field generated by the potential $(A_t(P), \mathbf{A}(P))$ in the ambient space.

5 Outlook and conclusions

In this paper we developed an algebraic method for constructing non-equilibrium steady states $\Omega_{\beta,\mu}$ on star graphs. Our approach is microscopic and our construction generalizes that of a Gibbs state over the algebra of canonical (anti)commutation relations. The Schrödinger and Dirac equations have been investigated in this framework. We considered in detail the case in which the interaction, driving the system away from equilibrium, is localized in the vertex of the graph. It turns out that the non-equilibrium dynamics, generated by such interactions, is exactly solvable. In fact, the $\Omega_{\beta,\mu}$ -expectation values of various observables (currents and charge densities) can be computed exactly, without resorting to any kind of approximation. We have shown in particular, that the expectation value of the electric current in the Schrödinger case reproduces precisely the famous L-B formula. Once the formalism has been tested on the L-B steady current, we applied it for the computation of the charge and energy densities, the

energy flow and the noise power. The presence of Friedel oscillations has been detected. We demonstrated also that point-like interaction in the junction modifies the linear dependence of the thermal noise on the temperature (Johnson-Nyquist formula). The formalism has been generalized in order to include the minimal coupling to an external time-independent electromagnetic field as well.

Summarizing, the star graph models proposed and analyzed in this paper represent relatively simple exactly solvable examples of quantum non-equilibrium systems in a steady state. For this reason they provide a nice laboratory for testing general ideas about non-equilibrium dynamics.

Our results can be generalized in various directions. First of all, one can consider more complicated networks with several junctions and loops, which can be crossed by magnetic fluxes. The basic idea for treating this case is to replace in the above formalism the scattering matrix \mathbb{S} with an effective one \mathbb{S}_{eff} , which takes into account all vertex interactions [64]-[68] and the presence of a magnetic field (see (3.101)). The derivation of the L-B steady current and the noise in this case is of particular physical interest and is currently under investigation [69].

Another possible generalization is the study of imperfect leads involving interactions with external potentials and/or self-interactions like those in the Luttinger liquid. More general boundary interactions, involving new vertex degrees of freedom of the type appearing in the resonant-level model [70], can be investigated in the above framework as well. We will discuss these issues elsewhere.

Acknowledgments:

It is a great pleasure to thank B. Bellazzini, V. Caudrelier, P. Sorba and E. Ragoucy for the intense collaboration on quantum field theory on graphs. Enlightening discussions and correspondence with P. Calabrese, B. Douçot, V. Kstrykin, I. Safi, R. Schrader and E. Vicari are also kindly acknowledged.

References

- [1] C. L. Kane and M. P. A. Fisher, Phys. Rev. Lett. **68**, 1220 (1992); Phys. Rev. **B 46**, 15233 (1992).
- [2] I. Safi, H. J. Schulz, Phys. Rev. B **52**, R17040 (1995).
- [3] C. Nayak, M. P. A. Fisher, A. W. W. Ludwig and H. H. Lin, Phys. Rev. B **59**, 15694 (1999).
- [4] I. Safi, P. Devillard, and T. Martin, Phys. Rev. Lett. **86**, 4628 (2001).
- [5] J.E. Moore and X.-G. Wen, Phys. Rev. B **66**, 115305 (2002).
- [6] H. Yi, Phys. Rev. B **65**, 195101 (2002).
- [7] S. Lal, S. Rao, and D. Sen, Phys. Rev. B **66**, 165327 (2002).
- [8] S. Chen, B. Trauzettel, and R. Egger, Phys. Rev. Lett. **89**, 226404 (2002).

- [9] K-V. Pham, F. Piechon, K-I Imura, P. Lederer, Phys. Rev. B **68**, 205110 (2003).
- [10] C. Chamon, M. Oshikawa, and I. Affleck, Phys. Rev. Lett. **91**, 206403 (2003); M. Oshikawa, C. Chamon, and I. Affleck, J. Stat. Mech. P02008 (2006).
- [11] F. Dolcini, H. Grabert, I. Safi and B. Trauzettel, Phys. Rev. Lett. **91**, 266402 (2003).
- [12] S. Rao and D. Sen, Phys. Rev. B **70**, 195115 (2004).
- [13] K. Kazymyrenko and B. Douçot, Phys. Rev. B **71**, 075110 (2005)
- [14] E.-A. Kim, M. J. Lawler, S. Vishveshwara, E. Fradkin, Phys. Rev. Lett. **95**, 176402 (2005); Phys. Rev. B **74**, 155324 (2006).
- [15] T. Enss, V. Meden, S. Andergassen, X. Barnabe-Theriat, W. Metzner, K. Schonhammer, Phys. Rev. B **71**, 155401 (2005); X. Barnabe-Theriat, A. Sedeki, V. Meden, K. Schonhammer, Phys. Rev. Lett. **94**, 136405 (2005); X. Barnabe-Theriat, A. Sedeki, V. Meden, K. Schonhammer, Phys. Rev. B **71**, 205327 (2005).
- [16] D. Friedan, cond-mat/0505084; cond-mat/0505085.
- [17] S. Das, S. Rao, D. Sen, Phys. Rev. B **74**, 045322 (2006).
- [18] B. Bellazzini and M. Mintchev, J. Phys. A **39**, 11101 (2006) [arXiv:hep-th/0605036].
- [19] B. Bellazzini, M. Mintchev and P. Sorba, J. Phys. A **40**, 2485 (2007) [arXiv:hep-th/0611090].
- [20] B. Bellazzini, M. Burrello, M. Mintchev and P. Sorba, Proc. Symp. Pure Math. **77**, 639 (2008), arXiv:0801.2852 [hep-th].
- [21] B. Bellazzini, P. Calabrese and M. Mintchev, Phys. Rev. B **79** 085122 (2009), [arXiv:0808.2719].
- [22] C.-Y. Hou and C. Chamon, Phys. Rev. B **77**, 155422 (2008).
- [23] S. Das, S. Rao, and A. Saha, Phys. Rev. B **77**, 155418 (2008); Europhys. Lett. **81**, 67001 (2008).
- [24] S. Das and S. Rao, Phys. Rev. B **78**, 205421 (2008).
- [25] C.-Y. Hou, E.-A. Kim, and C. Chamon, Phys. Rev. Lett. **102**, 076602 (2009).
- [26] A. Agarwal, S. Das, S. Rao and D. Sen, Phys. Rev. Lett. **103**, 026401 (2009); Erratum, Phys. Rev. Lett. **103**, 079903 (2009), arXiv:0810.3513 [cond-mat].
- [27] S. Das, S. Rao and A. Saha, Phys. Rev. B **79**, 155416 (2009).
- [28] B. Bellazzini, M. Mintchev and P. Sorba, J. Math. Phys. **51** 032302 (2010), arXiv:0810.3101 [hep-th].

- [29] I. Safi, arXiv:0906.2363 [cond-mat.]
- [30] B. Bellazzini, M. Mintchev and P. Sorba, Phys. Rev. B **80** 25441 (2009), arXiv:0907.4221[hep-th].
- [31] B. Bellazzini, M. Mintchev and P. Sorba, Phys. Rev. B **82** 195113 (2010), arXiv:1002.0206 [hep-th].
- [32] A. Soori, D. Sen, Europhys. Lett. **93** (2011) 57007.
- [33] J. A. McLennan, Phys. Rev. **115** (1959) 1405.
- [34] G. L. Eyink, J. L. Lebowitz and H. Spohn, J. Stat. Phys. **83** (1996) 385.
- [35] D. Ruelle, J. Stat. Phys. **98** (2000) 57.
- [36] L. Bertini, A. D. Sole, D. Gabrielli, G. Jona-Lasinio and C. Landim, Phys. Rev. Lett. **87** (2001) 040601.
- [37] V. Jaksic and C. A. Pillet, J. Stat. Phys. **108** (2002) 787.
- [38] T. Bodineau and B. Derrida, Phys. Rev. Lett. **92** (2004) 180601.
- [39] S. Sasa and H. Tasaki, J. Stat. Phys. **125** (2006) 125.
- [40] B. Derrida, arXiv:1012.1136 [cond-mat.stat-mech]
- [41] V. Kострыкин and R. Schrader, Fortschr. Phys. **48**, 703 (2000).
- [42] M. Harmer, J. Phys. A **33** (2000) 9015.
- [43] P. Kuchment, arXiv:0802.3442 [math-ph].
- [44] R. Landauer, IBM J. Res. Dev. **1** (1957) 233; Philos. Mag. **21** (1970) 863.
- [45] M. Büttiker, Phys. Rev. Lett. **57** (1986) 1761; IBM J. Res. Dev. **32** (1988) 317.
- [46] R. Kubo, M. Toda and N. Hashitsume, *Statistical Physics II*, (Springer, Berlin, 1985).
- [47] H. U. Baranger and A. D. Stone, Phys. Rev. B **40** (1989) 8169.
- [48] H. D. Cornean and A. Jensen, J. Math. Phys. **46** (2005) 042106-1.
- [49] R. Schrader, J. Phys. A **42** (2009) 495401, arXiv:0907.1522 [hep-th].
- [50] A. Liguori, M. Mintchev and L. Zhao, Commun. Math. Phys. **194**, 569 (1998) [arXiv:hep-th/9607085].
- [51] M. Mintchev, E. Ragoucy and P. Sorba, Phys. Lett. B **547**, 313 (2002) [arXiv:hep-th/0209052].

- [52] M. Mintchev, E. Ragoucy and P. Sorba, *J. Phys. A* **36**, 10407 (2003) [arXiv:hep-th/0303187].
- [53] M. Mintchev and P. Sorba, *J. Stat. Mech.* **0407**, P001 (2004) [arXiv:hep-th/0405264].
- [54] O. Bratteli and D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics 2*, (Springer, Berlin, 1996).
- [55] J. Friedel, *Phil. Mag.* **43** (1952) 153.
- [56] Y. Imry and R. Landauer, *Rev. Mod. Phys.* **71** (1999) S306.
- [57] Th. Martin and R. Landauer, *Phys. Rev. B* **45** (1992) 1742.
- [58] M. Büttiker, *Phys. Rev. B* **46** (1992) 12485.
- [59] Ya. M. Blanter and M. Büttiker, *Phys. Rep.* **336** (2000) 1.
- [60] I. S. Gradshteyn and I. M. Ryzhik, *Tables of Integrals, Series and Products*, (Elsevier, Amsterdam, 2007).
- [61] V. Kostrykin and R. Schrader, *Commun. Math. Phys.* **237** (2003) 161.
- [62] J. Bolte and J. Harrison, *J. Phys. A* **36** (2003) 2747.
- [63] B. Bellazzini, M. Burrello and M. Mintchev, unpublished.
- [64] V. Kostrykin and R. Schrader, *J. Math. Phys.* **42** (2001) 1563.
- [65] M. Mintchev and E. Ragoucy, *J. Phys. A* **40** (2007) 9515 [arXiv:0705.1322 [hep-th]].
- [66] E. Ragoucy, *J. Phys. A* **42** (2009) 295205, [arXiv:0901.2431 [hep-th]].
- [67] V. Caudrelier and E. Ragoucy, *Nucl. Phys. B* **828** (2010) 515, [arXiv:0907.5359 [math-ph]].
- [68] S. Khachatryan, A. Sedrakyan and P. Sorba, *Nucl. Phys. B* **825** (2010) 444, [arXiv:0904.2688 [cond-mat.mes-hall]].
- [69] V. Caudrelier, M. Mintchev and E. Ragoucy, (work in progress).
- [70] D. Bernard and B. Doyon, arXiv:1105.1695 [math-ph]