

# The link concordance invariant from Lee homology

John Pardon

23 July 2011

## Abstract

We use the knot homology of Khovanov and Lee to construct link concordance invariants generalizing the Rasmussen  $s$ -invariant of knots. The relevant invariant for a link is a filtration on a vector space of dimension  $2^{|L|}$ . The basic properties of the  $s$ -invariant all extend to the case of links; in particular, any orientable cobordism  $\Sigma$  between links induces a map between their corresponding vector spaces which is filtered of degree  $\chi(\Sigma)$ . A corollary of this construction is that any component preserving orientable cobordism from a Kh-thin link to a link split into  $k$  components must have genus at least  $\lfloor \frac{k}{2} \rfloor$ . In particular, no quasi-alternating link is concordant to a split link.

## 1 Introduction

Using Lee's modification [17] of Khovanov homology [13], Rasmussen [25] introduced for every knot  $K$  an even integer valued invariant, known as the  $s$ -invariant. It shares some of the basic properties of the knot signature; in particular it is a homomorphism from the group of smooth concordance classes of knots to  $2\mathbb{Z}$ , and gives a lower bound for twice the smooth slice genus (though the signature also does this in the *topological* category, whereas the  $s$ -invariant does not). The definition of the  $s$ -invariant is purely combinatorial, and, like many other knot invariants coming out of quantum algebra, it so far lacks any intrinsic geometric definition. One of the main reasons for interest in this invariant is that it is by definition algorithmically computable (though some cleverness is needed to do large calculations quickly, c.f. Bar-Natan [3] and Freedman, Gompf, Morrison, and Walker [8]), and is one of the few tools known to give useful lower bounds on the smooth slice genus of knots. In particular, Rasmussen [25] showed via direct calculation that  $s(T_{p,q}) = (p-1)(q-1)$ , thus proving the Milnor conjecture that  $g_4(T_{p,q}) = \frac{1}{2}(p-1)(q-1)$ , a hard theorem of Kronheimer and Mrowka [14, 16] [15] proved (twice) using gauge theory.

In this paper, we consider the natural generalization of the  $s$ -invariant to a concordance invariant of links. Everything we do will be in the smooth category. Since the  $s$ -invariant can detect the deep differences between the smooth and topological categories in four dimensions, this restriction is in fact necessary for this theory. In particular, knot and link concordance is meant in the smooth sense.

Let us denote the Khovanov–Lee homology groups of a link by  $\text{Kh}_{\text{Lee}}^*(L)$ . Lee [17] showed that  $\text{Kh}_{\text{Lee}}^*(L)$  is a surprisingly simple group: there is an isomorphism  $\bigoplus_{\text{orientations of } L} \mathbb{Q} \xrightarrow{\sim} \text{Kh}_{\text{Lee}}^*(L)$ . We denote the former group by  $\mathcal{O}(L)$ , and in Section 4, we will define it as a

functor (we will specify the maps associated to cobordisms). Kevin Walker (communication on [MathOverflow](#) [26]) informs the author that there is an equivalence of functors between  $\mathbb{O}$  and  $\text{Kh}_{\text{Lee}}^*$  (Rasmussen [25] [24] has proved a sort of approximate equivalence of functors). The natural generalization of the  $s$ -invariant is thus the pull-back of the  $s$ -filtration on  $\text{Kh}_{\text{Lee}}^*(L)$  to a filtration on  $\mathbb{O}(L)$ . To get a numerical invariant, we take the following (which is perhaps slightly coarser).

**Definition 1.1.** For a link  $L \subseteq \mathbb{R}^3$ , we associate a function  $d_L : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}_{\geq 0}$  defined by  $d_L(h, s) = \dim(\text{Kh}_{\text{Lee}}^h(L))^s / (\text{Kh}_{\text{Lee}}^h(L))^{s+1}$  (here  $(\text{Kh}^*(L))^s$  denotes the subspace of elements of filtration level  $\geq s$ ).

For a knot  $K$ , it is a theorem of Rasmussen [25] that  $d_K(0, s(K) \pm 1) = 1$  and  $d_K$  is otherwise zero (this is the defining property of the  $s$ -invariant). For a link  $L$ , the vector space  $\text{Kh}_{\text{Lee}}^*(L)$  has dimension  $2^{|L|}$ , and, as one might expect, the support of the function  $d_L$  can be much more complicated.

**Theorem 1.2.** *Let  $L$  be a link with orientation  $\mathfrak{o}_1$ . The invariant  $d_L : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}_{\geq 0}$  satisfies the following basic properties:*

1.  $\sum_{s \equiv |L| + k \pmod{4}} d_L(h, s)$  equals one half the number of orientations  $\mathfrak{o}$  of  $L$  such that  $\text{lk}(\mathfrak{o}) - \text{lk}(\mathfrak{o}_1) = -j$  if  $k$  is even, and zero otherwise. Here  $\text{lk}(\mathfrak{o}) = \sum_{i < j} \text{lk}(L_i^{\mathfrak{o}}, L_j^{\mathfrak{o}})$  (sum over the components of  $L$ ).
2.  $d_{L_1 \sqcup L_2} = d_{L_1} * d_{L_2}$  (convolution).
3.  $d_{\bar{L}}(h, s) = d_L(-h, -s)$ .
4. If  $\Sigma$  is a component preserving orientable cobordism between  $L_1$  and  $L_2$  (i.e.  $H_0(L_i) \xrightarrow{\sim} H_0(\Sigma)$ ), then  $\sum_{s \geq a} d_{L_1}(h, s) \leq \sum_{s \geq a + \chi(\Sigma)} d_{L_2}(h, s)$  for all  $h \in \mathbb{Z}$ .
5.  $d_L$  is a link concordance invariant.

For links with a large number of components, it is reasonable to expect that the invariant  $d_L$  will be a strong invariant of link concordance. As one sees in the theorem above, the invariant  $d_L$  is best suited for studying cobordisms which do not merge components of  $L$ . In general, if one wants to derive information about a given orientable cobordism, then the relevant object is the  $s$ -filtration restricted to the subspace of  $\mathbb{O}(L)$  generated by those orientations extending to orientations of the cobordism. The larger this subspace, the more likely the invariant is to be useful.

Beliakova and Wehrli have defined an integer  $s(L, \mathfrak{o})$  for a link with an orientation [5]. This corresponds to the  $s$ -filtration restricted to the 2-dimensional subspace of  $\text{Kh}_{\text{Lee}}^*(L)$  corresponding to that orientation and its reverse. Just like for knots, one shows that on this subspace, the filtration is supported in two levels  $s \pm 1$ , and this defines  $s(L, \mathfrak{o})$ . This invariant is best suited for studying oriented cobordisms which are allowed to merge components of  $L$ . Examples show that the function  $\mathfrak{o} \mapsto s(L, \mathfrak{o})$  is a weaker invariant than  $d_L$ . One expects that  $d_L$  is a weaker invariant than the filtration on  $\mathbb{O}(L)$  but we don't have any examples to prove this at present (mainly because  $d_L$  is often easy to derive from  $\text{Kh}^*$ —which there

exist programs to compute—whereas the  $s$ -filtration is not). We discuss examples in Section 5 and at the end of Section 4.1.

In Section 3, we use the invariants  $d_L$  to derive the following corollary, which appears to be new.

**Corollary 1.3.** *A component preserving orientable cobordism between a Kh-thin link and a link split into  $m$  components must have genus at least  $\lfloor \frac{m}{2} \rfloor$ . In particular, Kh-thin links (in particular quasi-alternating links [Definition 3.5]) are not concordant to split links.*

It is known (via properties of the Alexander module) that alternating links are not concordant to split links [12]. It would be interesting to try to prove Corollary 1.3 (say, restricted to quasi-alternating links) using the Alexander module.

This corollary is interesting because the  $s$ -invariant for alternating *knots* is equal to the knot signature, and thus gives no new information (the inequality  $g_4^{\text{top}}(K) \geq \frac{1}{2} |\sigma(K)|$  is classical, see Murasugi [20, p416, Theorem 9.1]). It is very interesting to note that Khovanov homology has a reputation for being easy to compute (at least, compared to gauge theoretic invariants which give results similar to the Milnor conjecture), but hard to use to prove general theorems, since its structure in general is still poorly understood. Thus the above corollary is interesting in that it is a *general* statement which doesn't intrinsically involve Khovanov homology (at least, if one restricts to quasi-alternating links).

There have recently been efforts (see Freedman, Gompf, Morrison, and Walker [8]) to prove that some specific proposed counterexamples to the smooth 4-dimensional Poincaré conjecture are in fact exotic by proving some links are not slice in the standard  $\mathbb{B}^4$  (links that, by virtue of coming from Kirby diagrams for the proposed counterexample, are by definition slice in the proposed exotic  $\mathbb{B}^4$ ). By slice, we mean *strongly slice*, i.e. bounding a disjoint union of disks in  $\mathbb{B}^4$ . Since these are usually multi-component links, it may be helpful to compute the entire filtration on  $\text{Kh}_{\text{Lee}}^*(L)$ : for a link with many many components, this *a priori* may be a much stronger invariant than the set of  $s$ -invariant values for some associated knots which are implied to be slice if the link is slice (computing these  $s$ -invariant values was the strategy employed in [8]). We should, however, also note that, in accordance with the growing relations between Khovanov homology and gauge theory, some would conjecture that the  $s$ -filtration should be invariant under concordance of links in any *homotopy*  $\mathbb{R}^3 \times [0, 1]$ , and thus would not imply in any straightforward manner that any homotopy  $\mathbb{B}^4$  is exotic.

One thinks that an invariant of links similar to  $d_L$  could be defined using the Link Floer Homology of Ozsvath–Szabó [21, 23] as an appropriate generalization of the  $\tau$ -invariant. One would expect this invariant to satisfy similar properties as the  $s$ -filtration on  $\text{Kh}_{\text{Lee}}^*(L)$ . It is perhaps interesting to note that the vector space  $\mathbb{O}(L)$  appears in the Link Floer Homology theory in the guise of  $\wedge^* H_1(\#^{|L|} \mathbb{S}^1 \times \mathbb{S}^2)$  (once we take the union of our link with the unknot).

## 1.1 Acknowledgments

This paper represents part of the author's Senior Thesis at Princeton University, advised by Zoltán Szabó. I thank him for lots of generous time spent meeting and discussing mathematics. John Baldwin, the second reader for my thesis, also made some useful comments.

## 2 Khovanov–Lee homology

In this section, we give a quick review of Lee’s version [17] of Khovanov homology [13] aimed at our intended application. In the following definition, we set the Lee deformation parameter to  $a = 2$ . This is slightly more convenient than Lee’s original  $a = 1$ , since the genus of the cobordism doesn’t introduce multiplicative factors. For a good introduction to Khovanov, see Bar-Natan’s articles [1] and [2]. The maps for cobordisms were first proved consistent by Jacobsson [11].

**Theorem 2.1.** *For every oriented link  $L$ , there is an associated  $\mathbb{Z}$ -graded vector space  $\mathrm{Kh}_{\mathrm{Lee}}^*(L)$  over  $\mathbb{Q}$  (the grading  $*$  is called the homological grading). Furthermore, each  $\mathrm{Kh}_{\mathrm{Lee}}^h(L)$  carries a descending filtration, called the  $s$ -filtration. Each oriented cobordism  $\Sigma \subseteq \mathbb{R}^3 \times [0, 1]$  from  $L_1$  to  $L_2$  induces a homomorphism  $F_\Sigma : \mathrm{Kh}_{\mathrm{Lee}}^*(L_1) \rightarrow \mathrm{Kh}_{\mathrm{Lee}}^*(L_2)$  (defined up to  $\pm 1$ ) which respects the homological grading, and which is filtered of degree  $\chi(\Sigma)$ . We have the following additional properties:*

1.  $\mathrm{Kh}_{\mathrm{Lee}}^h(L)$  carries an absolute  $\mathbb{Z}/4\mathbb{Z}$  grading which is supported in gradings  $\equiv |L| \pmod{2}$ , and these two pieces have equal dimensions. The  $s$ -filtration breaks up as a filtration on each of the pieces, and furthermore the  $s$ -filtration on the degree  $k \in \mathbb{Z}/4\mathbb{Z}$  piece is supported on integers  $s \equiv k \pmod{4}$ .
2.  $\mathrm{Kh}_{\mathrm{Lee}}^*(L_1 \sqcup L_2) = \mathrm{Kh}_{\mathrm{Lee}}^*(L_1) \otimes \mathrm{Kh}_{\mathrm{Lee}}^*(L_2)$  (naturally), and this is an isomorphism of the homological grading and the  $s$ -filtration.
3.  $\mathrm{Kh}_{\mathrm{Lee}}^*(\bar{L})$  is naturally the dual of  $\mathrm{Kh}_{\mathrm{Lee}}^*(L)$ .
4. (due to Lee [17])  $\dim \mathrm{Kh}_{\mathrm{Lee}}^*(L) = 2^{|L|}$ . In fact,  $\dim \mathrm{Kh}_{\mathrm{Lee}}^h(L)$  is the number of orientations  $\mathfrak{o}$  of  $L$  such that  $\mathrm{lk}(\mathfrak{o}) - \mathrm{lk}(\mathfrak{o}_1) = -h$ , where  $\mathfrak{o}_1$  is the given orientation of  $L$ , and  $\mathrm{lk}(\mathfrak{o}) = \sum_{i < j} \mathrm{lk}(L_i^{\mathfrak{o}}, L_j^{\mathfrak{o}})$  (sum over the components of  $L$ ).
5.  $\mathrm{Kh}_{\mathrm{Lee}}^*$  is a functor from the appropriately defined category of links and cobordisms (see [7]).

*Remark 2.2.* Clark, Morrison, and Walker [7] and Caprau [6] have shown how to define Khovanov homology (with indeterminate  $a$ ) so that the maps associated to cobordisms no longer have a sign ambiguity. This requires adjoining  $i$  to the coefficient ring.

**Definition 2.3.** For a link  $L \subseteq \mathbb{R}^3$ , we associate a function  $d_L : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}_{\geq 0}$  defined by  $d_L(h, s) = \dim (\mathrm{Kh}_{\mathrm{Lee}}^h(L))^s / (\mathrm{Kh}_{\mathrm{Lee}}^h(L))^{s+1}$ . Here  $(\mathrm{Kh}_{\mathrm{Lee}}^h(L))^s$  means the subspace of filtration level  $\geq s$ .

Theorem 1.2 (the basic properties of  $d_L$ ) follows directly from the basic properties of  $\mathrm{Kh}_{\mathrm{Lee}}^*$  listed in Theorem 2.1.

## 3 Applications to link concordance

**Definition 3.1.** A cobordism  $\Sigma$  between two links  $L_1$  and  $L_2$  is *component preserving* iff  $H_0(L_i) \xrightarrow{\sim} H_0(\Sigma)$  for  $i \in \{1, 2\}$ . Note that a component preserving orientable cobordism of genus 0 is exactly a link concordance.




Figure 1: Local pictures of  $L_0$ ,  $L_1$ ,  $L_\infty$ .

*Remark 3.2.* One is perhaps also interested in relaxing the restrictive notion of *component* preserving cobordism to *color* preserving cobordism, where multiple components of the link could have the same color. Now certainly this case is also easily handled using the invariant  $\text{Kh}_{\text{Lee}}^*(L)$ . The necessary data is a coloring of the link, and a choice of relative orientation on each colored component (by relative orientation, we mean an orientation up to overall reversal). Then one is just interested in the restriction of the  $s$ -filtration to the subspace of  $\text{Kh}_{\text{Lee}}^*(L)$  generated by all orientations agreeing with the given relative orientations on each colored component.

**Lemma 3.3.** *The map  $F_\Sigma : \text{Kh}_{\text{Lee}}^*(L_1) \rightarrow \text{Kh}_{\text{Lee}}^*(L_2)$  induced by a component preserving orientable cobordism is an isomorphism of vector spaces.*

*Proof.* This follows from Rasmussen [25, p434 Proposition 4.1].  $\square$

**Definition 3.4.** A link  $L$  is said to be Kh-thin iff  $\text{Kh}^*(L)$  is supported on exactly two diagonals of the form  $q = 2h + a \pm 1$  ( $h = *$  is the homological grading).

By virtue of the spectral sequence from  $\text{Kh}^*(L)$  to  $\text{Kh}_{\text{Lee}}^*(L)$ , it follows that for a Kh-thin link, we have that the support of  $d_L$  is contained in the same two diagonals  $s = 2h + a \pm 1$ .

**Definition 3.5.** Ozsváth and Szabó [22] define the set of *quasi-alternating* links to be the set of links generated by the unknot using the following skein operation: if  $L_0$  and  $L_1$  are quasi-alternating and  $\det L_\infty = \det L_0 + \det L_1$ , then so is  $L_\infty$  (where  $L_0, L_1, L_\infty$  are given as in Figure 1).

It is standard that all non-split alternating links are quasi-alternating. Quasi-alternating links are known to be both Kh-thin and  $\widehat{HFK}$ -thin by Manolescu–Ozsváth [19], though Greene [9] has shown that there are non-quasi-alternating links that are both Kh-thin and  $\widehat{HFK}$ -thin.

**Proposition 3.6** (Corollary 1.3). *Let  $L$  be a Kh-thin link, and suppose  $\Sigma$  is a component preserving orientable cobordism between  $L$  and  $M = M_1 \sqcup \cdots \sqcup M_k$ . Then  $g(\Sigma) \geq \lfloor \frac{k}{2} \rfloor$ .*

*Proof.* Fix an orientation on  $L$ , which thus orients each  $M_i$ .

We know (from Theorem 1.2 property 1) that the support of the  $s$ -filtration on  $\text{Kh}_{\text{Lee}}^0(M_i)$  has diameter at least 2. Thus (by Theorem 1.2 property 2)  $\text{Kh}_{\text{Lee}}^0(M)$  has  $s$ -filtration of diameter at least  $2k$ . Since  $L$  is Kh-thin, the  $s$ -filtration on  $\text{Kh}_{\text{Lee}}^0(L)$  has diameter equal to 2 (using the spectral sequence from  $\text{Kh}^*$  to  $\text{Kh}_{\text{Lee}}^*$ ).

The cobordism and its reverse induce two maps:

$$\text{Kh}_{\text{Lee}}^0(L) \xrightarrow{F_\Sigma} \text{Kh}_{\text{Lee}}^0(M) \xrightarrow{F_{-\Sigma}} \text{Kh}_{\text{Lee}}^0(L) \quad (3.1)$$

These are both isomorphisms by Lemma 3.3. Also, we know that both maps are filtered of degree  $-2g(\Sigma)$ .

Without loss of generality, say the  $s$ -filtration on  $\text{Kh}_{\text{Lee}}^0(L)$  is supported in degrees  $\pm 1$ . Then since  $\text{Kh}_{\text{Lee}}^0(M) \xrightarrow{F_{-\Sigma}} \text{Kh}_{\text{Lee}}^0(L)$  is filtered of degree  $-2g(\Sigma)$ , the  $s$ -filtration on  $\text{Kh}_{\text{Lee}}^0(M)$  must be supported in degrees  $\leq 1 + 2g(\Sigma)$ . Similarly, looking at  $\text{Kh}_{\text{Lee}}^0(L) \xrightarrow{F_{\Sigma}} \text{Kh}_{\text{Lee}}^0(M)$ , we see that the  $s$ -filtration on  $\text{Kh}_{\text{Lee}}^0(M)$  must be supported in degrees  $\geq -1 - 2g(\Sigma)$ . Thus we have  $2 + 4g(\Sigma) \geq 2k$ , so  $g(\Sigma) \geq \lceil \frac{k-1}{2} \rceil = \lfloor \frac{k}{2} \rfloor$ .  $\square$

The following corollary to Proposition 3.6 is already known via properties of the Alexander module [12].

**Corollary 3.7.** *No non-split alternating link is concordant to a split link.*

## 4 The orientation group

In this section we define a (almost tautological)  $(1 + 1)$ -dimensional TQFT which we call the *orientation group*. It is isomorphic to the TQFT used to define Lee homology (with Lee deformation parameter  $a = 2$ ). In fact, Kevin Walker (communication on `MathOverflow` [26]) informs us that the orientation group is isomorphic to  $\text{Kh}_{\text{Lee}}^*$  as a functor. The point of this construction in this section is to give a natural intrinsic description of the maps associated to cobordisms.

For any manifold  $X$ , we let  $|X|$  denote the number of connected components of  $X$ .

**Definition 4.1.** For an orientable manifold  $X$ , let  $O(X)$  denote the set of orientations of  $X$ . Let  $\mathbb{O}(X)$  denote the  $\mathbb{Q}$ -vector space with generators canonically associated to the elements of  $O(X)$ . We also define a natural inner product  $\langle \cdot, \cdot \rangle$  on  $\mathbb{O}(X)$  by declaring that this basis be orthonormal.

**Definition 4.2.** Let  $\mathfrak{o} \mapsto \bar{\mathfrak{o}}$  denote reversal of orientation; this is an involution of  $O(X)$  and of  $\mathbb{O}(X)$ .

By a *relative orientation* on a manifold  $X$ , we mean an orientation up to overall reversal of orientation, that is, an element of  $O(X)/(\mathfrak{o} \mapsto \bar{\mathfrak{o}})$ .

**Definition 4.3.** We define a mod 4 grading on  $\mathbb{O}(X)$  by declaring that the  $+1$  eigenspace of  $\mathfrak{o} \mapsto \bar{\mathfrak{o}}$  have grading  $-|X|$  and that the  $-1$  eigenspace of  $\mathfrak{o} \mapsto \bar{\mathfrak{o}}$  have grading  $2 - |X|$ .

**Lemma 4.4.** *We have a natural isomorphism  $\mathbb{O}(X_1 \cup X_2) = \mathbb{O}(X_1) \otimes \mathbb{O}(X_2)$  which respects the involution  $\mathfrak{o} \mapsto \bar{\mathfrak{o}}$  as well as the mod 4 grading.*

*Proof.* Clearly  $O(X_1 \cup X_2) = O(X_1) \times O(X_2)$ , and this gives us the desired isomorphism of vector spaces, which clearly respects reversal of orientation. Now by examining the definition of the mod 4 grading in terms of the map  $\mathfrak{o} \mapsto \bar{\mathfrak{o}}$ , one easily sees that this implies that the mod 4 grading is preserved as well.  $\square$

Henceforth we shall only be interested in  $\mathbb{O}(X)$  in the case that  $X$  is a 1-manifold.

**Definition 4.5.** If  $A$  is an orientable cobordism between  $X$  and  $Y$ , then we define a map  $F_A : \mathbb{O}(X) \rightarrow \mathbb{O}(Y)$  (up to overall multiplication by  $\pm 1$ ) as follows. Let  $\sigma : O(A) \rightarrow \{\pm 1\}$  satisfy the property that reversing the orientation on some component  $A_1 \subseteq A$  multiplies the value of  $\sigma$  by  $(-1)^{\chi(A_1) - |A_1 \cap X| + |A_1 \cap Y|/2}$  (note that since  $A$  is orientable,  $\chi(A_1) - |A_1 \cap X| + |A_1 \cap Y| \equiv \chi(\text{closed surface}) \equiv 0 \pmod{2}$ ). Clearly there are two such functions  $\sigma$ , differing by a sign. Then we define (up to  $\pm 1$ ):

$$F_A(\alpha) := \sum_{\mathfrak{o} \in O(A)} \sigma(\mathfrak{o}) \langle \alpha, \mathfrak{o}|_X \rangle \mathfrak{o}|_Y \quad (4.1)$$

By definition, orientations of  $X$  which do not extend to  $A$  get annihilated by  $F_A$ . More generally, an orientation is sent to a linear combination of those orientations on  $Y$  which are compatible with the cobordism  $A$  and the input orientation of  $X$ . Rasmussen [25, p434 Proposition 4.1] showed a similar property of  $\text{Kh}_{\text{Lee}}^*$  in the process of defining the  $s$ -invariant.

**Lemma 4.6.** *The maps associated to cobordisms are functorial in the sense that if  $A$  is a cobordism between  $X$  and  $Y$  and  $B$  is a cobordism between  $Y$  and  $Z$ , then  $F_{A \cup_Y B} = F_B \circ F_A$ .*

*Proof.*

$$\begin{aligned} F_B(F_A(\alpha)) &= \sum_{\mathfrak{o}_B \in O(B)} \sum_{\mathfrak{o}_A \in O(A)} \sigma(\mathfrak{o}_B) \sigma(\mathfrak{o}_A) \langle \alpha, \mathfrak{o}|_X \rangle \langle \mathfrak{o}_A|_Y, \mathfrak{o}_B|_Y \rangle \mathfrak{o}_B|_Z \\ &= \sum_{\mathfrak{o} \in O(A \cup_Y B)} \sigma(\mathfrak{o}|_A) \sigma(\mathfrak{o}|_B) \langle \alpha, \mathfrak{o}|_X \rangle \mathfrak{o}|_Z \end{aligned} \quad (4.2)$$

Now just observe that the function  $O(A \cup_Y B) \rightarrow \{\pm 1\}$  given by  $\sigma(\mathfrak{o}|_A) \sigma(\mathfrak{o}|_B)$  satisfies the property which defines  $\sigma : O(A \cup_Y B) \rightarrow \{\pm 1\}$  for the construction of  $F_{A \cup_Y B}$ .  $\square$

**Lemma 4.7.** *The map  $F_A$  on  $\mathbb{O}$  is homogeneous of degree  $\chi(A)$  with respect to the mod 4 grading.*

*Proof.* Note that by definition of the mod 4 grading, we have:

$$\begin{aligned} \overline{F_A(\bar{\alpha})} = F_A(\alpha) &\iff F_A \text{ homogeneous of degree } |X| - |Y| \\ \overline{F_A(\bar{\alpha})} = -F_A(\alpha) &\iff F_A \text{ homogeneous of degree } 2 + |X| - |Y| \end{aligned} \quad (4.3)$$

Now we calculate:

$$\begin{aligned} \overline{F_A(\bar{\alpha})} &= \sum_{\mathfrak{o} \in O(A)} \sigma(\mathfrak{o}) \langle \bar{\alpha}, \mathfrak{o}|_X \rangle \bar{\mathfrak{o}}|_Y \\ &= \sum_{\mathfrak{o} \in O(A)} \sigma(\bar{\mathfrak{o}}) \langle \alpha, \mathfrak{o}|_X \rangle \mathfrak{o}|_Y \end{aligned} \quad (4.4)$$

Now by the definition of  $\sigma$ , this equals  $(-1)^{\chi(A) - |X| + |Y|/2} F_A(\alpha)$ . Thus we have:

$$\begin{aligned} \chi(A) - |X| + |Y| \equiv 0 \pmod{4} &\implies F_A \text{ homogeneous of degree } |X| - |Y| \\ \chi(A) - |X| + |Y| \equiv 2 \pmod{4} &\implies F_A \text{ homogeneous of degree } 2 + |X| - |Y| \end{aligned} \quad (4.5)$$

which exactly says  $F_A$  is homogeneous of degree  $\chi(A)$ .  $\square$

The following description shows the isomorphism with Lee's TQFT (with  $a = 2$ ).

**Lemma 4.8.** *The map  $F_A$  has the following alternative description. We decompose  $A$  into iterated handle additions (e.g. using a Morse function), and then to each of the handle additions, we associate maps as follows.*

*For a 0-handle, we map  $\alpha$  to  $\alpha \otimes (\mathfrak{o} - \bar{\mathfrak{o}})$ , where  $\mathfrak{o}$  is an orientation on the new circle.*

*For a 1-handle which splits a component, the map sends every orientation to its extension to the new manifold.*

*For a 1-handle which joins two components, the map sends orientations which do not extend to the new manifold to zero, and sends orientations which do extend to their natural extension multiplied by  $\pm 1$  depending on the orientation of the new merged circle.*

*For a 2-handle, the map sends  $\mathfrak{o} \otimes \alpha$  to  $\alpha$  and  $\bar{\mathfrak{o}} \otimes \alpha$  to  $\alpha$ .*

*Proof.* That it suffices to splice together the maps for elementary cobordisms follows from Lemma 4.6. We just have to calculate the maps coming from  $k$ -handle additions,  $k \in \{0, 1, 2\}$ . These are given completely explicitly by Definition 4.5, which gives the result.  $\square$

It is interesting to note that even with this trivial construction, there is a good reason why if we want to make  $\mathbb{O}(L)$  into a functor, we have no choice but to stipulate that maps are only defined up to  $\pm 1$ . For instance, consider the birth of a circle. Since the birth of a circle is the same cobordism as the birth of a circle followed by a reflection, they must induce the same map. However, the image of the birth of a circle is  $\mathfrak{o} - \bar{\mathfrak{o}}$ , and this clearly changes sign under reflection.

If we are interested in links *embedded in  $\mathbb{R}^3$* , and we want functoriality with respect to orientable cobordisms *embedded in  $\mathbb{R}^3 \times [0, 1]$* , then it is probably possible to twist by an appropriate homomorphism  $\pi_1(\{\text{unoriented loops in } \mathbb{R}^3\}) \rightarrow \{\pm 1\}$  to get rid of the sign ambiguity in  $\mathbb{O}$ . We note that Hatcher [10] has proved the Smale Conjecture, which implies that the space of unoriented *unknotted* loops in  $\mathbb{R}^3$  deformation retracts onto the space of unoriented circles in  $\mathbb{R}^3$ , and the fundamental group of this space is indeed  $\mathbb{Z}/2\mathbb{Z}$ . We suspect this type of twisting is morally what fixes the functoriality of Khovanov homology as in [7] and [6].

## 4.1 Properties of the $s$ -filtration on $\mathbb{O}(L)$

Under the equivalence between  $\text{Kh}_{\text{Lee}}^*(L)$  and  $\mathbb{O}(L)$ , we get a natural definition of the  $s$ -filtration on  $\mathbb{O}(L)$ . The space  $\mathbb{O}(L)$  carries a number of natural operations, and it is reasonable to ask how they respect the  $s$ -filtration. We answer a few of these questions in this section, using only the functorial properties of  $\mathbb{O}(L)$  under cobordism. Because we use these soft methods, the properties we derive here would also be valid for a hypothetical generalization of the  $\tau$ -invariant to links.

The following is a rough analogue of Livingston's result [18] that  $s(K_-) \leq s(K_+) \leq s(K_-) + 2$  (here  $K_-$  and  $K_+$  are differ at exactly one crossing, which is positive for  $K_+$  and negative for  $K_-$ ).

**Lemma 4.9.** *Suppose  $L_1$  and  $L_2$  differ by a single crossing change. There is of course a natural isomorphism  $\phi : \mathbb{O}(L_1) \xrightarrow{\sim} \mathbb{O}(L_2)$ . Let  $\mathbb{O}(L_1)^+$  denote the space generated by*



orientations in which the given crossing is positive (and similarly define  $\mathbb{O}(L_1)^-$ ,  $\mathbb{O}(L_2)^+$ , and  $\mathbb{O}(L_2)^-$ ). Then we have:

1.  $\phi : \mathbb{O}(L_1)^- \rightarrow \mathbb{O}(L_2)^+$  is filtered of degree 0.
2.  $\phi : \mathbb{O}(L_1) \rightarrow \mathbb{O}(L_2)$  composed with a sign flip depending on the orientation of a strand at the crossing is filtered of degree  $-2$ .

*Proof.* Our strategy is to find cobordisms which induce the required maps.

For statement 2, it suffices to show that  $\phi : \mathbb{O}(L_1)^\pm \rightarrow \mathbb{O}(L_2)^\mp$  composed with a sign flip depending on the orientation of a strand at the crossing is filtered of degree  $-2$ . For this, consider that there is a cobordism of Euler characteristic  $-2$  from  $L_1$  to  $L_2$  (pass the strands through each other to get an immersed cobordism of Euler characteristic 0, and then resolve the double point). The induced map is  $\phi$  followed by the required sign flip depending on orientation. Thus this map is filtered of degree  $-2$ . We needed to split up  $\mathbb{O}(L_i)$  according to the sign of the crossing, since this sign dictates how we resolve the double point of the cobordism.

For statement 1, consider the following. By Rasmussen,  $\mathbb{O}(T_{2,3})$  is supported in  $s$ -filtration levels 1 and 3. Since the mod 4 grading agrees with the  $s$ -filtration, we see that  $\mathfrak{o} + \bar{\mathfrak{o}}$  lies in filtration level 3. Thus the map  $\mathbb{O}(L_1)^- \rightarrow \mathbb{O}(L_1)^- \otimes \mathbb{O}(T_{2,3})$  given by  $\alpha \mapsto \alpha \otimes (\mathfrak{o} + \bar{\mathfrak{o}})$  is filtered of degree 3. Now consider an immersed cobordism starting at  $L_1 \sqcup T_{2,3}$  which first passes the strands of the crossing of  $L_1$  through each other to get  $L_2$ , then unknots the  $T_{2,3}$  in a similar manner, and then merges the resulting unknot with  $L_2$ . The two double points are of opposite signs (when the crossing goes from negative in  $L_1$  to positive in  $L_2$ ), so they can be tubed together to obtain a cobordism of genus 1. Thus the resulting map  $\mathbb{O}(L_1)^- \otimes \mathbb{O}(T_{2,3}) \rightarrow \mathbb{O}(L_2)^+$  is filtered of degree  $-3$ . The composite is easily seen to be  $\phi : \mathbb{O}(L_1)^- \rightarrow \mathbb{O}(L_2)^+$ , so we are done.  $\square$

**Definition 4.10.** Given a specific orientation  $\mathfrak{o}_i$  on a component  $L_i$  of  $L$ , let  $\text{Res}_{\mathfrak{o}_i} : \mathbb{O}(L) \rightarrow \mathbb{O}(L)$  be orthogonal projection onto the subspace where  $L_i$  is oriented by  $\mathfrak{o}_i$ . For a relative orientation  $\mathfrak{o}_{ij}$  of  $L_i \cup L_j$  (two components of  $L$ ), let  $\text{Res}_{\mathfrak{o}_{ij}, \bar{\mathfrak{o}}_{ij}} : \mathbb{O}(L) \rightarrow \mathbb{O}(L)$  be projection onto the subspace where  $L_i \cup L_j$  has this relative orientation, composed with multiplication by  $\sigma : \mathbb{O}(L) \rightarrow \{\pm 1\}$  which flips sign depending on the orientation on  $L_i$ .

The following should be thought of as a generalization of Rasmussen's theorem that characterizes  $d_K$  for knots  $K$ .

**Lemma 4.11.** *The operators  $\text{Res}_{\mathfrak{o}_i}$  and  $\text{Res}_{\mathfrak{o}_{ij}, \bar{\mathfrak{o}}_{ij}}$  are both filtered of degree  $-2$ .*

*Proof.* For  $\text{Res}_{\mathfrak{o}_{ij}, \bar{\mathfrak{o}}_{ij}}$ , consider the cobordism formed by first adding a 1-handle connecting  $L_i$  and  $L_j$  (in such a way that the given relative orientation extends over the cobordism) and then adding a second 1-handle splitting the resulting component back into  $L_i \cup L_j$ . Clearly this cobordism induces the map  $\text{Res}_{\mathfrak{o}_{ij}, \bar{\mathfrak{o}}_{ij}} : \mathbb{O}(L) \rightarrow \mathbb{O}(L)$ , and it has Euler characteristic  $-2$ , so we are done.

For  $\text{Res}_{\mathfrak{o}_i}$ , let  $U$  be the unknot, and consider the map  $\mathbb{O}(L) \rightarrow \mathbb{O}(L) \otimes \mathbb{O}(U) = \mathbb{O}(L \sqcup U)$  which is  $\alpha \mapsto \alpha \otimes \mathfrak{o}$ . This is filtered of degree  $-1$ . Now compose with the map  $\mathbb{O}(L \sqcup U) \rightarrow \mathbb{O}(L)$  given by the cobordism obtained by adding a 1-handle to merge the unknot and  $L_i$

(such that the orientation  $\mathfrak{o}$  and the desired orientation  $\mathfrak{o}_i$  extend over the 1-handle). This cobordism has Euler characteristic  $-1$ . Thus the composition map is filtered of degree  $-2$ ; this map is also clearly  $\text{Res}_{\mathfrak{o}_i}$ .  $\square$

**Lemma 4.12.** *The map  $\mathbb{O}(L) \rightarrow \mathbb{O}(L)$  defined by flipping the sign according to the orientation of some particular component of  $L$  is filtered of degree  $-2$ .*

*Proof.* Such a map is a linear combination of  $\text{Res}_{\mathfrak{o}_i}$  and  $\text{Res}_{\bar{\mathfrak{o}}_i}$ .  $\square$

Suppose we have a relative orientation  $(\mathfrak{o}, \bar{\mathfrak{o}})$  of  $L$ . Let  $V_{\mathfrak{o}, \bar{\mathfrak{o}}} \subseteq \mathbb{O}(L)$  be the subspace generated by  $\mathfrak{o}$  and  $\bar{\mathfrak{o}}$ . Then let us consider the restriction of the  $s$ -filtration to  $V_{\mathfrak{o}, \bar{\mathfrak{o}}} = \mathbb{Q}(\mathfrak{o} + \bar{\mathfrak{o}}) \oplus \mathbb{Q}(\mathfrak{o} - \bar{\mathfrak{o}})$ . Note that this direct sum decomposition is into mod 4 graded pieces, thus the elements  $\mathfrak{o} + \bar{\mathfrak{o}}$  and  $\mathfrak{o} - \bar{\mathfrak{o}}$  are sent to different mod 4 gradings in  $\text{Kh}_{\text{Lee}}^*(L)$  which differ by exactly 2. Thus the  $s$ -filtration on  $V_{\mathfrak{o}, \bar{\mathfrak{o}}}$  is completely described by the two integers  $s(\mathfrak{o} + \bar{\mathfrak{o}})$  and  $s(\mathfrak{o} - \bar{\mathfrak{o}})$  (which differ by 2 mod 4). By Lemma 4.12,  $s(\mathfrak{o} + \bar{\mathfrak{o}})$  and  $s(\mathfrak{o} - \bar{\mathfrak{o}})$  differ by exactly two, and we let the oriented  $s(L, \mathfrak{o}) = \frac{1}{2}[s(\mathfrak{o} + \bar{\mathfrak{o}}) + s(\mathfrak{o} - \bar{\mathfrak{o}})]$ .

**Definition 4.13.** The invariant constructed in the previous paragraph is  $s(L, \mathfrak{o})$ . It was first defined by Beliakova and Wehrli [5].

For a knot  $K$ , there is just one relative orientation, and this gives Rasmussen's invariant  $s(K)$ . For links, however, there is much to the  $s$ -filtration on  $\text{Kh}_{\text{Lee}}^*(L)$  that is not captured by the function  $\mathfrak{o} \mapsto s(L, \mathfrak{o})$ . For example, for any alternating link with zero linking matrix, all  $s(L, \mathfrak{o})$  are equal (say, to  $s_0$ ), and  $d_L(h, s) = 2^{|L|-1} \delta(h) [\delta(s - s_0 - 1) + \delta(s - s_0 + 1)]$  (where  $\delta(0) = 1$  and  $\delta(\text{nonzero}) = 0$ ). The unlink on  $n$  components also has all  $s(L, \mathfrak{o})$  equal (this time to  $1 - n$ ), however in this case  $d_L(h, s) = \delta(h) \binom{|L|}{\frac{n}{2} + \frac{s}{2}}$  (the binomial coefficient is understood to equal zero when the lower entry is not an integer). Thus for link concordance,  $d_L$  is a stronger invariant than the function  $\mathfrak{o} \mapsto s(L, \mathfrak{o})$ .

## 5 Examples

We now summarize some calculations of the invariant  $d_L : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}_{\geq 0}$  for some links  $L$ . We used the package `KnotTheory` maintained by Bar-Natan [4], in particular the program to calculate Khovanov homology written by Scott Morrison. This allows us to calculate  $\text{Kh}^*(L)$  for the link in question. We use the simple fact that if  $\dim \text{Kh}^h(L) = \dim \text{Kh}_{\text{Lee}}^h(L)$ , then by virtue of the spectral sequence from  $\text{Kh}^*(L)$  to  $\text{Kh}_{\text{Lee}}^*(L)$ , the support of the  $s$ -filtration on  $\text{Kh}_{\text{Lee}}^h(L)$  is given exactly by the  $q$ -graded dimension of  $\text{Kh}^h(L)$ . Many interesting links have lots of crossings, and thus computing the Khovanov homology is time consuming on a computer; we just list the cases that we have been able to compute.

Most of the links in the standard link tables are quasi-alternating, so they do not present a particularly interesting case for the filtration on  $\text{Kh}_{\text{Lee}}^*(L)$  (it is just supported in two levels, so only their absolute height is interesting). So instead, we've taken as our examples some links with extra structure.

The function  $d_L$  is a link concordance invariant, and thus there are some easy corollaries using Theorem 1.2 distinguishing the link concordance classes of the links we consider below from other links whose  $d_L$  one could calculate (e.g. one can easily see which are concordant

to a quasi-alternating link). Theorem 1.2 also implies effective bounds on the genus of component preserving orientable cobordisms between these links and links with certain splitting numbers.

We use the following notation:

$$\delta(n) = \begin{cases} 1 & n = 0 \\ 0 & \text{otherwise} \end{cases} \quad (5.1)$$

## 5.1 Cablings of $T_{2,p}$

Let  $L_p$  be the  $(2, 0)$ -cabling of  $T_{2,p}$ . Then the linking matrix of  $L_p$  is zero, and we have (for  $p$  odd,  $1 \leq p \leq 11$ ):

$$d_{L_p}(h, s) = \delta(h)[\delta(s) + \delta(s - 2) + \delta(s - (2p - 4)) + \delta(s - (2p - 2))] \quad (5.2)$$

We conjecture that this is true for all odd  $p \geq 1$ . It should be the case also that  $\delta(s) + \delta(s - 2)$  represents the orientations where the two components are oriented in the opposite direction, and  $\delta(s - (2p - 4)) + \delta(s - (2p - 2))$  represents the orientations where the two components are oriented in the same direction.

## 5.2 $T_{p,p}$

We now consider the  $(p, p)$ -torus links for  $1 \leq p \leq 6$ . We have:

$$\begin{aligned} d_{T_{1,1}}(h, s) &= \delta(h)[\delta(s - 1) + \delta(s + 1)] \\ d_{T_{2,2}}(h, s) &= \delta(h)[\delta(s) + \delta(s - 2)] \\ &\quad + \delta(h - 2)[\delta(s + 2) + \delta(s)] \\ d_{T_{3,3}}(h, s) &= \delta(h)[\delta(s - 3) + \delta(s - 5)] \\ &\quad + \delta(h - 4)[\delta(s + 3) + 3\delta(s + 1) + 2\delta(s - 1)] \\ d_{T_{4,4}}(h, s) &= \delta(h)[\delta(s - 8) + \delta(s - 10)] \\ &\quad + \delta(h - 6)[\delta(s + 2) + 4\delta(s) + 3\delta(s - 2)] \\ &\quad + \delta(h - 8)[\delta(s + 4) + 3\delta(s + 2) + 2\delta(s)] \\ d_{T_{5,5}}(h, s) &= \delta(h)[\delta(s - 15) + \delta(s - 17)] \\ &\quad + \delta(h - 8)[\delta(s - 1) + 5\delta(s - 3) + 4\delta(s - 5)] \\ &\quad + \delta(h - 12)[\delta(s + 5) + 5\delta(s + 3) + 9\delta(s + 1) + 5\delta(s - 1)] \\ d_{T_{6,6}}(h, s) &= \delta(h)[\delta(s - 24) + \delta(s - 26)] \\ &\quad + \delta(h - 10)[\delta(s - 6) + 6\delta(s - 8) + 5\delta(s - 10)] \\ &\quad + \delta(h - 16)[\delta(s + 4) + 4\delta(s + 2) + 14\delta(s) + 9\delta(s - 2)] \\ &\quad + \delta(h - 18)[\delta(s + 6) + 5\delta(s + 4) + 9\delta(s + 2) + 5\delta(s)] \end{aligned} \quad (5.3)$$

For  $p = 5, 6$ , computing the final answer requires use of Rasmussen [25, p428 Lemma 3.5], in particular its consequence that the dimensions of  $\text{Kh}_{\text{Lee}}^*$  supported in  $s$ -filtration level  $|L|$

and  $|L| + 2$  are equal (this enables us to see which parts of  $\text{Kh}^*$  are killed in the spectral sequence).

Unfortunately, we see no obvious pattern in  $d_{T_{p,p}}$  which is likely to continue for larger values of  $p$ .

## References

- [1] Dror Bar-Natan. On Khovanov’s categorification of the Jones polynomial. *Algebr. Geom. Topol.*, 2:337–370 (electronic), 2002.
- [2] Dror Bar-Natan. Khovanov’s homology for tangles and cobordisms. *Geom. Topol.*, 9:1443–1499, 2005.
- [3] Dror Bar-Natan. Fast Khovanov homology computations. *J. Knot Theory Ramifications*, 16(3):243–255, 2007.
- [4] Dror Bar-Natan and Scott Morrison. KnotTheory‘ Package. [http://katlas.org/wiki/The\\_Mathematica\\_Package\\_KnotTheory‘](http://katlas.org/wiki/The_Mathematica_Package_KnotTheory%27), 2008.
- [5] Anna Beliakova and Stephan Wehrli. Categorification of the colored Jones polynomial and Rasmussen invariant of links. *Canad. J. Math.*, 60(6):1240–1266, 2008.
- [6] Carmen Livia Caprau.  $\text{sl}(2)$  tangle homology with a parameter and singular cobordisms. *Algebr. Geom. Topol.*, 8(2):729–756, 2008.
- [7] David Clark, Scott Morrison, and Kevin Walker. Fixing the functoriality of Khovanov homology. *Geom. Topol.*, 13(3):1499–1582, 2009.
- [8] Michael Freedman, Robert Gompf, Scott Morrison, and Kevin Walker. Man and machine thinking about the smooth 4-dimensional Poincaré conjecture. *Quantum Topol.*, 1(2):171–208, 2010.
- [9] Joshua Greene. Homologically thin, non-quasi-alternating links. *Math. Res. Lett.*, 17(1):39–49, 2010.
- [10] Allen E. Hatcher. A proof of the Smale conjecture,  $\text{Diff}(S^3) \simeq \text{O}(4)$ . *Ann. of Math. (2)*, 117(3):553–607, 1983.
- [11] Magnus Jacobsson. An invariant of link cobordisms from Khovanov homology. *Algebr. Geom. Topol.*, 4:1211–1251 (electronic), 2004.
- [12] Akio Kawauchi. On alternation numbers of links. *Topology Appl.*, 157(1):274–279, 2010.
- [13] Mikhail Khovanov. A categorification of the Jones polynomial. *Duke Math. J.*, 101(3):359–426, 2000.
- [14] P. B. Kronheimer and T. S. Mrowka. Gauge theory for embedded surfaces. I. *Topology*, 32(4):773–826, 1993.

- [15] P. B. Kronheimer and T. S. Mrowka. The genus of embedded surfaces in the projective plane. *Math. Res. Lett.*, 1(6):797–808, 1994.
- [16] P. B. Kronheimer and T. S. Mrowka. Gauge theory for embedded surfaces. II. *Topology*, 34(1):37–97, 1995.
- [17] Eun Soo Lee. An endomorphism of the Khovanov invariant. *Adv. Math.*, 197(2):554–586, 2005.
- [18] Charles Livingston. Computations of the Ozsváth-Szabó knot concordance invariant. *Geom. Topol.*, 8:735–742 (electronic), 2004.
- [19] Ciprian Manolescu and Peter Ozsváth. On the Khovanov and knot Floer homologies of quasi-alternating links. In *Proceedings of Gökova Geometry-Topology Conference 2007*, pages 60–81. Gökova Geometry/Topology Conference (GGT), Gökova, 2008.
- [20] Kunio Murasugi. On a certain numerical invariant of link types. *Trans. Amer. Math. Soc.*, 117:387–422, 1965.
- [21] Peter Ozsváth and Zoltán Szabó. Holomorphic disks and knot invariants. *Adv. Math.*, 186(1):58–116, 2004.
- [22] Peter Ozsváth and Zoltán Szabó. On the Heegaard Floer homology of branched double-covers. *Adv. Math.*, 194(1):1–33, 2005.
- [23] Peter Ozsváth and Zoltán Szabó. Holomorphic disks, link invariants and the multi-variable Alexander polynomial. *Algebr. Geom. Topol.*, 8(2):615–692, 2008.
- [24] Jacob Rasmussen. Khovanov’s invariant for closed surfaces. *arXiv math.GT 0502527*, 2005.
- [25] Jacob Rasmussen. Khovanov homology and the slice genus. *Invent. Math.*, 182(2):419–447, 2010.
- [26] Kevin Walker (mathoverflow.net/users/284). Is the complete functorial structure for Khovanov–Lee homology known? MathOverflow. <http://mathoverflow.net/questions/70902> (version: 2011-07-22).