

**NEW INEQUALITIES OF OSTROWSKI TYPE FOR MAPPINGS
WHOSE DERIVATIVES ARE s -CONVEX IN THE SECOND
SENSE VIA FRACTIONAL INTEGRALS**

ERHAN SET

ABSTRACT. New identity similar to an identity of [13] for fractional integrals have been defined. Then making use of this identity, some new Ostrowski type inequalities for Riemann-Liouville fractional integral have been developed. Our results have some relationships with the results of Alomari et. al., proved in [13] [published in. Appl. Math. Lett. 23 (2010) 1071-1076] and the analysis used in the proofs is simple.

1. INTRODUCTION AND PRELIMINARY RESULTS

In 1938, A.M. Ostrowski proved the following interesting and useful integral inequality ([30], see also [29, page 468]):

Theorem 1. *Let $f : I \rightarrow \mathbb{R}$, where $I \subseteq \mathbb{R}$ is an interval, be a mapping differentiable in the interior I° of I , and let $a, b \in I^\circ$ with $a < b$. If $|f'(x)| \leq M$ for all $x \in [a, b]$, then the following inequality holds:*

$$(1.1) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq M(b-a) \left[\frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right]$$

for all $x \in [a, b]$. The constant $\frac{1}{4}$ is the best possible in the sense that it cannot be replaced by a smaller one.

This inequality gives an upper bound for the approximation of the integral average $\frac{1}{(b-a)} \int_a^b f(t) dt$ by the value $f(x)$ at point $x \in [a, b]$. In recent years, such inequalities were studied extensively by many researchers and numerous generalizations, extensions and variants of them appeared in a number of papers see ([1]-[13])

In [15], the class of functions which are s -convex in the second sense has been introduced by Hudzik and Maligranda as the following:

Definition 1. *A function $f : [0, \infty) \rightarrow \mathbb{R}$ is said to be s -convex in the second sense if*

$$f(\lambda x + (1 - \lambda)y) \leq \lambda^s f(x) + (1 - \lambda)^s f(y)$$

for all $x, y \in [0, \infty)$, $\lambda \in [0, 1]$ and for some fixed $s \in (0, 1]$. This class of s -convex functions is usually denoted by K_s^2 .

2000 *Mathematics Subject Classification.* 26A33, 26A51, 26D07, 26D10, 26D15.

Key words and phrases. Ostrowski type inequality, s -convex function, , Riemann-Liouville fractional integral.

It can be easily seen that for $s = 1$, s -convexity reduces to ordinary convexity of functions defined on $[0, \infty)$.

In [14], S.S. Dragomir and S. Fitzpatrick proved a variant of Hadamard's inequality which holds for s -convex functions in the second sense:

Theorem 2. *Suppose that $f : [0, \infty) \rightarrow [0, \infty)$ is an s -convex function in the second sense, where $s \in (0, 1)$, and let $a, b \in [0, \infty)$, $a < b$. If $f' \in L^1([a, b])$, then the following inequalities hold:*

$$(1.2) \quad 2^{s-1} f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{s+1}.$$

The constant $k = \frac{1}{s+1}$ is the best possible in the second inequality in (1.2).

The following identity is proved by Alomari et.al. (see [13])

Lemma 1. *Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable mapping on I° where $a, b \in I$ with $a < b$. If $f' \in L[a, b]$, then we have the equality:*

$$\begin{aligned} f(x) - \frac{1}{b-a} \int_a^b f(t) dt &= \frac{(x-a)^2}{b-a} \int_0^1 t f'(tx + (1-t)a) dt \\ &\quad - \frac{(b-x)^2}{b-a} \int_0^1 t f'(tx + (1-t)b) dt \end{aligned}$$

for each $x \in [a, b]$.

Using the Lemma 1, Alomari et al. in [13] established the following results which holds for s -convex functions in the second sense.

Theorem 3. *Let $f : I \subset [0, \infty) \rightarrow \mathbb{R}$ be a differentiable mapping on I° such that $f' \in L[a, b]$, where $a, b \in I$ with $a < b$. If $|f'|$ is s -convex in the second sense on $[a, b]$ for some fixed $s \in (0, 1]$ and $|f'(x)| \leq M$, $x \in [a, b]$, then we have the inequality:*

$$(1.3) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{M}{b-a} \left[\frac{(x-a)^2 + (b-x)^2}{s+1} \right],$$

for each $x \in [a, b]$.

Theorem 4. *Let $f : I \subset [0, \infty) \rightarrow \mathbb{R}$ be a differentiable mapping on I° such that $f' \in L[a, b]$, where $a, b \in I$ with $a < b$. If $|f'|^q$ is s -convex in the second sense on $[a, b]$ for some fixed $s \in (0, 1]$, $q > 1$, $p = \frac{q}{q-1}$ and $|f'(x)| \leq M$, $x \in [a, b]$, then we have the inequality:*

$$(1.4) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{M}{b-a} \left[\frac{(x-a)^2 + (b-x)^2}{s+1} \right],$$

for each $x \in [a, b]$.

Theorem 5. *Let $f : I \subset [0, \infty) \rightarrow \mathbb{R}$ be a differentiable mapping on I° such that $f' \in L[a, b]$, where $a, b \in I$ with $a < b$. If $|f'|^q$ is s -convex in the second sense on $[a, b]$ for some fixed $s \in (0, 1]$, $q \geq 1$ and $|f'(x)| \leq M$, $x \in [a, b]$, then we have the inequality:*

$$(1.5) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq M \left(\frac{2}{s+1} \right)^{\frac{1}{q}} \left[\frac{(x-a)^2 + (b-x)^2}{2(b-a)} \right],$$

for each $x \in [a, b]$.

Theorem 6. Let $f : I \subset [0, \infty) \rightarrow \mathbb{R}$ be a differentiable mapping on I° such that $f' \in L[a, b]$, where $a, b \in I$ with $a < b$. If $|f'|^q$ is s -concave in the second sense on $[a, b]$ for some fixed $s \in (0, 1]$, $q > 1$ and $p = \frac{q}{q-1}$, then we have the inequality:

$$(1.6) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{2^{(s-1)/q}}{(1+p)^{1/p} (b-a)} \left[(x-a)^2 \left| f' \left(\frac{x+a}{2} \right) \right| + (b-x)^2 \left| f' \left(\frac{b+x}{2} \right) \right| \right],$$

for each $x \in [a, b]$.

For other recent results concerning s -convex functions see [13]-[17].

We give some necessary definitions and mathematical preliminaries of fractional calculus theory which are used throughout this paper.

Definition 2. Let $f \in L_1[a, b]$. The Riemann-Liouville integrals $J_{a+}^\alpha f$ and $J_{b-}^\alpha f$ of order $\alpha > 0$ with $a \geq 0$ are defined by

$$J_{a+}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t) dt, \quad x > a$$

and

$$J_{b-}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t) dt, \quad x < b,$$

where $\Gamma(\alpha) = \int_0^\infty e^{-t} t^{\alpha-1} dt$. Here, $J_{a+}^0 f(x) = J_{b-}^0 f(x) = f(x)$.

In the case of $\alpha = 1$, the fractional integral reduces to the classical integral. Some recent results and properties concerning this operator can be found ([18]-[28]).

Inspired and motivated by the recent results given in [13], [18]-[23] and [27], in the present note, we establish new Ostrowski type inequalities for s -convex functions in the second sense via Riemann-Liouville fractional integral. An interesting feature of our results is that they provide new estimates on these types of inequalities for fractional integrals.

2. OSTROWSKI TYPE INEQUALITIES VIA FRACTIONAL INTEGRALS

In order to prove our main results we need the following identity:

Lemma 2. Let $f : [a, b] \rightarrow \mathbb{R}$ be a differentiable mapping on (a, b) with $a < b$. If $f' \in L[a, b]$, then for all $x \in [a, b]$ and $\alpha > 0$ we have:

$$(2.1) \quad \left(\frac{(x-a)^\alpha + (b-x)^\alpha}{b-a} \right) f(x) - \frac{\Gamma(\alpha+1)}{(b-a)} [J_{x-}^\alpha f(a) + J_{x+}^\alpha f(b)] = \frac{(x-a)^{\alpha+1}}{b-a} \int_0^1 t^\alpha f'(tx + (1-t)a) dt - \frac{(b-x)^{\alpha+1}}{b-a} \int_0^1 t^\alpha f'(tx + (1-t)b) dt$$

where $\Gamma(\alpha) = \int_0^\infty e^{-t} t^{\alpha-1} dt$.

Proof. By integration by parts, we can state

$$\begin{aligned}
& \int_0^1 t^\alpha f'(tx + (1-t)a) dt \\
&= t^\alpha \frac{f(tx + (1-t)a)}{x-a} \Big|_0^1 - \int_0^1 \alpha t^{\alpha-1} \frac{f(tx + (1-t)a)}{x-a} dt \\
(2.2) \quad &= \frac{f(x)}{x-a} - \frac{\alpha}{x-a} \int_a^x \frac{(a-u)^{\alpha-1}}{(a-x)^{\alpha-1}} \frac{f(u)}{x-a} du \\
&= \frac{f(x)}{x-a} - \frac{\alpha \Gamma(\alpha)}{(x-a)^{\alpha+1}} \frac{1}{\Gamma(\alpha)} \int_a^x (u-a)^{\alpha-1} f(u) du
\end{aligned}$$

and

$$\begin{aligned}
& \int_0^1 t^\alpha f'(tx + (1-t)b) dt \\
&= t^\alpha \frac{f(tx + (1-t)b)}{x-b} \Big|_0^1 - \int_0^1 \alpha t^{\alpha-1} \frac{f(tx + (1-t)b)}{x-b} dt \\
(2.3) \quad &= \frac{f(x)}{x-b} - \frac{\alpha}{x-b} \int_b^x \frac{(b-u)^{\alpha-1}}{(b-x)^{\alpha-1}} \frac{f(u)}{x-b} du \\
&= \frac{f(x)}{x-b} + \frac{\alpha \Gamma(\alpha)}{(b-x)^{\alpha+1}} \frac{1}{\Gamma(\alpha)} \int_x^b (b-u)^{\alpha-1} f(u) du.
\end{aligned}$$

Multiplying the both sides of (2.2) and (2.3) by $\frac{(x-a)^{\alpha+1}}{b-a}$ and $\frac{(b-x)^{\alpha+1}}{b-a}$, respectively, we have

$$(2.4) \quad \frac{(x-a)^{\alpha+1}}{b-a} \int_0^1 t^\alpha f'(tx + (1-t)a) dt = \frac{(x-a)^\alpha f(x)}{b-a} - \frac{\Gamma(\alpha+1)}{b-a} J_{x-}^\alpha f(a)$$

and

$$(2.5) \quad \frac{(b-x)^{\alpha+1}}{b-a} \int_0^1 t^\alpha f'(tx + (1-t)b) dt = -\frac{(b-x)^\alpha f(x)}{b-a} + \frac{\Gamma(\alpha+1)}{b-a} J_{x+}^\alpha f(b).$$

From (2.4) and (2.5), it is obtained desired result. \square

Using Lemma 2, we can obtain the following fractional integral inequalities:

Theorem 7. *Let $f : [a, b] \subset [0, \infty) \rightarrow \mathbb{R}$, be a differentiable mapping on (a, b) with $a < b$ such that $f' \in L[a, b]$. If $|f'|$ is s -convex in the second sense on $[a, b]$ for some fixed $s \in (0, 1]$ and $|f'(x)| \leq M$, $x \in [a, b]$, then the following inequality for fractional integrals with $\alpha > 0$ holds:*

$$\begin{aligned}
(2.6) \quad & \left| \left(\frac{(x-a)^\alpha + (b-x)^\alpha}{b-a} \right) f(x) - \frac{\Gamma(\alpha+1)}{(b-a)} [J_{x-}^\alpha f(a) + J_{x+}^\alpha f(b)] \right| \\
& \leq \frac{M}{b-a} \left(1 + \frac{\Gamma(\alpha+1)\Gamma(s+1)}{\Gamma(\alpha+s+1)} \right) \left[\frac{(x-a)^{\alpha+1} + (b-x)^{\alpha+1}}{\alpha+s+1} \right]
\end{aligned}$$

where Γ is Euler Gamma function.

Proof. From (2.1) and since $|f'|$ is a s -convex mapping in the second sense on $[a, b]$, we have

$$\begin{aligned}
 & \left| \left(\frac{(x-a)^\alpha + (b-x)^\alpha}{b-a} \right) f(x) - \frac{\Gamma(\alpha+1)}{(b-a)} [J_{x-}^\alpha f(a) + J_{x+}^\alpha f(b)] \right| \\
 & \leq \frac{(x-a)^{\alpha+1}}{b-a} \int_0^1 t^\alpha |f'(tx + (1-t)a)| dt \\
 & \quad + \frac{(b-x)^{\alpha+1}}{b-a} \int_0^1 t^\alpha |f'(tx + (1-t)b)| dt \\
 & \leq \frac{(x-a)^{\alpha+1}}{b-a} \int_0^1 t^{\alpha+s} |f'(x)| + t^\alpha(1-t)^s |f'(a)| dt \\
 & \quad + \frac{(b-x)^{\alpha+1}}{b-a} \int_0^1 t^{\alpha+s} |f'(x)| + t^\alpha(1-t)^s |f'(b)| dt \\
 & \leq \frac{M}{b-a} \left(\frac{1}{\alpha+s+1} + \frac{\Gamma(\alpha+1)\Gamma(s+1)}{\Gamma(\alpha+s+2)} \right) [(x-a)^{\alpha+1} + (b-x)^{\alpha+1}]
 \end{aligned}$$

where we have used the fact that

$$\int_0^1 t^{\alpha+s} dt = \frac{1}{\alpha+s+1} \quad \text{and} \quad \int_0^1 t^\alpha(1-t)^s dt = \frac{\Gamma(\alpha+1)\Gamma(s+1)}{\Gamma(\alpha+s+2)}.$$

Hence, using the reduction formula $\Gamma(n+1) = n\Gamma(n)$ ($n > 0$) for Euler Gamma function, the proof is complete. \square

Remark 1. In Theorem 7, if we choose $\alpha = 1$, then (2.6) reduces the inequality (1.3) of Theorem 3.

Theorem 8. Let $f : [a, b] \subset [0, \infty) \rightarrow \mathbb{R}$, be a differentiable mapping on (a, b) with $a < b$ such that $f' \in L[a, b]$. If $|f'|^q$ is s -convex in the second sense on $[a, b]$ for some fixed $s \in (0, 1]$, $p, q > 1$ and $|f'(x)| \leq M$, $x \in [a, b]$, then the following inequality for fractional integrals holds:

$$\begin{aligned}
 & \left| \left(\frac{(x-a)^\alpha + (b-x)^\alpha}{b-a} \right) f(x) - \frac{\Gamma(\alpha+1)}{(b-a)} [J_{x-}^\alpha f(a) + J_{x+}^\alpha f(b)] \right| \\
 (2.7) \quad & \leq \frac{M}{(1+p\alpha)^{\frac{1}{p}}} \left(\frac{2}{s+1} \right)^{\frac{1}{q}} \left[\frac{(x-a)^{\alpha+1} + (b-x)^{\alpha+1}}{b-a} \right],
 \end{aligned}$$

where $\frac{1}{p} + \frac{1}{q} = 1$, $\alpha > 0$ and Γ is Euler Gamma function.

Proof. From Lemma 2 and using the well known Hölder inequality , we have

$$\begin{aligned}
& \left| \left(\frac{(x-a)^\alpha + (b-x)^\alpha}{b-a} \right) f(x) - \frac{\Gamma(\alpha+1)}{(b-a)} [J_{x-}^\alpha f(a) + J_{x+}^\alpha f(b)] \right| \\
& \leq \frac{(x-a)^{\alpha+1}}{b-a} \int_0^1 t^\alpha |f'(tx + (1-t)a)| dt \\
& \quad + \frac{(b-x)^{\alpha+1}}{b-a} \int_0^1 t^\alpha |f'(tx + (1-t)b)| dt \\
& \leq \frac{(x-a)^{\alpha+1}}{b-a} \left(\int_0^1 t^{p\alpha} dt \right)^{\frac{1}{p}} \left(\int_0^1 |f'(tx + (1-t)a)|^q dt \right)^{\frac{1}{q}} \\
& \quad + \frac{(b-x)^{\alpha+1}}{b-a} \left(\int_0^1 t^{p\alpha} dt \right)^{\frac{1}{p}} \left(\int_0^1 |f'(tx + (1-t)b)|^q dt \right)^{\frac{1}{q}}.
\end{aligned}$$

Since $|f'|^q$ is s -convex in the second sense on $[a, b]$ and $|f'(x)| \leq M$, we get (see [13, p. 1073])

$$\int_0^1 |f'(tx + (1-t)a)|^q dt \leq \frac{2M^q}{s+1} \quad \text{and} \quad \int_0^1 |f'(tx + (1-t)b)|^q dt \leq \frac{2M^q}{s+1}$$

and by simple computation

$$\int_0^1 t^{p\alpha} dt = \frac{1}{p\alpha + 1}.$$

Hence, we have

$$\begin{aligned}
& \left| \left(\frac{(x-a)^\alpha + (b-x)^\alpha}{b-a} \right) f(x) - \frac{\Gamma(\alpha+1)}{(b-a)} [J_{x-}^\alpha f(a) + J_{x+}^\alpha f(b)] \right| \\
& \leq \frac{M}{(1+p\alpha)^{\frac{1}{p}}} \left(\frac{2}{s+1} \right)^{\frac{1}{q}} \left[\frac{(x-a)^{\alpha+1} + (b-x)^{\alpha+1}}{b-a} \right]
\end{aligned}$$

which completes the proof. \square

Remark 2. In Theorem 8, if we choose $\alpha = 1$, then (2.7) reduces the inequality (1.4) of Theorem 4.

Theorem 9. Let $f : [a, b] \subset [0, \infty) \rightarrow \mathbb{R}$, be a differentiable mapping on (a, b) with $a < b$ such that $f' \in L[a, b]$. If $|f'|^q$ is s -convex in the second sense on $[a, b]$ for some fixed $s \in (0, 1]$, $q \geq 1$, and $|f'(x)| \leq M$, $x \in [a, b]$, then the following inequality for fractional integrals holds:

$$\begin{aligned}
(2.8) \quad & \left| \left(\frac{(x-a)^\alpha + (b-x)^\alpha}{b-a} \right) f(x) - \frac{\Gamma(\alpha+1)}{(b-a)} [J_{x-}^\alpha f(a) + J_{x+}^\alpha f(b)] \right| \\
& \leq \frac{M}{(1+p\alpha)^{\frac{1}{p}}} \left(\frac{2}{s+1} \right)^{\frac{1}{q}} \left[\frac{(x-a)^{\alpha+1} + (b-x)^{\alpha+1}}{b-a} \right],
\end{aligned}$$

where $\alpha > 0$ and Γ is Euler Gamma function.

Proof. From Lemma 2 and using the well known power mean inequality, we have

$$\begin{aligned}
 & \left| \left(\frac{(x-a)^\alpha + (b-x)^\alpha}{b-a} \right) f(x) - \frac{\Gamma(\alpha+1)}{(b-a)} [J_{x-}^\alpha f(a) + J_{x+}^\alpha f(b)] \right| \\
 & \leq \frac{(x-a)^{\alpha+1}}{b-a} \int_0^1 t^\alpha |f'(tx + (1-t)a)| dt \\
 & \quad + \frac{(b-x)^{\alpha+1}}{b-a} \int_0^1 t^\alpha |f'(tx + (1-t)b)| dt \\
 & \leq \frac{(x-a)^{\alpha+1}}{b-a} \left(\int_0^1 t^\alpha dt \right)^{1-\frac{1}{q}} \left(\int_0^1 t^\alpha |f'(tx + (1-t)a)|^q dt \right)^{\frac{1}{q}} \\
 & \quad + \frac{(b-x)^{\alpha+1}}{b-a} \left(\int_0^1 t^\alpha dt \right)^{1-\frac{1}{q}} \left(\int_0^1 t^\alpha |f'(tx + (1-t)b)|^q dt \right)^{\frac{1}{q}}.
 \end{aligned}$$

Since $|f'|^q$ is s -convex in the second sense on $[a, b]$ and $|f'(x)| \leq M$, we get (see [13, p. 1073])

$$\begin{aligned}
 \int_0^1 t^\alpha |f'(tx + (1-t)a)|^q dt & \leq \int_0^1 [t^{s+\alpha} |f'(x)|^q + t^\alpha (1-t)^s |f'(a)|^q] dt \\
 & = \frac{|f'(x)|^q}{\alpha+s+1} + |f'(a)|^q \int_0^1 t^\alpha (1-t)^s dt \\
 & = \frac{|f'(x)|^q}{\alpha+s+1} + |f'(a)|^q \beta(\alpha+1, s+1) \\
 & = \frac{|f'(x)|^q}{\alpha+s+1} + |f'(a)|^q \frac{\Gamma(\alpha+1)\Gamma(s+1)}{(\alpha+s+1)\Gamma(\alpha+s+1)} \\
 & \leq \frac{M^q}{\alpha+s+1} \left(1 + \frac{\Gamma(\alpha+1)\Gamma(s+1)}{\Gamma(\alpha+s+1)} \right)
 \end{aligned}$$

and similarly

$$\begin{aligned}
 \int_0^1 t^\alpha |f'(tx + (1-t)b)|^q dt & \leq \int_0^1 [t^{s+\alpha} |f'(x)|^q + t^\alpha (1-t)^s |f'(b)|^q] dt \\
 & = \frac{|f'(x)|^q}{\alpha+s+1} + |f'(b)|^q \int_0^1 t^\alpha (1-t)^s dt \\
 & = \frac{|f'(x)|^q}{\alpha+s+1} + |f'(b)|^q \beta(\alpha+1, s+1) \\
 & = \frac{|f'(x)|^q}{\alpha+s+1} + |f'(b)|^q \frac{\Gamma(\alpha+1)\Gamma(s+1)}{(\alpha+s+1)\Gamma(\alpha+s+1)} \\
 & \leq \frac{M^q}{\alpha+s+1} \left(1 + \frac{\Gamma(\alpha+1)\Gamma(s+1)}{\Gamma(\alpha+s+1)} \right),
 \end{aligned}$$

where β is Euler Beta function defined by

$$\beta(x, y) = \int_0^1 t^{x-1}(1-t)^{y-1} dt \quad (x, y > 0).$$

We used the fact that

$$\beta(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)} \quad \text{and} \quad \Gamma(n+1) = n\Gamma(n) (n > 0).$$

Hence, we have

$$\begin{aligned} & \left| \left(\frac{(x-a)^\alpha + (b-x)^\alpha}{b-a} \right) f(x) - \frac{\Gamma(\alpha+1)}{(b-a)} [J_{x-}^\alpha f(a) + J_{x+}^\alpha f(b)] \right| \\ & \leq M \left(\frac{1}{1+\alpha} \right)^{1-\frac{1}{q}} \left(\frac{1}{\alpha+s+1} \right)^{\frac{1}{q}} \\ & \quad \times \left(1 + \frac{\Gamma(\alpha+1)\Gamma(s+1)}{\Gamma(\alpha+s+1)} \right)^{\frac{1}{q}} \left[\frac{(x-a)^{\alpha+1} + (b-x)^{\alpha+1}}{b-a} \right] \end{aligned}$$

which completes the proof. \square

Remark 3. In Theorem 9, if we choose $\alpha = 1$, then (2.8) reduces the inequality (1.5) of Theorem 5.

The following result holds for s -concavity:

Theorem 10. Let $f : [a, b] \subset [0, \infty) \rightarrow \mathbb{R}$, be a differentiable mapping on (a, b) with $a < b$ such that $f' \in L[a, b]$. If $|f'|^q$ is s -concave in the second sense on $[a, b]$ for some fixed $s \in (0, 1]$ and $p, q > 1$, then the following inequality for fractional integrals holds:

$$\begin{aligned} & \left| \left(\frac{(x-a)^\alpha + (b-x)^\alpha}{b-a} \right) f(x) - \frac{\Gamma(\alpha+1)}{(b-a)} [J_{x-}^\alpha f(a) + J_{x+}^\alpha f(b)] \right| \\ (2.9) \quad & \leq \frac{2^{(s-1)/q}}{(1+p\alpha)^{\frac{1}{p}}(b-a)} \left[(x-a)^{\alpha+1} \left| f' \left(\frac{x+a}{2} \right) \right| + (b-x)^{\alpha+1} \left| f' \left(\frac{b+x}{2} \right) \right| \right], \end{aligned}$$

where $\frac{1}{p} + \frac{1}{q} = 1$, $\alpha > 0$ and Γ is Euler Gamma function.

Proof. From Lemma 2 and using the well known Hölder inequality , we have

$$\begin{aligned}
 & \left| \left(\frac{(x-a)^\alpha + (b-x)^\alpha}{b-a} \right) f(x) - \frac{\Gamma(\alpha+1)}{(b-a)} [J_{x-}^\alpha f(a) + J_{x+}^\alpha f(b)] \right| \\
 & \leq \frac{(x-a)^{\alpha+1}}{b-a} \int_0^1 t^\alpha |f'(tx + (1-t)a)| dt \\
 (2.10) \quad & + \frac{(b-x)^{\alpha+1}}{b-a} \int_0^1 t^\alpha |f'(tx + (1-t)b)| dt \\
 & \leq \frac{(x-a)^{\alpha+1}}{b-a} \left(\int_0^1 t^{p\alpha} dt \right)^{\frac{1}{p}} \left(\int_0^1 |f'(tx + (1-t)a)|^q dt \right)^{\frac{1}{q}} \\
 & \quad + \frac{(b-x)^{\alpha+1}}{b-a} \left(\int_0^1 t^{p\alpha} dt \right)^{\frac{1}{p}} \left(\int_0^1 |f'(tx + (1-t)b)|^q dt \right)^{\frac{1}{q}} .
 \end{aligned}$$

Since $|f'|^q$ is s -concave, using the inequality (1.2) we get (see [13, p. 1074])

$$(2.11) \quad \int_0^1 |f'(tx + (1-t)a)|^q dt \leq 2^{s-1} \left| f' \left(\frac{x+a}{2} \right) \right|^q$$

and

$$(2.12) \quad \int_0^1 |f'(tx + (1-t)b)|^q dt \leq 2^{s-1} \left| f' \left(\frac{b+x}{2} \right) \right|^q .$$

Using (2.11) and (2.12) in (2.10), we have

$$\begin{aligned}
 & \left| \left(\frac{(x-a)^\alpha + (b-x)^\alpha}{b-a} \right) f(x) - \frac{\Gamma(\alpha+1)}{(b-a)} [J_{x-}^\alpha f(a) + J_{x+}^\alpha f(b)] \right| \\
 & \leq \frac{2^{(s-1)/q}}{(1+p\alpha)^{\frac{1}{p}} (b-a)} \left[(x-a)^{\alpha+1} \left| f' \left(\frac{x+a}{2} \right) \right| + (b-x)^{\alpha+1} \left| f' \left(\frac{b+x}{2} \right) \right| \right]
 \end{aligned}$$

which completes the proof. □

Remark 4. In Theorem 10, if we choose $\alpha = 1$, then (2.9) reduces the inequality (1.6) of Theorem 6.

REFERENCES

- [1] S.S. Dragomir, On the Ostrowski's integral inequality for mappings with bounded variation and applications, *Math. Ineq. & Appl.*, 1(2) (1998).
- [2] S.S. Dragomir, The Ostrowski integral inequality for Lipschitzian mappings and applications, *Comput. Math. Appl.*, 38 (1999), 33-37.
- [3] S. S. Dragomir, S. Wang, A new inequality of Ostrowski's type in L_1 -norm and applications to some special means and to some numerical quadrature rules, *Tamkang J. of Math.*, 28 (1997), 239-244.
- [4] Z. Liu, Some companions of an Ostrowski type inequality and application, *J. Inequal. in Pure and Appl. Math.*, 10(2), 2009, Art. 52, 12 pp.
- [5] M.E. Özdemir, H. Kavurmacı and E. Set, Ostrowski's type inequalities for (α, m) -convex functions, *Kyungpook Math. J.*, 50 (2010), 371-378.
- [6] B. G. Pachpatte, On an inequality of Ostrowski type in three independent variables, *J. Math. Anal. Appl.*, 249(2000), 583-591.
- [7] B. G. Pachpatte, On a new Ostrowski type inequality in two independent variables, *Tamkang J. Math.*, 32(1), (2001), 45-49

- [8] A. Rafiq, N.A. Mir and F. Ahmad, Weighted Čebyšev-Ostrowski type inequalities, Applied Math. Mechanics (English Edition), 2007, 28(7), 901-906.
- [9] M. Z. Sarikaya, On the Ostrowski type integral inequality, Acta Math. Univ. Comenianae, Vol. LXXIX, 1(2010), pp. 129-134.
- [10] N. Ujević, Sharp inequalities of Simpson type and Ostrowski type, Comput. Math. Appl. 48 (2004) 145-151.
- [11] L. Zhongxue, On sharp inequalities of Simpson type and Ostrowski type in two independent variables, Comput. Math. Appl., 56 (2008) 2043-2047.
- [12] M. Alomari and M. Darus, Some Ostrowski type inequalities for convex functions with applications, RGMIA 13 (1) (2010) article No. 3. Preprint.
- [13] M. Alomari, M. Darus, S.S. Dragomir, P. Cerone, Ostrowski type inequalities for functions whose derivatives are s -convex in the second sense, Appl. Math. Lett. 23 (2010) 1071-1076.
- [14] S. S. Dragomir and S. Fitzpatrick, The Hadamard's inequality for s -convex functions in the second sense, Demonstratio Math. 32(4), (1999), 687-696.
- [15] H. Hudzik and L. Maligranda, Some remarks on s -convex functions, Aequationes Math. 48 (1994), 100-111.
- [16] U.S. Kirmaci, M.K. Bakula, M.E. Özdemir, J. Pečarić, Hadamard-type inequalities for s -convex functions, Appl. Math. and Comp., 193 (2007), 26-35.
- [17] M.Z. Sarikaya, E. Set and M.E. Özdemir "On new inequalities of Simpson's type for s -convex functions", Comput. Math. Appl., 60(8), (2010) 2191-2199.
- [18] G. Anastassiou, M.R. Hooshmandasl, A. Ghasemi and F. Moftakharzadeh, Montgomery identities for fractional integrals and related fractional inequalities, J. Ineq. Pure and Appl. Math., 10(4) (2009), Art. 97.
- [19] S. Belarbi and Z. Dahmani, On some new fractional integral inequalities, J. Ineq. Pure and Appl. Math., 10(3) (2009), Art. 86.
- [20] Z. Dahmani, New inequalities in fractional integrals, International Journal of Nonlinear Science, 9(4) (2010), 493-497.
- [21] Z. Dahmani, On Minkowski and Hermite-Hadamard integral inequalities via fractional integration, Ann. Funct. Anal. 1(1) (2010), 51-58.
- [22] Z. Dahmani, L. Tabharit, S. Taf, Some fractional integral inequalities, Nonl. Sci. Lett. A, 1(2) (2010), 155-160.
- [23] Z. Dahmani, L. Tabharit, S. Taf, New generalizations of Grüss inequality using Riemann-Liouville fractional integrals, Bull. Math. Anal. Appl., 2(3) (2010), 93-99.
- [24] R. Gorenflo, F. Mainardi, Fractional calculus: integral and differential equations of fractional order, Springer Verlag, Wien (1997), 223-276.
- [25] S. Miller and B. Ross, An introduction to the Fractional Calculus and Fractional Differential Equations, John Wiley & Sons, USA, 1993, p.2.
- [26] I. Podlubni, Fractional Differential Equations, Academic Press, San Diego, 1999.
- [27] M.Z. Sarikaya, E. Set, H. Yıldız and N. Başak, Hermite-Hadamard's inequalities for fractional integrals and related fractional inequalities, Submitted.
- [28] M.Z. Sarikaya and H. Ogunmez, On new inequalities via Riemann-Liouville fractional integration, arXiv:1005.1167v1, submitted.
- [29] D.S. Mitrinović, J.E. Pečarić and A.M. Fink, Inequalities Involving Functions and Their Integrals and Derivatives, Kluwer Academic Publishers, Dordrecht, 1991
- [30] A.M. Ostrowski, Über die Absolutabweichung einer differentierbaren Funktion von ihren Integralmittelwert, Comment. Math. Helv., 10, 226-227, (1938).
- [31] D.S. Mitrinović, J.E. Pečarić and A.M. Fink, Classical and New Inequalities in Analysis, Kluwer Academic Publishers, Dordrecht, 1993, p. 106.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, DÜZCE-TURKEY

E-mail address: erhanset@yahoo.com