ON DISCRETIZATION OF TORI OF COMPACT SIMPLE LIE GROUPS II

JIŘÍ HRIVNÁK¹, LENKA MOTLOCHOVÁ³, AND JIŘÍ PATERA^{2,3,4}

ABSTRACT. The discrete orthogonality of special function families, called C- and S-functions, which are derived from the characters of compact simple Lie groups, is described in [3]. Here, the results of [3] are extended to two additional recently discovered families of special functions, called S^s – and S^l -functions. The main result is an explicit description of their pairwise discrete orthogonality within each family, when the functions are sampled on finite fragments F_M^s and F_M^l of a lattice in any dimension $n \geq 2$ and of any density controlled by M, and of the symmetry of the weight lattice of any compact simple Lie group with two different lengths of roots.

¹ Department of physics, Faculty of nuclear sciences and physical engineering, Czech Technical University in Prague, Břehová 7, CZ-115 19 Prague, Czech Republic

² Centre de recherches mathématiques, Université de Montréal, C. P. 6128 – Centre ville, Montréal, H3C 3J7, Québec, Canada

³ Département de Mathématiques et de Statistique, Université de Montréal, Québec, Canada;

⁴ MIND Research Institute, 3631 S. Harbor Blvd., Suite 200, Santa Ana, CA 92704, USA

E-mail: jiri.hrivnak@fjfi.cvut.cz, motlochova@dms.umontreal.ca, patera@crm.umontreal.ca

1. INTRODUCTION

This paper focuses on the Fourier transform of data sampled on lattices of any dimension and any symmetry [13, 15]. The main problem is to find families of expansion functions that are complete in their space and orthogonal over finite fragments of the lattices. Generality of results is possible because the expansion function is built using properties that are uniformly valid over the series of semisimple Lie groups. Results of [3] on the discrete orthogonality of C- and S-functions of compact simple Lie groups are extended to the recently discovered families of S^s- and S^l- functions. The new families of functions add new possibilities of transforms for the same data.

Uniform discretization of tori of all semisimple Lie groups became possible after the classification of conjugacy classes of elements of finite order in compact simple Lie groups [6]. This was accomplished in [11, 12] for C-functions and extended to S-functions in [13]. Functions of C- and S-families are ingredients of irreducible characters of representations. They are uniformly defined for all semisimple Lie groups. Discretization here refers to their orthogonality when sampled on a fraction of a lattice F_M in the fundamental region F of the corresponding Lie group and summed up over all lattice points in F_M . This lattice is necessarily isomorphic to the weight lattice of the underlying Lie group, but its density is controlled by the choice of $M \in \mathbb{N}$.

C-functions are Weyl group invariant constituents of characters of irreducible representations. They are well known, even if unfrequently used [7]. S-functions appear in the Weyl character formula. They are skew-invariant with respect to the Weyl group [8]. In the new families of S^s - and S^l -functions, the Weyl group acts differently when reflections are, with respect to hyperplanes, orthogonal to short and long roots of the Lie group. The functions are 'half invariant and half skew-invariant' under the action of the Weyl group.

The key point of the discretization of S^s - and S^l -functions lies in finding appropriate subsets $F_M^s \subset F_M$ and $F_M^l \subset F_M$, which play the role of sampling points of a given data. The solution involves determining the sets of weights Λ_M^s and Λ_M^l , which label the discretely orthogonal S^s - and S^l -functions over the sets F_M^s and F_M^l . In order to verify the completeness of the found sets of functions, the last step involves comparing the number of points in F_M^s , F_M^l to the number of weights in Λ_M^s , Λ_M^l .

Date: June 4, 2012.

The pertinent standard properties of affine Weyl groups and their dual versions are recalled in section 2. Two types of sign homomorphisms and the corresponding fundamental domains are defined in section 3. The S^s – and S^l –functions and their behavior on the given discrete grids are studied in section 4. In section 5, the number of points in F_M^s , F_M^l are shown to be equal to the number of weights in Λ_M^s , Λ_M^l . Explicit formulas for these numbers are also given. Section 6 contains the detailed description of the discrete orthogonality and discrete transforms of S^{s} – and S^{l} –functions. Comments and follow-up questions are in the last section.

2. Pertinent properties of Affine Weyl groups

2.1. Roots and reflections.

We use the notation established in [3]. Recall that, to the Lie algebra of the compact simple Lie group G of rank n, corresponds the set of simple roots $\Delta = \{\alpha_1, \ldots, \alpha_n\}$ [1, 5, 17]. The set Δ spans the Euclidean space \mathbb{R}^n , with the scalar product denoted by \langle , \rangle . We consider here only simple algebras with two different lengths of roots, namely B_n , $n \ge 3$, C_n , $n \ge 2$, G_2 and F_4 . For these algebras, the set of simple roots consists of short simple roots Δ_s and long simple roots Δ_l . Thus, we have the disjoint decomposition

$$\Delta = \Delta_s \cup \Delta_l. \tag{1}$$

We then use the following well-known objects related to the set Δ :

• The marks m_1, \ldots, m_n of the highest root $\xi \equiv -\alpha_0 = m_1 \alpha_1 + \cdots + m_n \alpha_n$.

c

- The Coxeter number $m = 1 + m_1 + \dots + m_n$ of G.
- The Cartan matrix C and its determinant

$$= \det C. \tag{2}$$

- The root lattice $Q = \mathbb{Z}\alpha_1 + \cdots + \mathbb{Z}\alpha_n$.
- The \mathbb{Z} -dual lattice to Q,

$$P^{\vee} = \left\{ \omega^{\vee} \in \mathbb{R}^n \, | \, \langle \omega^{\vee}, \, \alpha \rangle \in \mathbb{Z}, \, \forall \alpha \in \Delta \right\} = \mathbb{Z} \omega_1^{\vee} + \dots + \mathbb{Z} \omega_n^{\vee} \, .$$

- The dual root lattice Q[∨] = Zα₁[∨] + ··· + Zα_n[∨], where α_i[∨] = 2α_i/⟨α_i, α_i⟩.
 The dual marks m₁[∨],..., m_n[∨] of the highest dual root η ≡ -α₀[∨] = m₁[∨]α₁[∨] + ··· + m_n[∨]α_n[∨]. The marks and the dual marks are summarized in Table 1 in [3].
- The \mathbb{Z} -dual lattice to Q^{\vee}

$$P = \left\{ \omega \in \mathbb{R}^n \, | \, \langle \omega, \, \alpha^{\vee} \rangle \in \mathbb{Z}, \, \forall \alpha^{\vee} \in Q^{\vee} \right\} = \mathbb{Z}\omega_1 + \dots + \mathbb{Z}\omega_n.$$

Recall that n reflections r_{α} , $\alpha \in \Delta$ in (n-1)-dimensional 'mirrors' orthogonal to simple roots intersecting at the origin are given explicitly by

$$r_{\alpha}a = a - \frac{2\langle a, \alpha \rangle}{\langle \alpha, \alpha \rangle} \alpha, \qquad a \in \mathbb{R}^n$$

The affine reflection r_0 with respect to the highest root ξ is given by

$$r_0 a = r_{\xi} a + \frac{2\xi}{\langle \xi, \xi \rangle}, \qquad r_{\xi} a = a - \frac{2\langle a, \xi \rangle}{\langle \xi, \xi \rangle} \xi, \qquad a \in \mathbb{R}^n.$$

We denote the set of reflections $r_1 \equiv r_{\alpha_1}, \ldots, r_n \equiv r_{\alpha_n}$, together with the affine reflection r_0 , by R, i.e.

$$R = \{r_0, r_1, \dots, r_n\}.$$
 (3)

Analogously to (1), we divide the reflections of R into two subsets

$$R^{s} = \{r_{\alpha} \mid \alpha \in \Delta_{s}\}$$
$$R^{l} = \{r_{\alpha} \mid \alpha \in \Delta_{l}\} \cup \{r_{0}\}$$

We then obtain the disjoint decomposition

$$R = R^s \cup R^l. \tag{4}$$

Type	R^{s}	R^l	$R^{s\vee}$	$R^{l\vee}$	m^s	m^l
$B_n \ (n \ge 3)$	r_n	$r_0, r_1, \ldots, r_{n-1}$	r_0^{\lor}, r_n^{\lor}	$r_1^{\vee},\ldots,r_{n-1}^{\vee}$	2	2n-2
$C_n \ (n \ge 2)$	r_1,\ldots,r_{n-1}	r_0, r_n	$r_0^{\vee}, r_1^{\vee}, \dots, r_{n-1}^{\vee}$	r_n^{\vee}	2n-2	2
G_2	r_2	r_0, r_1	r_0^{\lor}, r_2^{\lor}	r_1^{\vee}	3	3
F_4	r_3, r_4	r_0, r_1, r_2	r_0^ee, r_3^ee, r_4^ee	r_1^{\lor}, r_2^{\lor}	6	6

TABLE 1. The decomposition of the sets of generators R, R^{\vee} and the Coxeter numbers m^s , m^l . Numbering of the simple roots is standard (see e.g. Figure 1 in [3]).

We can also call the sums of marks corresponding to the long or short roots, i.e. the numbers

$$m^{s} = \sum_{\alpha_{i} \in \Delta_{s}} m_{i}$$
$$m^{l} = \sum_{\alpha_{i} \in \Delta_{l}} m_{i} + 1,$$

the short and the long Coxeter numbers. Their sum gives the Coxeter number $m = m^s + m^l$.

The dual affine reflection r_0^{\vee} , with respect to the dual highest root η , is given by

$$r_0^{\vee}a = r_\eta a + \frac{2\eta}{\langle \eta, \eta \rangle}, \quad r_\eta a = a - \frac{2\langle a, \eta \rangle}{\langle \eta, \eta \rangle} \eta, \quad a \in \mathbb{R}^n.$$

We denote the set of reflections $r_1^{\vee} \equiv r_{\alpha_1}, \ldots, r_n^{\vee} \equiv r_{\alpha_n}$, together with the dual affine reflection r_0^{\vee} , by R^{\vee} , i.e.

$$R^{\vee} = \{r_0^{\vee}, r_1^{\vee}, \dots, r_n^{\vee}\}.$$
 (5)

Analogously to (1), (4), we divide the reflections of R^{\vee} into two subsets

$$R^{s\vee} = \{r_{\alpha} \mid \alpha \in \Delta_s\} \cup \{r_0^{\vee}\}$$
$$R^{l\vee} = \{r_{\alpha} \mid \alpha \in \Delta_l\}.$$

The disjoint decomposition of R^{\vee} is then

$$R^{\vee} = R^{s} \vee \cup R^{l} \vee. \tag{6}$$

We can also call the sums of the dual marks corresponding to generators from $R^{s\vee}$, $R^{l\vee}$ i.e. the numbers

$$\begin{split} m^{s\vee} &= \sum_{\alpha_i \in \Delta_s} m_i^{\vee} + 1 \\ m^{l\vee} &= \sum_{\alpha_i \in \Delta_l} m_i^{\vee}, \end{split}$$

the short and the long dual Coxeter numbers. Again, their sum gives the Coxeter number $m = m^{s\vee} + m^{l\vee}$. Direct calculation of these numbers yields the following crucial result:

Proposition 2.1. For the numbers m^s , m^l and $m^{s\vee}$, $m^{l\vee}$, it holds that

$$m^s = m^{s\vee}, \quad m^l = m^{l\vee}. \tag{7}$$

The explicit form of decompositions (4) and (6) of sets R and R^{\vee} as well as Coxeter numbers m^s , m^l is given in Table 1.

2.2. Weyl group and affine Weyl group.

Weyl group W is generated by n reflections r_{α} , $\alpha \in \Delta$. Applying the action of W on the set of simple roots Δ , we obtain the entire root system $W\Delta$. The root system $W\Delta$ contains two subsystems $W\Delta_s$ and $W\Delta_l$, i.e. we have the disjoint decomposition

$$W\Delta = W\Delta_s \cup W\Delta_l. \tag{8}$$

The set of n + 1 generators R generates the affine Weyl group W^{aff} . The affine Weyl group W^{aff} can be viewed as the semidirect product of the Abelian group of translations Q^{\vee} and of the Weyl group W:

$$W^{\text{aff}} = Q^{\vee} \rtimes W. \tag{9}$$

Thus, for any $w^{\text{aff}} \in W^{\text{aff}}$, there exist a unique $w \in W$ and a unique shift $T(q^{\vee})$ such that $w^{\text{aff}} = T(q^{\vee})w$. The retraction homomorphism $\psi: W^{\text{aff}} \to W$ for $w^{\text{aff}} \in W^{\text{aff}}$ is given by

$$\psi(w^{\text{aff}}) = \psi(T(q^{\vee})w) = w.$$
(10)

The fundamental domain F of W^{aff} , which consists of precisely one point of each W^{aff} -orbit, is the convex hull of the points $\left\{0, \frac{\omega_1^{\vee}}{m_1}, \ldots, \frac{\omega_n^{\vee}}{m_n}\right\}$. Considering n+1 real parameters $y_0, \ldots, y_n \ge 0$, we have

$$F = \left\{ y_1 \omega_1^{\vee} + \dots + y_n \omega_n^{\vee} \mid y_0 + y_1 m_1 + \dots + y_n m_n = 1 \right\}.$$
 (11)

Recall that the stabilizer

$$\operatorname{Stab}_{W^{\operatorname{aff}}}(a) = \left\{ w^{\operatorname{aff}} \in W^{\operatorname{aff}} \mid w^{\operatorname{aff}}a = a \right\}$$
(12)

of a point $a = y_1 \omega_1^{\vee} + \cdots + y_n \omega_n^{\vee} \in F$ is trivial, $\operatorname{Stab}_{W^{\operatorname{aff}}}(a) = 1$ if the point a is in the interior of F, $a \in \operatorname{int}(F)$. Otherwise the group $\operatorname{Stab}_{W^{\operatorname{aff}}}(a)$ is generated by such r_i for which $y_i = 0, i = 0, \ldots, n$.

Considering the standard action of W on the torus \mathbb{R}^n/Q^{\vee} , we denote for $x \in \mathbb{R}^n/Q^{\vee}$ the isotropy group and its order by

$$\operatorname{Stab}(x) = \{ w \in W \mid wx = x \}, \quad h_x \equiv |\operatorname{Stab}(x)|$$

and denote the orbit and its order by

$$Wx = \left\{ wx \in \mathbb{R}^n / Q^{\vee} \, | \, w \in W \right\}, \quad \varepsilon(x) \equiv |Wx|.$$

Then we have

$$\varepsilon(x) = \frac{|W|}{h_x}.$$
(13)

Recall the following three properties from Proposition 2.2 in [3] of the action of W on the torus \mathbb{R}^n/Q^{\vee} :

(1) For any $x \in \mathbb{R}^n/Q^{\vee}$, there exists $x' \in F \cap \mathbb{R}^n/Q^{\vee}$ and $w \in W$ such that

$$x = wx'. (14)$$

(2) If
$$x, x' \in F \cap \mathbb{R}^n/Q^{\vee}$$
 and $x' = wx, w \in W$, then

$$x' = x = wx. (15)$$

(3) If
$$x \in F \cap \mathbb{R}^n / Q^{\vee}$$
, i.e. $x = a + Q^{\vee}$, $a \in F$, then $\psi(\operatorname{Stab}_{W^{\operatorname{aff}}}(a)) = \operatorname{Stab}(x)$ and
 $\operatorname{Stab}(x) \cong \operatorname{Stab}_{W^{\operatorname{aff}}}(a).$
(16)

2.3. Dual affine Weyl group.

The dual affine Weyl group \widehat{W}^{aff} is generated by the set R^{\vee} . Moreover, \widehat{W}^{aff} is a semidirect product of the group of shifts Q and the Weyl group W

$$\widehat{W}^{\text{aff}} = Q \rtimes W. \tag{17}$$

Thus, for any $w^{\text{aff}} \in \widehat{W}^{\text{aff}}$, there exist a unique $w \in W$ and a unique shift T(q) such that $w^{\text{aff}} = T(q)w$. The dual retraction homomorphism $\widehat{\psi} : \widehat{W}^{\text{aff}} \to W$ for $w^{\text{aff}} \in \widehat{W}^{\text{aff}}$ is given by

$$\widehat{\psi}(w^{\text{aff}}) = \widehat{\psi}(T(q)w) = w.$$
(18)

The dual fundamental domain F^{\vee} of \widehat{W}^{aff} is the convex hull of vertices $\left\{0, \frac{\omega_1}{m_1^{\vee}}, \ldots, \frac{\omega_n}{m_n^{\vee}}\right\}$. Considering n+1 real parameters $z_0, \ldots, z_n \ge 0$, we have

$$F^{\vee} = \left\{ z_1 \omega_1 + \dots + z_n \omega_n \mid z_0 + z_1 m_1^{\vee} + \dots + z_n m_n^{\vee} = 1 \right\}.$$
 (19)

Consider the point $a = z_1 \omega_1 + \cdots + z_n \omega_n \in F^{\vee}$ such that $z_0 + z_1 m_1^{\vee} + \cdots + z_n m_n^{\vee} = 1$. The isotropy group

$$\operatorname{Stab}_{\widehat{W}^{\operatorname{aff}}}(a) = \left\{ w^{\operatorname{aff}} \in \widehat{W}^{\operatorname{aff}} \mid w^{\operatorname{aff}}a = a \right\}$$
(20)

of point a is trivial, $\operatorname{Stab}_{\widehat{W}^{\operatorname{aff}}}(a) = 1$, if $a \in \operatorname{int}(F^{\vee})$, i.e. all $z_i > 0, i = 0, \ldots, n$. Otherwise the group $\operatorname{Stab}_{\widehat{W}^{\operatorname{aff}}}(a)$ is generated by such r_i^{\vee} for which $z_i = 0, i = 0, \ldots, n$.

Recall from [3] that, for an arbitrary $M \in \mathbb{N}$, the grid Λ_M is defined as cosets from the *W*-invariant group P/MQ with a representative element in MF^{\vee} , i.e.

$$\Lambda_M \equiv MF^{\vee} \cap P/MQ.$$

Considering a natural action of W on the quotient group \mathbb{R}^n/MQ , we denote for $\lambda \in \mathbb{R}^n/MQ$ the isotropy group and its order by

$$\operatorname{Stab}^{\vee}(\lambda) = \{ w \in W \, | \, w\lambda = \lambda \} \,, \quad h_{\lambda}^{\vee} \equiv |\operatorname{Stab}^{\vee}(\lambda)|.$$

$$(21)$$

Recall the following three properties from Proposition 3.6 in [3] of the action of W on the quotient group \mathbb{R}^n/MQ .

(1) For any $\lambda \in P/MQ$, there exists $\lambda' \in \Lambda_M$ and $w \in W$ such that

$$\lambda = w\lambda'. \tag{22}$$

(2) If $\lambda, \lambda' \in \Lambda_M$ and $\lambda' = w\lambda, w \in W$, then

$$\lambda' = \lambda = w\lambda. \tag{23}$$

(3) If
$$\lambda \in MF^{\vee} \cap \mathbb{R}^n/MQ$$
, i.e. $\lambda = b + MQ$, $b \in MF^{\vee}$, then $\widehat{\psi}(\operatorname{Stab}_{\widehat{W}^{\operatorname{aff}}}(b/M)) = \operatorname{Stab}^{\vee}(\lambda)$ and

$$\operatorname{Stab}^{\vee}(\lambda) \cong \operatorname{Stab}_{\widehat{W}^{\operatorname{aff}}}(b/M).$$
 (24)

3. SIGN HOMOMORPHISMS AND ORBIT FUNCTIONS

3.1. Sign homomorphisms.

The Weyl group W has the following abstract presentation [1, 5]

$$r_i^2 = 1, \quad (r_i r_j)^{m_{ij}} = 1, \quad i, j = 1, \dots, n$$
 (25)

where integers m_{ij} are elements of the Coxeter matrix. To introduce various classes of orbit functions, we consider 'sign' homomorphisms $\sigma : W \to \{\pm 1\}$. An admissible mapping σ must satisfy the presentation condition (25)

$$\sigma(r_i)^2 = 1, \quad (\sigma(r_i)\sigma(r_j))^{m_{ij}} = 1, \quad i, j = 1, \dots, n.$$
 (26)

If condition (26) is satisfied, then it follows from the universality property (see e.g. [1]) that σ is a well-defined homomorphism and its values on any $w \in W$ are given as products of generator values. The following two choices of homomorphism values of generators r_{α} , $\alpha \in \Delta$, obviously satisfying (26), lead to the well-known homomorphisms:

$$\mathbf{1}(r_{\alpha}) = 1 \tag{27}$$

$$\sigma^e(r_\alpha) = -1 \tag{28}$$

which yield for any $w \in W$

$$\mathbf{1}(w) = 1 \tag{29}$$

$$\sigma^e(w) = \det w. \tag{30}$$

It is shown in [10] that, for root systems with two different lengths of roots, there are two other available choices. Using the decomposition (1), these two new homomorphisms are given as follows [10]:

$$\sigma^{s}(r_{\alpha}) = \begin{cases} 1, & \alpha \in \Delta_{l} \\ -1, & \alpha \in \Delta_{s} \end{cases}$$
(31)

$$\sigma^{l}(r_{\alpha}) = \begin{cases} 1, & \alpha \in \Delta_{s} \\ -1, & \alpha \in \Delta_{l}. \end{cases}$$
(32)

Since the highest root $\xi \in W\Delta_l$, there exist $w \in W$ and $\alpha \in \Delta_l$ such that $\xi = w\alpha$. Then from the relation $r_{\xi} = wr_{\alpha}w^{-1}$, we obtain for any sign homomorphism that $\sigma(r_{\xi}) = \sigma(r_{\alpha})$ holds. Thus we have

$$\sigma^{s}(r_{\xi}) = 1, \quad \sigma^{l}(r_{\xi}) = -1.$$
 (33)

Similarly, for the highest dual root η there exists a root $\beta \in W\Delta_s$ such that $\eta = 2\beta/\langle \beta, \beta \rangle$, and we obtain

$$\sigma^s(r_\eta) = -1, \quad \sigma^l(r_\eta) = 1. \tag{34}$$

3.2. Fundamental domains.

Each of the sign homomorphisms σ^s and σ^l determines a decomposition of the fundamental domain F. The factors of this decomposition will be crucial for the study of the orbit functions. We introduce two subsets of F:

$$F^{s} = \left\{ a \in F \mid \sigma^{s} \circ \psi \left(\operatorname{Stab}_{W^{\operatorname{aff}}}(a) \right) = \left\{ 1 \right\} \right\}$$
$$F^{l} = \left\{ a \in F \mid \sigma^{l} \circ \psi \left(\operatorname{Stab}_{W^{\operatorname{aff}}}(a) \right) = \left\{ 1 \right\} \right\}$$

where ψ is the retraction homomorphism (10). Since for all points of the interior of F the stabilizer is trivial, i.e. $\operatorname{Stab}_{W^{\operatorname{aff}}}(a) = 1$, $a \in \operatorname{int}(F)$, the interior $\operatorname{int}(F)$ is a subset of both F^s and F^l . In order to determine the analytic form of the sets F^s and F^l , we define two subsets of the boundaries of F

$$H^{s} = \{a \in F \mid (\exists r \in R^{s})(ra = a)\}$$
$$H^{l} = \left\{a \in F \mid (\exists r \in R^{l})(ra = a)\right\}.$$

Note that, since for the affine reflection $r_0 \in R^l$ it holds that $\psi(r_0) = r_{\xi}$, we have from (33) that $\sigma^s \circ \psi(r_0) = 1$ and $\sigma^l \circ \psi(r_0) = -1$. Taking into account the disjoint decomposition (4), we obtain for any $r \in R$ the following two exclusive choices:

$$\sigma^{s} \circ \psi(r) = -1, \quad \sigma^{l} \circ \psi(r) = 1, \qquad r \in \mathbb{R}^{s}$$

$$\sigma^{s} \circ \psi(r) = 1, \qquad \sigma^{l} \circ \psi(r) = -1, \quad r \in \mathbb{R}^{l}.$$
(35)

Proposition 3.1. For the sets F^s and F^l , the following holds:

- (1) $F^s = F \setminus H^s$.
- (2) $F^l = F \setminus H^l$.

Proof. Let $a \in F$.

- (1) If $a \notin F \setminus H^s$, then $a \in H^s$, and there exists $r \in R^s$ such that $r \in \operatorname{Stab}_{W^{\operatorname{aff}}}(a)$. Then according to (35), we have $\sigma^s \circ \psi(r) = -1$. Thus, $\sigma^s \circ \psi(\operatorname{Stab}_{W^{\operatorname{aff}}}(a)) = \{\pm 1\}$ and consequently $a \notin F^s$. Conversely, if $a \in F \setminus H^s$, then the stabilizer $\operatorname{Stab}_{W^{\operatorname{aff}}}(a)$ is either trivial or generated by generators from R^l only. Then, since for any generator $r \in R^l$ it follows from (35) that $\sigma^s \circ \psi(r) = 1$, we obtain $\sigma^s \circ \psi(\operatorname{Stab}_{W^{\operatorname{aff}}}(a)) = \{1\}$, i.e. $a \in F^s$.
- (2) This case is completely analogous to case (1).

The explicit description of domains F^s and F^l now follows from (11) and Proposition 3.1. We introduce the symbols y_i^s , $y_i^l \in \mathbb{R}$, i = 0, ..., n in the following way:

$$y_{i}^{s} > 0, \quad y_{i}^{l} \ge 0, \quad r_{i} \in \mathbb{R}^{s}$$

$$y_{i}^{s} \ge 0, \quad y_{i}^{l} > 0, \quad r_{i} \in \mathbb{R}^{l}.$$
(36)

Thus, the explicit form of F^s and F^l is given by

$$F^{s} = \left\{ y_{1}^{s} \omega_{1}^{\vee} + \dots + y_{n}^{s} \omega_{n}^{\vee} \mid y_{0}^{s} + y_{1}^{s} m_{1} + \dots + y_{n}^{s} m_{n} = 1 \right\}$$

$$F^{l} = \left\{ y_{1}^{l} \omega_{1}^{\vee} + \dots + y_{n}^{l} \omega_{n}^{\vee} \mid y_{0}^{l} + y_{1}^{l} m_{1} + \dots + y_{n}^{l} m_{n} = 1 \right\}.$$
(37)

3.3. Dual fundamental domains.

The sign homomorphisms σ^s and σ^l also determine a decomposition of the dual fundamental domain F^{\vee} . The factors of this decomposition will be needed for the study of the discretized orbit functions. We introduce two subsets of F^{\vee} :

$$F^{s\vee} = \left\{ a \in F^{\vee} \mid \sigma^{s} \circ \widehat{\psi} \left(\operatorname{Stab}_{\widehat{W}^{\operatorname{aff}}}(a) \right) = \{1\} \right\}$$

$$F^{l\vee} = \left\{ a \in F^{\vee} \mid \sigma^{l} \circ \widehat{\psi} \left(\operatorname{Stab}_{\widehat{W}^{\operatorname{aff}}}(a) \right) = \{1\} \right\}$$
(38)

where $\widehat{\psi}$ is the dual retraction homomorphism (18). Since for all points of the interior of F^{\vee} the stabilizer is trivial, i.e. $\operatorname{Stab}_{\widehat{W}^{\operatorname{aff}}}(a) = 1$, $a \in \operatorname{int}(F^{\vee})$, the interior $\operatorname{int}(F^{\vee})$ is a subset of both $F^{s^{\vee}}$ and $F^{l^{\vee}}$. In order to determine the analytic form of sets $F^{s\vee}$ and $F^{l\vee}$, we define two subsets of the boundaries of F^{\vee}

$$\begin{aligned} H^{s\vee} &= \left\{ a \in F^{\vee} \, | \, (\exists r \in R^{s\vee})(ra=a) \right\} \\ H^{l\vee} &= \left\{ a \in F^{\vee} \, | \, (\exists r \in R^{l\vee})(ra=a) \right\}. \end{aligned}$$

Note that, since for the affine reflection $r_0^{\vee} \in \mathbb{R}^{s^{\vee}}$ it holds that $\widehat{\psi}(r_0^{\vee}) = r_{\eta}$, we have from (34) that $\sigma^s \circ \widehat{\psi}(r_0^{\vee}) = -1$ and $\sigma^l \circ \widehat{\psi}(r_0^{\vee}) = 1$. Taking into account the disjoint decomposition (6), we obtain for any $r \in \mathbb{R}^{\vee}$ the following two exclusive choices:

$$\sigma^{s} \circ \widehat{\psi}(r) = -1, \quad \sigma^{l} \circ \widehat{\psi}(r) = 1, \qquad r \in R^{s \vee}$$

$$\sigma^{s} \circ \widehat{\psi}(r) = 1, \qquad \sigma^{l} \circ \widehat{\psi}(r) = -1, \quad r \in R^{l \vee}.$$
(39)

Similarly to Proposition 3.1, we obtain:

Proposition 3.2. For sets $F^{s\vee}$ and $F^{l\vee}$, the following holds:

- (1) $F^{s\vee} = F^{\vee} \setminus H^{s\vee}$ (2) $F^{l\vee} = F^{\vee} \setminus H^{l\vee}$.

The explicit description of domains $F^{s\vee}$ and $F^{l\vee}$ now follows from (19) and Proposition 3.2. We introduce the symbols $z_i^s, z_i^l \in \mathbb{R}, i = 0, \dots, n$ in the following way:

$$z_{i}^{s} > 0, \quad z_{i}^{l} \ge 0, \quad r_{i} \in \mathbb{R}^{s \vee}$$

$$z_{i}^{s} \ge 0, \quad z_{i}^{l} > 0, \quad r_{i} \in \mathbb{R}^{l \vee}.$$
(40)

Thus, the explicit form of $F^{s\vee}$ and $F^{l\vee}$ is given by

$$F^{s\vee} = \left\{ z_1^s \omega_1 + \dots + z_n^s \omega_n \mid z_0^s + z_1^s m_1^{\vee} + \dots + z_n^s m_n^{\vee} = 1 \right\}$$

$$F^{l\vee} = \left\{ z_1^l \omega_1 + \dots + z_n^l \omega_n \mid z_0^l + z_1^l m_1^{\vee} + \dots + z_n^l m_n^{\vee} = 1 \right\}.$$
(41)

4. S^s - AND S^l - FUNCTIONS

Four sign homomorphisms 1, σ^e , σ^s and σ^l induce four types of families of complex orbit functions. Within each family, determined by $\sigma \in \{1, \sigma^e, \sigma^s, \sigma^l\}$, are the complex functions $\varphi_b^{\sigma} : \mathbb{R}^n \to \mathbb{C}$ labeled by weights $b \in P$ and in the general form

$$\varphi_b^{\sigma}(a) = \sum_{w \in W} \sigma(w) \, e^{2\pi i \langle wb, a \rangle}, \quad a \in \mathbb{R}^n.$$
(42)

The resulting functions for $\sigma = 1$ in (42) are called *C*-functions; for their detailed review, see [8]. For $\sigma = \sigma^e$, we obtain the well-known *S*-functions [9]. The discretization properties of both *C*- and *S*-functions on a finite fragment of the grid $\frac{1}{M}P^{\vee}$ were described in [3]. The remaining two options of homomorphisms σ^s and σ^l and corresponding functions $\varphi_{\lambda}^{\sigma^s}$, $\varphi_{\lambda}^{\sigma^l}$, called S^l – and S^s -functions [10], were studied in detail for G_2 only [16]. In order to describe the discretization of functions $\varphi_{\lambda}^{\sigma^s}$ and $\varphi_{\lambda}^{\sigma^l}$ in full generality, we first review their basic properties.

4.1. S^s -functions.

4.1.1. Symmetries of S^s-functions. Choosing $\sigma = \sigma^s$ in (42), we obtain S^s-functions $\varphi_b^{\sigma^s}$; we abbreviate the notation by denoting $\varphi_b^s \equiv \varphi_b^{\sigma^s}$, i.e.

$$\varphi_b^s(a) = \sum_{w \in W} \sigma^s(w) \, e^{2\pi i \langle wb, a \rangle}, \quad a \in \mathbb{R}^n, \, b \in P.$$
(43)

The following properties of S^s -functions are crucial:

JIŘÍ HRIVNÁK¹, LENKA MOTLOCHOVÁ³, AND JIŘÍ PATERA^{2,3,4}

• (anti)symmetry with respect to $w \in W$

$$\varphi_b^s(wa) = \sigma^s(w)\varphi_b^s(a) \tag{44}$$

$$\varphi^s_{wb}(a) = \sigma^s(w)\varphi^s_b(a) \tag{45}$$

• invariance with respect to shifts from $q^{\vee} \in Q^{\vee}$

$$\varphi_b^s(a+q^{\vee}) = \varphi_b^s(a). \tag{46}$$

Thus, the S^s -functions are (anti)symmetric with respect to the affine Weyl group W^{aff} . This allows us to consider the values $\varphi_b^s(a)$ only for points of the fundamental domain $a \in F$. Moreover, from (35), (44) we deduce that

$$\varphi_b^s(ra) = -\varphi_b^s(a), \quad r \in \mathbb{R}^s.$$
(47)

This antisymmetry implies that the functions φ_b^s are for all $b \in P$ zero on part H^s of the boundary of F

$$\varphi_b^s(a') = 0, \quad a' \in H^s \tag{48}$$

and therefore we consider the functions φ_b^s on the fundamental domain $F^s = F \setminus H^s$ only.

4.1.2. Discretization of S^s -functions. In order to develop discrete calculus of S^s -functions, we investigate the behavior of these functions on the grid $\frac{1}{M}P^{\vee}$. Suppose we have fixed $M \in \mathbb{N}$ and $u \in \frac{1}{M}P^{\vee}$. It follows from (46) that we can consider φ_b^s as a function on cosets from $\frac{1}{M}P^{\vee}/Q^{\vee}$. It follows from (48) that we can consider φ_b^s only on the set

$$F_M^s \equiv \frac{1}{M} P^{\vee} / Q^{\vee} \cap F^s.$$
(49)

Next we have

$$\varphi_{b+MQ}^s(u) = \varphi_b^s(u), \quad u \in F_M^s$$

and thus we can consider the functions φ_{λ}^{s} on F_{M}^{s} parametrized by cosets from $\lambda \in P/MQ$. Moreover, it follows from (22) and (45) that we can consider φ_{λ}^{s} on F_{M}^{s} parameterized by classes from Λ_{M} . Taking any $\lambda \in \Lambda_{M}$ and any reflection $r^{\vee} \in R^{s^{\vee}}$ we calculate directly using (34), (45) and (46) that

$$\varphi^s_{Mr^\vee(\frac{\lambda}{M})}(u) = -\varphi^s_\lambda(u), \quad u \in F^s_M$$

This implies that, for $\lambda \in MH^{s\vee} \cap \Lambda_M$, the functions φ^s_{λ} are zero on F^s_M , i.e.

 $\varphi_{\lambda}^{s}(u) = 0, \quad \lambda \in MH^{s \vee} \cap \Lambda_{M}, \, u \in F_{M}^{s}.$

Defining the set

$$\Lambda^s_M \equiv P/MQ \cap MF^{s\vee} \tag{50}$$

we conclude that we can consider S^s -functions φ^s_{λ} on F^s_M parameterized by $\lambda \in \Lambda^s_M$ only.

4.2. S^l -functions.

4.2.1. Symmetries of S^l -functions. Choosing $\sigma = \sigma^l$ in (42), we obtain S^l -functions $\varphi_b^{\sigma^l}$; we abbreviate the notation by denoting $\varphi_b^l \equiv \varphi_b^{\sigma^l}$, i.e.

$$\varphi_b^l(a) = \sum_{w \in W} \sigma^l(w) \, e^{2\pi i \langle wb, a \rangle}, \quad a \in \mathbb{R}^n, \, b \in P.$$
(51)

The following properties of S^l -functions are crucial

• (anti)symmetry with respect to $w \in W$

$$\varphi_b^l(wa) = \sigma^l(w)\varphi_b^l(a) \tag{52}$$

$$\varphi_{wb}^l(a) = \sigma^l(w)\varphi_b^l(a) \tag{53}$$

• invariance with respect to shifts from $q^{\vee} \in Q^{\vee}$

$$\varphi_b^l(a+q^{\vee}) = \varphi_b^l(a). \tag{54}$$

Thus, the S^l -functions are (anti)symmetric with respect to the affine Weyl group W^{aff} . This allows us to consider the values $\varphi_b^l(a)$ only for points of the fundamental domain $a \in F$. Moreover, from (35), (52) and (54) we deduce that

$$\varphi_b^l(ra) = -\varphi_b^l(a), \quad r \in \mathbb{R}^l.$$
(55)

This antisymmetry implies that the functions φ_b^l are for all $b \in P$ zero on part H^l of the boundary of F

$$\varphi_b^l(a') = 0, \quad a' \in H^l \tag{56}$$

and therefore we consider the functions φ_b^l on the fundamental domain $F^l = F \setminus H^l$ only.

4.2.2. Discretization of S^l -functions. In order to develop discrete calculus of S^l -functions, we investigate the behavior of these functions on the grid $\frac{1}{M}P^{\vee}$. Suppose we have fixed $M \in \mathbb{N}$ and $u \in \frac{1}{M}P^{\vee}$. It follows from (54) that we can consider φ_b^l as a function on cosets from $\frac{1}{M}P^{\vee}/Q^{\vee}$. It follows from (56) that we can consider φ_b^l only on the set

$$F_M^l \equiv \frac{1}{M} P^{\vee} / Q^{\vee} \cap F^l.$$
(57)

Next we have

 $\varphi_{b+MQ}^l(u)=\varphi_b^l(u),\quad u\in F_M^l$

and thus we can consider the functions φ_{λ}^{l} on F_{M}^{l} parametrized by cosets from $\lambda \in P/MQ$. Moreover, it follows from (22) and (53) that we can consider φ_{λ}^{l} on F_{M}^{l} parameterized by classes from Λ_{M} . Taking any $\lambda \in \Lambda_{M}$ and any reflection $r^{\vee} \in \mathbb{R}^{l^{\vee}}$, we calculate directly using (34), (53) that

$$\varphi_{r^{\vee}\lambda}^{l}(u) = -\varphi_{\lambda}^{l}(u), \quad u \in F_{M}^{l}$$

This implies that for $\lambda \in MH^{l\vee} \cap \Lambda_M$, the functions φ_{λ}^l are zero on F_M^l , i.e.

$$\varphi_{\lambda}^{l}(u) = 0, \quad \lambda \in MH^{l \vee} \cap \Lambda_{M}, \, u \in F_{M}^{l}.$$

Defining the set

$$\Lambda^l_M \equiv P/MQ \cap MF^{l\vee} \tag{58}$$

we conclude that we can consider S^l -functions φ^l_{λ} on F^l_M parameterized by $\lambda \in \Lambda^l_M$ only.

5. Number of grid elements

5.1. Number of elements of F_M^s and F_M^l .

Recall from [3] that, for an arbitrary $M \in \mathbb{N}$, the grid F_M is given as cosets from the W-invariant group $\frac{1}{M}P^{\vee}/Q^{\vee}$ with a representative element in the fundamental domain F

$$F_M \equiv \frac{1}{M} P^{\vee} / Q^{\vee} \cap F$$

and the following property holds

$$WF_M = \frac{1}{M} P^{\vee} / Q^{\vee}.$$
(59)

The representative points of F_M can be explicitly written as

$$F_M = \left\{ \frac{u_1}{M} \omega_1^{\vee} + \dots + \frac{u_n}{M} \omega_n^{\vee} \mid u_0, u_1, \dots, u_n \in \mathbb{Z}^{\ge 0}, u_0 + u_1 m_1 + \dots + u_n m_n = M \right\}.$$
(60)

The number of elements of F_M , denoted by $|F_M|$, are also calculated in [3] for all simple Lie algebras. Using these results, we derive the number of elements of F_M^s and F_M^l . Firstly, we describe explicitly the sets F_M^s and F_M^l . Similarly to (36), we introduce the symbols u_i^s , $u_i^l \in \mathbb{R}$, $i = 0, \ldots, n$:

$$u_i^s \in \mathbb{N}, \qquad u_i^l \in \mathbb{Z}^{\ge 0}, \quad r_i \in R^s$$

$$u_i^s \in \mathbb{Z}^{\ge 0}, \quad u_i^l \in \mathbb{N}, \qquad r_i \in R^l.$$
 (61)

The explicit form of F_M^s and F_M^l then follows from the explicit form of F^s and F^l in (37):

$$F_M^s = \left\{ \frac{u_1^s}{M} \omega_1^{\vee} + \dots + \frac{u_n^s}{M} \omega_n^{\vee} \mid u_0^s + u_1^s m_1 + \dots + u_n^s m_n = M \right\}$$
(62)

$$F_{M}^{l} = \left\{ \frac{u_{1}^{l}}{M} \omega_{1}^{\vee} + \dots + \frac{u_{n}^{l}}{M} \omega_{n}^{\vee} \mid u_{0}^{l} + u_{1}^{l} m_{1} + \dots + u_{n}^{l} m_{n} = M \right\}.$$
 (63)

Using the following proposition, the number of elements of F_M^s and F_M^l can be obtained from the formulas for $|F_M|$.

Proposition 5.1. Let m^s and m^l be the short and long Coxeter numbers, respectively. Then

$$|F_M^s| = \begin{cases} 0 & M < m^s \\ 1 & M = m^s \\ |F_{M-m^s}| & M > m^s. \end{cases} \quad |F_M^l| = \begin{cases} 0 & M < m^l \\ 1 & M = m^l \\ |F_{M-m^l}| & M > m^l. \end{cases}$$
(64)

Proof. Taking non-negative numbers $u_i \in \mathbb{Z}^{\geq 0}$ and substituting the relations $u_i^s = 1 + u_i$ if $r_i \in \mathbb{R}^s$ and $u_i^s = u_i$ if $r_i \in \mathbb{R}^l$ into the defining relation (62), we obtain

$$u_0 + m_1 u_1 + \dots + m_n u_n = M - m^s, \quad u_0, \dots, u_n \in \mathbb{Z}^{\ge 0}.$$

This equation has one solution $[0, \ldots, 0]$ if $M = m^s$, no solution if $M < m^s$, and is equal to the defining relation (60) of F_{M-m^s} if $M > m^s$. The case of F_M^l is similar.

Theorem 5.2. The numbers of points of grids F_M^s and F_M^l of Lie algebras B_n , C_n , G_2 and F_4 are given by the following relations.

$$(1) C_n, n \ge 2,$$

$$|F_{2k}^{s}(C_{n})| = \binom{k+1}{n} + \binom{k}{n}$$
$$|F_{2k+1}^{s}(C_{n})| = 2\binom{k+1}{n}$$

$$|F_{2k}^{l}(C_{n})| = \binom{n+k-1}{n} + \binom{n+k-2}{n}$$
$$|F_{2k+1}^{l}(C_{n})| = 2\binom{n+k-1}{n}$$

(2) $B_n, n \ge 3,$

$$|F_M^s(B_n)| = |F_M^l(C_n)|$$
$$|F_M^l(B_n)| = |F_M^s(C_n)|$$

 $(3) G_2$

$$|F_{6k}^{s}(G_{2})| = 3k^{2}, \qquad |F_{6k+1}^{s}(G_{2})| = 3k^{2} + k$$

$$|F_{6k+2}^{s}(G_{2})| = 3k^{2} + 2k, \qquad |F_{6k+3}^{s}(G_{2})| = 3k^{2} + 3k + 1$$

$$|F_{6k+4}^{s}(G_{2})| = 3k^{2} + 4k + 1, \qquad |F_{6k+5}^{s}(G_{2})| = 3k^{2} + 5k + 2.$$

$$|F_M^l(G_2)| = |F_M^s(G_2)|$$

$$\begin{split} |F_{12k}^{s}(F_{4})| &= 18k^{4} - k^{2}, & |F_{12k+1}^{s}(F_{4})| = 18k^{4} + 6k^{3} - \frac{5}{2}k^{2} - \frac{1}{2}k \\ |F_{12k+2}^{s}(F_{4})| &= 18k^{4} + 12k^{3} + 2k^{2}, & |F_{12k+3}^{s}(F_{4})| = 18k^{4} + 18k^{3} + \frac{7}{2}k^{2} - \frac{1}{2}k \\ |F_{12k+4}^{s}(F_{4})| &= 18k^{4} + 24k^{3} + 11k^{2} + 2k, & |F_{12k+5}^{s}(F_{4})| = 18k^{4} + 30k^{3} + \frac{31}{2}k^{2} + \frac{5}{2}k \\ |F_{12k+6}^{s}(F_{4})| &= 18k^{4} + 36k^{3} + 26k^{2} + 8k + 1, & |F_{12k+7}^{s}(F_{4})| = 18k^{4} + 42k^{3} + \frac{67}{2}k^{2} + \frac{21}{2}k + 1 \\ |F_{12k+8}^{s}(F_{4})| &= 18k^{4} + 48k^{3} + 47k^{2} + 20k + 3, & |F_{12k+9}^{s}(F_{4})| = 18k^{4} + 54k^{3} + \frac{115}{2}k^{2} + \frac{51}{2}k + 4 \\ |F_{12k+10}^{s}(F_{4})| &= 18k^{4} + 60k^{3} + 74k^{2} + 40k + 8, & |F_{12k+11}^{s}(F_{4})| = 18k^{4} + 66k^{3} + \frac{175}{2}k^{2} + \frac{99}{2}k + 10 \\ |F_{M}^{l}(F_{4})| &= |F_{M}^{s}(F_{4})|. \end{split}$$

Proof. For the case C_n , we have that $|F_{2k}(C_n)| = \binom{n+k}{n} + \binom{n+k-1}{n}$ from [3] and $m^s = 2n-2$ from Table 1. It can be verified directly that the formula

$$|F_{2k}^s(C_n)| = \binom{k+1}{n} + \binom{k}{n}$$

satisfies (64) for all values of $k \in \mathbb{N}$. Analogously, we obtain formulas for the remaining cases.

Example 5.1. For the Lie algebra C_2 , we have Coxeter number m = 4 and c = 2. For M = 4, the order of the group $\frac{1}{4}P^{\vee}/Q^{\vee}$ is equal to 32, and according to Theorem 5.2 we calculate

$$|F_4^s(C_2)| = |F_4^l(C_2)| = {3 \choose 2} + {2 \choose 2} = 4.$$

The coset representatives of $\frac{1}{4}P^{\vee}/Q^{\vee}$ and the fundamental domains F^s and F^l are depicted in Figure 1.

5.2. Number of elements of Λ^s_M and Λ^l_M .

In this section, we relate the numbers of elements of F_M^s , F_M^l to the numbers of elements Λ_M^s , Λ_M^l , defined by (50), (58). Firstly, we describe explicitly the sets Λ_M^s and Λ_M^l . Similarly to (40), we introduce the symbols t_i^s , $t_i^l \in \mathbb{R}$, $i = 0, \ldots, n$:

$$t_i^s \in \mathbb{N}, \qquad t_i^l \in \mathbb{Z}^{\ge 0}, \quad r_i \in \mathbb{R}^{s \vee}$$

$$t_i^s \in \mathbb{Z}^{\ge 0}, \quad t_i^l \in \mathbb{N}, \qquad r_i \in \mathbb{R}^{l \vee}.$$
(65)

The explicit form of Λ_M^s and Λ_M^l then follows from the explicit form of $F^{s\vee}$ and $F^{l\vee}$ in (41):

$$\Lambda_{M}^{s} = \left\{ t_{1}^{s}\omega_{1} + \dots + t_{n}^{s}\omega_{n} \mid t_{0}^{s} + t_{1}^{s}m_{1}^{\vee} + \dots + t_{n}^{s}m_{n}^{\vee} = M \right\}$$

$$\Lambda_{M}^{l} = \left\{ t_{1}^{l}\omega_{1} + \dots + t_{n}^{l}\omega_{n} \mid t_{0}^{l} + t_{1}^{l}m_{1}^{\vee} + \dots + t_{n}^{l}m_{n}^{\vee} = M \right\}.$$
(66)

Similarly to Proposition 5.1, we obtain the following one.

Proposition 5.3. Let $m^{s\vee}$ and $m^{l\vee}$ be the short and the long dual Coxeter numbers, respectively. Then

$$|\Lambda_{M}^{s}| = \begin{cases} 0 & M < m^{s\vee} \\ 1 & M = m^{s\vee} \\ |F_{M-m^{s\vee}}| & M > m^{s\vee} \end{cases} \quad |\Lambda_{M}^{l}| = \begin{cases} 0 & M < m^{l\vee} \\ 1 & M = m^{l\vee} \\ |F_{M-m^{l\vee}}| & M > m^{l\vee} \end{cases}$$
(67)

Combining Propositions 5.1, 5.3 and 2.1 and taking into account that $|F_M| = |\Lambda_M|$ we conclude with the following crucial result.



FIGURE 1. The fundamental domains F^s and F^l of C_2 . The fundamental domain F is depicted as the grey triangle containing borders H^s and H^l , depicted as the thick dashed line and dot-and-dashed lines, respectively. The coset representatives of $\frac{1}{4}P^{\vee}/Q^{\vee}$ are shown as 32 black dots. The four representatives belonging to F_4^s and F_4^l are crossed with '+' and '×', respectively. The dashed lines represent 'mirrors' r_0, r_1 and r_2 . Circles are elements of the root lattice Q; together with the squares they are elements of the weight lattice P.

Corollary 5.4. For the numbers of elements of the sets Λ_M^s and Λ_M^l it holds that

$$\begin{aligned} |\Lambda_M^s| &= |F_M^s|,\\ |\Lambda_M^l| &= |F_M^l|. \end{aligned}$$

$$\tag{68}$$

Example 5.2. For the Lie algebra C_2 we have |P/4Q| = 32 and according to Theorem 5.2 and Corollary 5.4 we have

$$|\Lambda_4^s(C_2)| = |\Lambda_4^l(C_2)| = 4.$$

The cosets representants of P/4Q together with the grids of weights $\Lambda_4^s(C_2)$ and $\Lambda_4^l(C_2)$ are depicted in Figure 2.

6. Discrete orthogonality and transforms of S^s – and S^l – functions

6.1. Discrete orthogonality of S^s - and S^l -functions.

To describe the discrete orthogonality of the S^{l} – and S^{s} –functions, we use the ideas discussed in [13] and reformulated in [3]. Recall that basic orthogonality relations from [3, 13] are, for any $\lambda, \lambda' \in P/MQ$, of the form:

$$\sum_{y \in \frac{1}{M} P^{\vee}/Q^{\vee}} e^{2\pi i \langle \lambda - \lambda', y \rangle} = c M^n \delta_{\lambda, \lambda'}.$$
(69)



FIGURE 2. The grids of weights $\Lambda_4^s(C_2)$ and $\Lambda_4^l(C_2)$ of C_2 . The darker grey triangle is the fundamental domain F^{\vee} and the lighter grey triangle is the domain $4F^{\vee}$. The borders $4H^{s\vee}$ and $4H^{l\vee}$ are depicted as the thick dashed lines and dot-and-dashed lines, respectively. The cosets representants of P/4Q of C_2 are shown as 32 black dots. The four representants belonging to $\Lambda_4^s(C_2)$ and $\Lambda_4^l(C_2)$ are crossed with '+' and '×', respectively. The dashed lines represent dual 'mirrors' r_0^{\vee}, r_1, r_2 and the affine mirror $r_{0,4}^{\vee}$ is defined by $r_{0,4}^{\vee}\lambda = 4r_0^{\vee}(\lambda/4)$. The circles and squares coincide with those in Figure 1.

We define the scalar product of two functions $f,g:F^s_M\to \mathbb{C}$ or $f,g:F^l_M\to \mathbb{C}$ by

$$\langle f, g \rangle_{F_M^s} = \sum_{x \in F_M^s} \varepsilon(x) f(x) \overline{g(x)}, \quad \langle f, g \rangle_{F_M^l} = \sum_{x \in F_M^l} \varepsilon(x) f(x) \overline{g(x)}, \tag{70}$$

where the numbers $\varepsilon(x)$ are determined by (13). We show that Λ_M^s and Λ_M^l are the lowest maximal sets of pairwise orthogonal S^s - and S^l -functions.

Theorem 6.1. For $\lambda, \lambda' \in \Lambda_M^s$ it holds that

$$\langle \varphi_{\lambda}^{s}, \varphi_{\lambda'}^{s} \rangle_{F_{M}^{s}} = c \left| W \right| M^{n} h_{\lambda}^{\vee} \delta_{\lambda,\lambda'} \tag{71}$$

and for $\lambda, \lambda' \in \Lambda_M^l$ it holds that

$$\langle \varphi^l_{\lambda}, \varphi^l_{\lambda'} \rangle_{F^l_M} = c \left| W \right| M^n h^{\vee}_{\lambda} \delta_{\lambda,\lambda'}, \tag{72}$$

where c, h_{λ}^{\vee} were defined by (2), (21), respectively, |W| is the number of elements of the Weyl group W and n is the rank of G.

Proof. Since φ_{λ}^s vanishes on $F_M \setminus F_M^s$, we have

(

$$\varphi_{\lambda}^{s}, \varphi_{\lambda'}^{s}\rangle_{F_{M}^{s}} = \sum_{x \in F_{M}^{s}} \varepsilon(x)\varphi_{\lambda}^{s}(x)\overline{\varphi_{\lambda'}^{s}(x)} = \sum_{x \in F_{M}} \varepsilon(x)\varphi_{\lambda}^{s}(x)\overline{\varphi_{\lambda'}^{s}(x)}.$$

The equality

$$\sum_{x\in F_M}\varepsilon(x)\varphi_{\lambda}^s(x)\overline{\varphi_{\lambda'}^s(x)}=\sum_{y\in \frac{1}{M}P^\vee/Q^\vee}\varphi_{\lambda}^s(y)\overline{\varphi_{\lambda'}^s(y)}$$

follows from (15) and (59) and the *W*-invariance of the expression $\varphi_{\lambda}^{s}(x)\overline{\varphi_{\lambda'}^{s}(x)}$. Then, using the *W*-invariance of $\frac{1}{M}P^{\vee}/Q^{\vee}$ and (69), we have

$$\begin{split} \langle \varphi_{\lambda}^{s}, \, \varphi_{\lambda'}^{s} \rangle_{F_{M}^{s}} &= \sum_{w' \in W} \sum_{w \in W} \sum_{y \in \frac{1}{M} P^{\vee}/Q^{\vee}} \sigma^{s}(ww') e^{2\pi \mathrm{i}\langle w\lambda - w'\lambda', \, y \rangle} \\ &= |W| \sum_{w' \in W} \sum_{y \in \frac{1}{M} P^{\vee}/Q^{\vee}} \sigma^{s}(w') e^{2\pi \mathrm{i}\langle \lambda - w'\lambda', \, y \rangle} \\ &= c \, |W| \, M^{n} \sum_{w' \in W} \sigma^{s}(w') \delta_{w'\lambda',\lambda}. \end{split}$$

If $\lambda = w'\lambda'$, then we have from (23) that $w'\lambda = \lambda = \lambda'$, i.e. $w' \in \operatorname{Stab}^{\vee}(\lambda)$. Any $\lambda \in \Lambda_M^s$ is of the form $\lambda = b + MQ$ with $b \in MF^{s\vee}$. Then, considering (24) and (38), we have

$$\sigma^{s}(\mathrm{Stab}^{\vee}(\lambda)) = \sigma^{s} \circ \widehat{\psi}\left(\mathrm{Stab}_{\widehat{W}^{\mathrm{aff}}}(b/M)\right) = \{1\}$$

i.e. we obtain $\sigma^s(w') = 1$ for any $w' \in \operatorname{Stab}^{\vee}(\lambda)$, and consequently

$$\sum_{w' \in W} \sigma^s(w') \delta_{w'\lambda',\lambda} = \sum_{w' \in W} \delta_{w'\lambda',\lambda} = h_{\lambda}^{\vee} \delta_{\lambda',\lambda}.$$

The case of S^l -functions is similar.

Example 6.1. The highest root ξ and the highest dual root η of C_2 are given by the formulas

$$\xi = 2\alpha_1 + \alpha_2, \quad \eta = \alpha_1^{\vee} + 2\alpha_2^{\vee},$$

The Weyl group of C_2 has eight elements, |W| = 8, and we calculate the determinant of the Cartan matrix c = 2. For a parameter with coordinates in ω -basis (a, b) and for a point with coordinates in α^{\vee} -basis (x, y), we have the following explicit form of S^s - and S^l -functions of C_2 :

$$\begin{split} \varphi^s_{(a,b)}(x,y) =& 2\{\cos(2\pi((a+2b)x-by)) + \cos(2\pi(ax+by)) \\ &- \cos(2\pi((a+2b)x-(a+b)y)) - \cos(2\pi(ax-(a+b)y))\} \\ \varphi^l_{(a,b)}(x,y) =& 2\{-\cos(2\pi((a+2b)x-by)) + \cos(2\pi(ax+by)) \\ &- \cos(2\pi((a+2b)x-(a+b)y)) + \cos(2\pi(ax-(a+b)y))\}. \end{split}$$

The grids F_M^s and F_M^l are given by

$$\begin{split} F_M^s(C_2) &= \left\{ \frac{u_1^s}{M} \omega_1^{\vee} + \frac{u_2^s}{M} \omega_2^{\vee} \mid u_0^s, \, u_2^s \in \mathbb{Z}^{\ge 0}, \, u_1^s \in \mathbb{N}, \, u_0^s + 2u_1^s + u_2^s = M \right\} \\ F_M^l(C_2) &= \left\{ \frac{u_1^l}{M} \omega_1^{\vee} + \frac{u_2^l}{M} \omega_2^{\vee} \mid u_0^l, \, u_2^l \in \mathbb{N}, \, u_1^l \in \mathbb{Z}^{\ge 0}, \, u_0^l + 2u_1^l + u_2^l = M \right\} \end{split}$$

and the grids of weights Λ^s_M and Λ^l_M are determined by

$$\Lambda_M^s(C_2) = \left\{ t_1^s \omega_1 + t_2^s \omega_2 \mid t_0^s, t_1^s \in \mathbb{N}, t_2^s \in \mathbb{Z}^{\ge 0}, t_0^s + t_1^s + 2t_2^s = M \right\}$$
$$\Lambda_M^l(C_2) = \left\{ t_1^l \omega_1 + t_2^l \omega_2 \mid t_0^l, t_1^l \in \mathbb{Z}^{\ge 0}, t_2^l \in \mathbb{N}, t_0^l + t_1^l + 2t_2^l = M \right\}.$$

The discrete orthogonality relations of S^s – and S^l – functions of C_2 , which hold for any two functions φ_{λ}^s , $\varphi_{\lambda'}^s$ labeled by $\lambda, \lambda' \in \Lambda_M^s(C_2)$, and φ_{λ}^l , $\varphi_{\lambda'}^l$ labeled by $\lambda, \lambda' \in \Lambda_M^l(C_2)$, are of the form (71) and (72), respectively. The calculation procedure of the coefficients $\varepsilon(x)$, h_{λ}^{\vee} , which appear in (70),(71) and (72), is detailed in §3.7 in [3]. The values of the coefficients $\varepsilon(x)$, h_{λ}^{\vee} for $x \in F_M^s(C_2)$, $\lambda \in \Lambda_M^s(C_2)$ and for $x \in F_M^l(C_2)$, $\lambda \in \Lambda_M^l(C_2)$ are listed in Table 2. We represent each point $x \in F_M^s(C_2)$ and each weight $\lambda^s \in \Lambda_M(C_2)$ by the coordinates $[u_0^s, u_1^s, u_2^s]$ and $[t_0^s, t_1^s, t_2^s]$. Similarly, we represent each point $x \in F_M^l(C_2)$ and each weight $\lambda^l \in \Lambda_M(C_2)$ by the coordinates $[u_0^l, u_1^l, u_2^l]$ and $[t_0^l, t_1^l, t_2^l]$.

$x \in F^s_M(C_2)$	$\varepsilon(x)$	$\lambda \in \Lambda^s_M(C_2)$	h_{λ}^{\vee}
$[u_0^s, u_1^s, u_2^s]$	8	$[t_0^s, t_1^s, t_2^s]$	1
$[0, u_1^s, u_2^s]$	4	$[t_0^s, t_1^s, 0]$	2
$[u_0^s, u_1^s, 0]$	4		
$[0, u_1^s, 0]$	2		
	•		
$x \in F_M^l(C_2)$	$\varepsilon(x)$	$\lambda \in \Lambda^l_M(C_2)$	$ h_{\lambda}^{\vee}$
$\frac{x \in F_M^l(C_2)}{[u_0^l, u_1^l, u_2^l]}$	$\frac{\varepsilon(x)}{8}$	$\frac{\lambda \in \Lambda^l_M(C_2)}{[t^l_0, t^l_1, t^l_2]}$	$\frac{h_{\lambda}^{\vee}}{1}$
$ \begin{array}{c} x \in F_M^l(C_2) \\ \hline [u_0^l, u_1^l, u_2^l] \\ [u_0^l, 0, u_2^l] \end{array} $	$\begin{array}{c}\varepsilon(x)\\ 8\\ 4\end{array}$	$\frac{\lambda \in \Lambda_M^l(C_2)}{[t_0^l, t_1^l, t_2^l]} \\ [0, t_1^l, t_2^l]$	$\frac{h_{\lambda}^{\vee}}{1}$
$\begin{array}{c} x \in F_M^l(C_2) \\ \hline [u_0^l, u_1^l, u_2^l] \\ [u_0^l, 0, u_2^l] \end{array}$	$\frac{\varepsilon(x)}{8}$	$\frac{\lambda \in \Lambda^l_M(C_2)}{ \begin{bmatrix} t^l_0, t^l_1, t^l_2 \end{bmatrix} } \\ \begin{bmatrix} 0, t^l_1, t^l_2 \end{bmatrix} \\ \begin{bmatrix} 0, t^l_1, t^l_2 \end{bmatrix} \\ \begin{bmatrix} t^l_0, 0, t^l_2 \end{bmatrix}$	$ \begin{array}{c c} h_{\lambda}^{\vee} \\ \hline 1 \\ 2 \\ 2 \end{array} $

TABLE 2. The coefficients $\varepsilon(x)$ and h_{λ}^{\vee} of C_2 . All variables $u_0^s, u_1^s, u_2^s, t_0^s, t_1^s, t_2^s$ and $u_0^l, u_1^l, u_2^l, t_0^l, t_1^l, t_2^l$ are assumed to be natural numbers.

6.2. Discrete S^s - and S^l -transforms.

Analogously to ordinary Fourier analysis, we define interpolating functions I_M^s and I_M^l

$$I_M^s(x) := \sum_{\lambda \in \Lambda_M^s} c_\lambda^s \varphi_\lambda^s(x), \qquad I_M^l(x) := \sum_{\lambda \in \Lambda_M^l} c_\lambda^l \varphi_\lambda^l(x), \quad x \in \mathbb{R}^n$$
(73)

which are given in terms of expansion functions φ_{λ}^{s} and φ_{λ}^{l} and expansion coefficients c_{λ}^{s} , c_{λ}^{l} , whose values need to be determined. These interpolating functions can also be understood as finite cut-offs of infinite expansions. Suppose we have some function f sampled on the grid F_{M}^{s} or F_{M}^{l} . The interpolation of f consists in finding the coefficients c_{λ}^{s} or c_{λ}^{l} in the interpolating functions (73) such that

$$I_M^s(x) = f(x), \quad x \in F_M^s$$

$$I_M^l(x) = f(x), \quad x \in F_M^l$$
(74)

Relations (68) and (71), (72) allow us to view the values $\varphi_{\lambda}^{s}(x)$ with $x \in F_{M}^{s}$, $\lambda \in \Lambda_{M}^{s}$ and the values $\varphi_{\lambda}^{l}(x)$ with $x \in F_{M}^{l}$, $\lambda \in \Lambda_{M}^{l}$ as elements of non-singular square matrices. These invertible matrices coincide with the matrices of the linear systems (74). Thus, the coefficients c_{λ}^{s} and c_{λ}^{l} can be uniquely determined. The formulas for calculation of c_{λ}^{s} and c_{λ}^{l} , which we call discrete S^{s} and S^{l} -transforms, can obtained by means of calculation of standard Fourier coefficients

$$c_{\lambda}^{s} = \frac{\langle f, \varphi_{\lambda}^{s} \rangle_{F_{M}^{s}}}{\langle \varphi_{\lambda}^{s}, \varphi_{\lambda}^{s} \rangle_{F_{M}^{s}}} = (c |W| M^{n} h_{\lambda}^{\vee})^{-1} \sum_{x \in F_{M}^{s}} \varepsilon(x) f(x) \overline{\varphi_{\lambda}^{s}(x)}$$

$$c_{\lambda}^{l} = \frac{\langle f, \varphi_{\lambda}^{l} \rangle_{F_{M}^{l}}}{\langle \varphi_{\lambda}^{l}, \varphi_{\lambda}^{l} \rangle_{F_{M}^{l}}} = (c |W| M^{n} h_{\lambda}^{\vee})^{-1} \sum_{x \in F_{M}^{l}} \varepsilon(x) f(x) \overline{\varphi_{\lambda}^{l}(x)}$$
(75)

and the corresponding Plancherel formulas also hold

$$\sum_{x \in F_M^s} \varepsilon(x) |f(x)|^2 = c |W| M^n \sum_{\lambda \in \Lambda_M^s} h_{\lambda}^{\vee} |c_{\lambda}^s|^2$$
$$\sum_{x \in F_M^l} \varepsilon(x) |f(x)|^2 = c |W| M^n \sum_{\lambda \in \Lambda_M^l} h_{\lambda}^{\vee} |c_{\lambda}^l|^2.$$

7. Concluding Remarks

• In view of the ever-increasing amount of digital data, practically the most valuable property of the orbit functions of C-, S-, S^l- and S^s- families is their discrete orthogonality. The four families are distinguished most notably by their behavior at the boundary of F. The functions of S^l- and S^s- families do not have an analog in one variable, i.e. rank 1 simple Lie group.

• The product of two S^s -functions or two S^l -functions with the same underlying Lie group and the same arguments $x \in \mathbb{R}^n$ but different dominant weights, say λ and λ' , decomposes into the sum of C-functions:

$$\varphi_{\lambda}^{s}(x) \cdot \varphi_{\lambda'}^{s}(x) = \sum_{w \in W} \sigma^{s}(w) \Phi_{\lambda + w\lambda'}(x), \quad \varphi_{\lambda}^{l}(x) \cdot \varphi_{\lambda'}^{l}(x) = \sum_{w \in W} \sigma^{l}(w) \Phi_{\lambda + w\lambda'}(x).$$

where Φ_{λ} denotes the (normalized) *C*-function $\Phi_{\lambda} = \varphi_{\lambda}^{1}$.

- The present work raises the question under which conditions converge the functional series $\{I_M^s\}_{M=1}^{\infty}, \{I_M^l\}_{M=1}^{\infty}$ assigned to a function $f: F \to \mathbb{C}$ by the relations (73), (75).
- In addition to the C- and S-functions, which are multidimensional generalizations of common cosine and sine functions, the E-functions generalizing the exponential functions [9] is also defined [15]. The E-functions also admit discrete orthogonality [4, 13]. For these 'standard' E-functions, the kernel of the homomorphism σ^e , given by (30), is crucial. It turns out that there are altogether six types of E-functions once the kernels of the sign homomorphisms σ^s and σ^l are included in the definition. So far, these six types have been studied in full detail only for rank 2 Lie groups [2].
- A general one-to-one link between the orbit functions and orthogonal polynomials in *n* variables was pointed out in [14]. Extensive literature exists about orthogonal polynomials, although most of it pertains to 2-variable polynomials. It cannot be assumed that our Lie group defined polynomials of rank two were not taken into account. For a greater number of variables, not all of the polynomials defined from the simple Lie groups have been noticed. Discrete orthogonality of the polynomials in more than one variable is outside the scope of traditional approaches.

Acknowledgments

We gratefully acknowledge the support of this work by the Natural Sciences and Engineering Research Council of Canada and by the Doppler Institute of the Czech Technical University in Prague. JH is grateful for the hospitality extended to him at the Centre de recherches mathématiques, Université de Montréal. JH gratefully acknowledges support by the Ministry of Education of Czech Republic (project MSM6840770039). JP expresses his gratitude for the hospitality of the Doppler Institute.

References

- A. Bjorner, F. Brenti, Combinatorics of Coxeter groups, Graduate Texts in Mathematics, 231 (2005) Springer, New York.
- [2] L. Háková, J. Hrivnák, J. Patera, Six types of E-functions of the Lie groups O(5) and G(2), J. Phys. A: Math. Theor. 45 (2012), 125201 arXiv:1202.5031.
- [3] J. Hrivnák, J. Patera, On discretization of tori of compact simple Lie groups, J. Phys. A: Math. Theor. 42 (2009) 385208, arXiv:0905.2395.
- [4] J. Hrivnák, J. Patera, On E-discretization of tori of compact simple Lie groups, J. Phys. A: Math. Theor. 43 (2010) 165206, arXiv:0912.4194.
- [5] J. E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge Studies in Advanced Mathematics, 29 (1990) Cambridge University Press, Cambridge.
- [6] V. Kac, Automorphisms of finite order of semi-simple Lie algebras, Funct. Anal. and Appl., 3 (1969) 252–254.
- [7] A. U. Klimyk, J. Patera, Orbit functions, SIGMA (Symmetry, Integrability and Geometry: Methods and Applications) 2 (2006), 006, 60. pages, math-ph/0601037.
- [8] A. U. Klimyk, J. Patera, Antisymmetric orbit functions, SIGMA (Symmetry, Integrability and Geometry: Methods and Applications) 3 (2007), paper 023, 83 pages; math-ph/0702040.
- [9] A. U. Klimyk, J. Patera, *E-orbit functions*, SIGMA (Symmetry, Integrability and Geometry: Methods and Applications) 4 (2008), 002, 57 pages; arXiv:0801.0822.
- [10] R. V. Moody, L. Motlochová, and J. Patera, New families of Weyl group orbit functions,, arXiv:1202.4415.
- [11] R. V. Moody, J. Patera, Characters of elements of finite order in simple Lie groups, SIAM J. on Algebraic and Discrete Methods 5 (1984) 359-383.
- [12] R. V. Moody, J. Patera, Computation of character decompositions of class functions on compact semisimple Lie groups, Mathematics of Computation 48 (1987) 799–827.
- [13] R. V. Moody and J. Patera, Orthogonality within the families of C-, S-, and E-functions of any compact semisimple Lie group, SIGMA (Symmetry, Integrability and Geometry: Methods and Applications) 2 (2006) 076, 14 pages, mathph/0611020.
- [14] M. Nesterenko, J. Patera, A. Tereszkiewicz Orthogonal polynomials of compact simple Lie groups, International Journal of Mathematics and Mathematical Sciences (2011) 969424.

- [15] J. Patera, Compact simple Lie groups and theirs C-, S-, and E-transforms, SIGMA (Symmetry, Integrability and Geometry: Methods and Applications) 1 (2005) 025, 6 pages, math-ph/0512029.
- [16] M. Szajewska, Four types of special functions of G_2 and their discretization, Integral Transform. Spec. Funct. **23** (6) (2012) 445-472, arXiv:1101.2502.
- [17] E. B. Vinberg, A. L. Onishchik, Lie groups and Lie algebras, Springer, New York, 1994.