

SYMPLECTIC 4-MANIFOLDS WITH FIXED POINT FREE CIRCLE ACTIONS

JONATHAN BOWDEN

ABSTRACT. We show that recent results of Friedl-Vidussi and Chen imply that a symplectic manifold admits a fixed point free circle action if and only if it admits a symplectic circle action and we give a complete description of the symplectic cone in this case. This then completes the characterisation of symplectic 4-manifolds that admit non-trivial circle actions.

1. INTRODUCTION

Recently Friedl and Vidussi, [6] solved the long standing Taubes Conjecture, which classifies which 4-manifolds of the form $M \times S^1$ admit symplectic forms. Moreover, they determined exactly which cohomology classes can be represented by symplectic forms. Using recent results of D. Wise, [12] they have extended their results to the case of non-trivial S^1 -bundles in [8]. In this note we observe that their results as well of those of Chen, who obtained partial results in the fixed point free case in [3], imply the analogue of ([8], Theorem 1.3) for all fixed point free circle actions.

Before stating our main result we fix some notation and terminology. Let $X \xrightarrow{p} M$ be an orientable 4-manifold with a fixed point free circle action and quotient space $M = X/S^1$. The quotient space is an orbifold whose underlying topological space $|M|$ is a manifold since all the stabilisers of the S^1 -action are necessarily cyclic and the singular locus consists of a collection of branching circles.

We let M_{reg} denote the complement of a regular neighbourhood of the singular locus of M and $X_{reg} = p^{-1}(M_{reg})$, which is an honest S^1 -bundle so that the pushforward map p_* is well-defined for cohomology classes in $H^*(X_{reg}, \mathbb{R})$. The manifold M_{reg} has toroidal boundary and thus one may define the Thurston norm on $H^1(M_{reg}, \mathbb{R})$ in the usual fashion. Finally for $\psi \in H^2(X_{reg}, \mathbb{R})$ we let ψ_{reg} denote the restriction of ψ to X_{reg} .

Theorem 1. *Let $X \xrightarrow{p} M$ be an oriented manifold admitting a fixed point free S^1 -action with quotient space M and let $\psi \in H^2(X, \mathbb{R})$. Then the following are equivalent:*

- (1) ψ can be represented by a symplectic form,
- (2) ψ can be represented by an S^1 -invariant symplectic form,
- (3) $\psi^2 > 0$ and $p_*\psi_{reg} \in H^1(M_{reg}, \mathbb{R})$ lies in the open cone over a fibered face of the Thurston norm ball and is the restriction of a class in $H^1(|M|, \mathbb{R})$.

Note that if a class ϕ is integral, then ϕ can be represented by a fibration over S^1 that is transverse to the singular locus of $|M|$ if and only if its restriction of ϕ to M_{reg} , which we denote by ϕ_{reg} , is fibered. Recall that a fibration of a manifold with boundary is required

Date: June 5, 2012.

2010 Mathematics Subject Classification. Primary 57R17; Secondary 57N10, 57N13.

to be transverse to the boundary. Furthermore, since ϕ_{reg} is the restriction of a class in $H^1(|M|, \mathbb{R})$, it automatically vanishes on the meridian classes in ∂M_{reg} so that if ϕ_{reg} is fibered, then the induced fibration on the boundary is necessarily meridional. Thus the fibration dual to ϕ_{reg} extends to $|M|$ in the desired way by filling in discs near the singular locus. In particular, part (3) of Theorem 1 implies that the underlying manifold $|M|$ is fibered and we obtain a positive answer to the following conjecture, which implies ([3], Conjecture 1.7) as a special case.

Conjecture 1 (Generalised Taubes conjecture). *Let X be a symplectic 4-manifold that admits a non-trivial fixed point free circle action with quotient orbifold M . Then the (possibly empty) singular locus L of M is a meridionally fibered link.*

Furthermore, as noted in ([3], p. 6), Theorem 1 completes the characterisation of which symplectic manifolds admit non-trivial S^1 -actions. For Baldridge, [1] showed that if a non-trivial S^1 -action on a symplectic 4-manifold has fixed points then X is rational or ruled and thus admits an S^1 -invariant symplectic form for *some* non-trivial S^1 -action. In view of this we obtain the following corollary.

Corollary 1. *Let X be a symplectic 4-manifold that admits a non-trivial S^1 -action. Then either the action is fixed point free and the quotient space fibers over S^1 or X is rational or ruled. In either case, X admits a non-trivial symplectic S^1 -action.*

2. PROOF OF THEOREM 1

The proof of Theorem 1 is based on the following lemma, which provides a generalisation of ([4], Theorem 5.2) to include irrational classes. For the proof we assume a certain familiarity with the basic properties of the Thurston norm (cf. [11]).

Lemma 1. *Let \overline{M} be a 3-manifold with an orientation preserving smooth action of a finite group G and quotient orbifold $M = \overline{M}/G$. An element $\overline{\phi}$ in the invariant subspace $H^1(\overline{M}, \mathbb{R})^G$ admits a non-degenerate de Rham representative if and only if it admits a non-degenerate de Rham representative that is G -invariant.*

In particular, the restriction of the associated class $\phi \in H^1(|M|, \mathbb{R}) \cong H^1(\overline{M}, \mathbb{R})^G$ to M_{reg} lies in the open cone over a fibered face of the Thurston norm ball.

Proof. We first assume that $\overline{\phi}$ is rational. Since nothing changes after multiplying with positive constants, we may assume that $\overline{\phi}$ is in fact integral. In this case the first claim is just a restatement of ([4], Theorem 5.2), which can be applied in complete generality in view of ([10], Theorem 8.1). Note that the assumption $H^1(\overline{M}, \mathbb{Q})^G = \mathbb{Q}$ in ([4], Theorem 5.2) can be replaced by the fact that the fibration is given by a fibered class $\overline{\phi}$ that is G -invariant. Moreover, the proof in [4] actually gives a fibration that is transverse to the branching locus in \overline{M} . The quotient map π induces an isomorphism $H^1(\overline{M}, \mathbb{R})^G \cong H^1(|M|, \mathbb{R})$ so that there is a unique class ϕ with $\overline{\phi} = \pi^*\phi$ and the fibration dual to $\overline{\phi}$ descends to a fibration of $|M|$ dual to ϕ . Finally since the fibration is transverse to the singular locus it follows that the restriction of ϕ to M_{reg} is fibered.

We next assume that $\overline{\phi}$ is irrational and let ϕ be the unique class with $\overline{\phi} = \pi^*\phi$. We let ι_{reg} denote the natural inclusion $M_{reg} \hookrightarrow |M|$ and set $V = \text{Im}(\iota_{reg}^*)$. By the previous case all rational classes in V that are sufficiently close to $\phi_{reg} = \iota_{reg}^*\phi$ are fibered. If ϕ_{reg} itself did not lie in the open cone over a fibered face of the Thurston unit ball, then it must lie in the closed

cone over the boundary of a fibered face by the assumption that it can be approximated by fibered elements. Since the Thurston unit ball is rational, the intersection of the closed cone containing ϕ_{reg} with V must contain non-fibered rational points arbitrarily close to ϕ_{reg} , which gives a contradiction. Thus ϕ_{reg} admits a non-degenerate de Rham representative η_{reg} and $\pi^*\eta_{reg}$ extends to a G -equivariant representative of $\bar{\phi}$. \square

Proof of Theorem 1. The implication (2) \implies (1) is trivial.

(1) \implies (3): Let (X, ω) be a symplectic manifold with a fixed point free S^1 -action and quotient space M . By ([3], Proposition 1.8) there is a manifold \bar{M} and a smooth action by a finite group so that $M = \bar{M}/G$. Furthermore, we have the following commutative diagram:

$$\begin{array}{ccc} \pi^*X = \bar{X} & \xrightarrow{\bar{p}} & \bar{M} \\ \bar{\pi} \downarrow & & \downarrow \pi \\ X & \xrightarrow{p} & M, \end{array}$$

where π is the quotient map, $\bar{\pi}$ is an unramified covering and the induced S^1 -action on \bar{X} is free. Moreover, the group G acts naturally on \bar{X} as the group of deck transformations of $\bar{\pi}$.

Thus $\bar{\omega} = \pi^*\omega$ is a symplectic form and by ([8], Theorem 1.4) its image under the pushforward map $\bar{p}_*(\bar{\omega}) \in H^1(\bar{M}, \mathbb{R})$ lies in the open cone over a fibered face of the Thurston norm ball. Since $\bar{\omega}$ is G -invariant and the action on \bar{X} is fiber preserving, the class $\bar{\phi} = \bar{p}_*(\bar{\omega})$ is also G -invariant. We let ϕ be the unique class such that $\pi^*\phi = \bar{\phi}$. By Lemma 1 the restriction ϕ_{reg} to M_{reg} lies in the open cone over a fibered face. Finally the naturality of the transfer homomorphism implies that the restriction of ϕ_{reg} agrees with $p_*\omega_{reg}$.

(3) \implies (2): By assumption $\phi_{reg} = p_*\psi_{reg}$ lies in the open cone over a fibered face of the Thurston norm ball and ϕ_{reg} is the restriction of a class $\phi \in H^1(|M|, \mathbb{R})$. In particular, $|M|$ fibers over S^1 . We first note that M is a *very good* orbifold so that it is a quotient of a manifold \bar{M} by a smooth action of a finite group G . For this it suffices to rule out bad 2-suborbifolds by ([2], Corollary 3.28). However, a bad 2-suborbifold is topologically a sphere that is essential in $H_2(|M|, \mathbb{Z})$ and as in the proof of ([3], Lemma 2.3) this implies that $b_2^+(X) = b_2(|M|) - 1$. Thus $|M| = S^2 \times S^1$ and $b_2^+(X) = 0$, contradicting the assumption that $\psi^2 > 0$.

We are thus in the situation as above and we set $\bar{\phi} = \pi^*\phi$. Since a degree one cohomology class on \bar{M} is determined by its restriction to the complement of the branching locus, we deduce that $\bar{\phi}$ and $p_*(\pi^*\psi)$ agree as cohomology classes. We then note that the construction of S^1 -invariant forms in [5] and its extension to irrational classes ([7], Theorem 1.1) can be done G -equivariantly.

First choose a G -invariant representative γ of $e(\bar{X})$, which can be obtained as the curvature of a G -equivariant angular form. By ([7], Lemma 2.1) we may write $\gamma = \bar{\phi} \wedge \beta$. After averaging over G this equation still holds, so β can be assumed to be G -equivariant. Let η be a G -invariant angular 1-form so that $d\eta = \bar{p}^*\gamma$ and let $\Omega \in H^2(|M|, \mathbb{R})$ be such that the following holds in cohomology

$$\bar{\pi}^*\psi - \eta \wedge \bar{p}^*\bar{\phi} = \pi^*\Omega.$$

Such an Ω exists since the left hand side is G -equivariant and π^* is an isomorphism onto $H^1(\bar{M}, \mathbb{R})^G$. The fact that $\pi^*\psi^2 > 0$ implies that $\bar{p}^*\bar{\phi} \wedge \pi^*\Omega > 0$. Thus by ([7], Lemma 2.2) there is a non-vanishing 2-form $\bar{\Omega}$ representing the class $\pi^*\Omega$ so that $\bar{\phi} \wedge \bar{\Omega} > 0$, again after

averaging we may assume that $\bar{\Omega}$ is G -invariant. Thus the S^1 -invariant form

$$\bar{\omega}_{inv} = \eta \wedge \bar{p}^* \bar{\phi} + \bar{p}^* \bar{\Omega}$$

represents $\pi^* \psi$ and descends to an S^1 -invariant form ω_{inv} on X which is cohomologous to ψ . \square

Remark 1. A vital step in the proof of Theorem 1 was Chen's observation that the base orbifold of a symplectic manifold X admitting a fixed point free S^1 -action is a quotient of a manifold by a finite group action. The main technical point in the proof of ([3], Proposition 1.8) is to rule out bad 2-orbifolds in the base. This is achieved by results relating the Seiberg-Witten invariants of the base orbifold to those of the underlying manifold.

We sketch a different proof which uses more standard Seiberg-Witten vanishing results. For background on the Seiberg-Witten invariants we refer to [9] and the references therein. First observe that a bad 2-suborbifold Σ in the quotient orbifold $M = X/S^1$ can intersect at most 2 singular curves L_1, L_2 each in at most one point. Taking a neighbourhood N of $|\Sigma| \cup L_1$ gives a topological splitting of the base $|M| = (S^2 \times S^1) \# M'$ so that preimage of the splitting sphere in $|M|$ induces a splitting $X = X_1 \cup_S X_2$, where S is either $S^2 \times S^1$ or S^3 depending on whether L_2 is empty or not. Moreover, as in the proof of ([3], Lemma 2.3) we must have $b_1(M') > 0$ by the assumption that $b_2^+(X) > 0$. If S is a 3-sphere, then $b_2^+(X_2) > 0$ and by taking the covering \bar{X} of X induced by the natural surjection

$$\pi_1(X) \rightarrow \pi_1(S^2 \times S^1) \rightarrow \mathbb{Z}_n$$

we obtain a splitting of $\bar{X} = \bar{X}_1 \cup_{S^3} \bar{X}_2$, where $b_2^+(X_1), b_2^+(X_2) \geq 1$. It follows that the Seiberg-Witten invariants of \bar{X} are trivial.

If $S = S^2 \times S^1$, then we take the covering \bar{X} of X induced by a surjection

$$\pi_1(X) \rightarrow \pi_1(S^2 \times S^1) * \pi_1(|M'|) \rightarrow \mathbb{Z}_n \times \mathbb{Z}_n.$$

The embedded 2-sphere in S then becomes essential in the covering and $b_1(\bar{X})$, and hence $b_2^+(\bar{X})$, may be assumed to be arbitrarily large. Again the Seiberg-Witten invariants are trivial and in both cases we obtain a contradiction to the non-vanishing results of Taubes for the Seiberg-Witten invariants of a symplectic 4-manifold.

ACKNOWLEDGMENTS:

We thank P. Suárez-Serrato for his stimulating questions and S. Friedl for helpful comments. The hospitality of the Max Planck Institute für Mathematik in Bonn, where this research was carried out, is also gratefully acknowledged.

REFERENCES

- [1] S. Baldridge, *Seiberg-Witten vanishing theorem for S^1 -manifolds with fixed points*, Pacific J. Math. **217** (2004), no. 1, 1–10.
- [2] M. Boileau, S. Maillot and J. Porti, *Three-Dimensional Orbifolds and their Geometric Structures*, Panoramas et Synthèses, **15**, Société Mathématique de France, Paris, 2003.
- [3] W. Chen, *Seifert fibered four-manifolds with nonzero Seiberg-Witten invariant*, Preprint: arXiv:1103.5681v3, 2011.
- [4] A. Edmonds and C. Livingston, *Group actions on fibered three-manifolds*, Comment. Math. Helv. **58** (1983), no. 4, 529–542.
- [5] M. Fernández, A. Gray and J. Morgan, *Compact symplectic manifolds with free circle actions, and Massey products*, Michigan Math. J. **38** (1991), 271–283.

- [6] S. Friedl and S. Vidussi, *Twisted Alexander polynomials detect fibered 3-manifolds*, Ann. Math. **173** (2011), 1587–1643.
- [7] S. Friedl and S. Vidussi, *Construction of symplectic structures on 4-manifolds with a free circle action*, Preprint: arXiv:1102.0821v1, 2011 (to appear Proc. Royal Soc. Edinburgh).
- [8] S. Friedl and S. Vidussi, *A Vanishing Theorem for Twisted Alexander Polynomials with Applications to Symplectic 4-manifolds*, Preprint: arXiv:1205.2434v1, 2012.
- [9] R. Gompf and A. Stipsicz, *Kirby calculus and the topology of 4-manifolds*, Graduate Studies in Mathematics, **20** AMS, Providence RI 1999.
- [10] W. Meeks III and P. Scott, *Finite group actions on 3-manifolds*, Invent. Math. **86** (1986), no. 2, 287–346.
- [11] W. Thurston, *A norm for the homology of 3-manifolds*, Mem. Amer. Math. Soc. **59** (1986), no. 339, i-vi and 99-130.
- [12] D. Wise, *The structure of groups with a quasiconvex hierarchy*, 187 pages, Preprint: available on the webpage for the NSF-CBMS conference.

MATHEMATISCHES INSTITUT, UNIVERSITÄT AUGSBURG, UNIVERSITÄTSSTR. 14, 86159 AUGSBURG, GERMANY

Current address: Max-Planck-Institut für Mathematik, Vivatsgasse 7, 53129 Bonn, Germany

E-mail address: jonathan.bowden@math.uni-augsburg.de