

DIFFERENTIAL HARNACK ESTIMATES FOR PARABOLIC EQUATIONS

XIAODONG CAO * AND ZHOU ZHANG †

ABSTRACT. Let $(M, g(t))$ be a solution to the Ricci flow on a closed Riemannian manifold. In this paper, we prove differential Harnack inequalities for positive solutions of nonlinear parabolic equations of the type

$$\frac{\partial}{\partial t} f = \Delta f - f \ln f + Rf.$$

We also comment on an earlier result of the first author on positive solutions of the conjugate heat equation under the Ricci flow.

1. Introduction

Let $(M, g(t))$, $t \in [0, T)$, be a solution to the Ricci flow on a closed manifold M . In the first part of this paper, we deal with positive solutions of nonlinear parabolic equations on M . We establish Li-Yau type differential Harnack inequalities for such positive solutions. To be more precisely, $g(t)$ evolves under the Ricci flow

$$(1.1) \quad \frac{\partial g(t)}{\partial t} = -2Rc,$$

where Rc denote the Ricci curvature of $g(t)$. We first assume that the initial metric $g(0)$ has nonnegative curvature operator, which implies that for all time $t \in [0, T)$, $g(t)$ has nonnegative curvature operator (c.f. [7]). Consider a positive function $f(x, t)$ defined on $M \times [0, T)$, which solves the following nonlinear parabolic equation,

$$(1.2) \quad \frac{\partial f}{\partial t} = \Delta f - f \ln f + Rf.$$

The symbol Δ here stands for the Laplacian of the evolving metric $g(t)$ and R is the scalar curvature of $g(t)$. For simplicity, we omit $g(t)$ in the above notations. All geometry operators are with respect to the evolving metric $g(t)$.

Differential Harnack inequalities were originated by P. Li and S.-T. Yau in [12] for positive solutions of the heat equation (therefore it is also known as Li-Yau type Harnack estimates). The technique was then brought into the study of geometric evolution equation by R. Hamilton (for example, see [8]) and ever since has been playing an important role in the study of geometric flows. Applications include estimates

Date: Jan. 8, 2010.

2000 Mathematics Subject Classification. Primary 53C44.

* Research partially supported by NSF grant DMS 0904432.

† Research partially supported by NSF grant DMS 0904760.

on the heat kernel; curvature growth control; understanding the ancient solutions for geometric flows; proving noncollapsing result in the Ricci flow ([17]); etc. See [16] for a recent survey on this subject by L. Ni.

Using maximum principle, one can see that the solution for (1.2) will remain positive along the flow. It exists as long as the solution for (1.1) exists. The study of the Ricci flow coupled with a heat-type (or backward heat-type) equation started from R. Hamilton's paper [9]. Recently, there has been some interesting study on this topic. In [17], G. Perelman proved a differential Harnack inequality for the fundamental solution of the conjugate heat equation under the Ricci flow. In [2], the first author proved a differential Harnack inequality for general positive solutions of the conjugate heat equation, which was also proved independently by S. Kuang and Q. S. Zhang in [11]. For the Ricci flow coupled with the heat equation, the study was pursued in [3, 6, 15, 20]. Various estimates are obtained recently by M. Bailesteanu, A. Pulemotov and the first author in [1], and by S. Liu in [13]. For nonlinear parabolic equations under the Ricci flow, local gradient estimates for positive solutions of equation

$$\frac{\partial}{\partial t} f = \Delta f + af \ln f + bf,$$

where a and b are constants, has been studied by Y. Yang in [19]. For general evolving metrics, similar estimate has been obtained by A. Chau, L.-F. Tam and C. Yu in [4], by S.-Y. Hsu in [10], and by J. Sun in [18]. In [14], L. Ma proved a gradient estimate for the elliptic equation

$$\Delta f + af \ln f + bf = 0.$$

In (1.2), if one defines

$$u(x, t) = -\ln f(x, t),$$

then the function $u = u(x, t)$ satisfies the evolution equation

$$(1.3) \quad \frac{\partial u}{\partial t} = \Delta u - |\nabla u|^2 - R - u.$$

The computation from (1.2) to (1.3) is standard, which also gives the explicit relation between these two equations.

Our motivation to study (1.2) under the Ricci flow comes from the geometric interpretation of (1.3), which arises from the study of expanding Ricci solitons. Recall that if we are on a gradient expanding Ricci soliton (M, g) satisfying

$$R_{ij} + \nabla_i \nabla_j w = -\frac{1}{4} g_{ij},$$

where w is called soliton potential function, then we have

$$R(g) + \Delta_g w = -\frac{n}{4}.$$

In sight of this, by taking covariant derivative for the soliton equation and applying the second Bianchi identity, one can see that

$$R(g) + |\nabla_g w|_g^2 + \frac{w}{2} = \text{constant}.$$

Also notice that the Ricci soliton potential function w can be differed by a constant in the above equations. So by choosing this constant carefully, we have

$$R(g) + |\nabla_g w|_g^2 = -\frac{w}{2} - \frac{n}{8}.$$

One consequence of the above identities is the following

$$(1.4) \quad |\nabla_g w|_g^2 = \Delta_g w - |\nabla_g w|_g^2 - R(g) - w.$$

Recall that the Ricci flow solution for an expanding soliton is $g(t) = c(t) \cdot \phi(t)^*g$ (c.f. [5]), where $c(t) = 1 + \frac{t}{2}$ and the family of diffeomorphism $\phi(t)$ satisfies, for any $x \in M$,

$$\frac{\partial}{\partial t}(\phi(t)(x)) = \frac{1}{c(t)} \cdot (\nabla_g w)(\phi(t)(x)).$$

Thus the corresponding Ricci soliton potential $\phi(t)^*w$ satisfies

$$\frac{\partial \phi(t)^*w}{\partial t}(x) = \frac{1}{c(t)}(\nabla_g w)(w)(\phi(t)(x)) = |\nabla \phi(t)^*w|^2(x).$$

Along the Ricci flow, (1.4) becomes

$$|\nabla \phi^*w|^2 = \Delta \phi^*w - |\nabla \phi^*w|^2 - R - \frac{\phi^*w}{c(t)}.$$

Hence the evolution equation for the Ricci soliton potential is

$$(1.5) \quad \frac{\partial \phi(t)^*w}{\partial t} = \Delta \phi^*w - |\nabla \phi^*w|^2 - R - \frac{\phi^*w}{c(t)}.$$

The second nonlinear parabolic equation that we investigate in this paper is

$$(1.6) \quad \frac{\partial u}{\partial t} = \Delta u - |\nabla u|^2 - R - \frac{u}{1 + \frac{t}{2}}.$$

Notice that (1.3) and (1.6) are closely related and only differ by their last terms.

Our first result deals with (1.2) and (1.3).

Theorem 1.1. *Let $(M, g(t))$, $t \in [0, T)$, be a solution to the Ricci flow on a closed manifold, and suppose that $g(0)$ (and so $g(t)$) has weakly positive curvature operator. Let f be a positive solution to the heat equation (1.2), $u = -\ln f$ and*

$$(1.7) \quad H = 2\Delta u - |\nabla u|^2 - 3R - \frac{2n}{t}.$$

Then for all time $t \in (0, T)$

$$H \leq \frac{n}{4}.$$

Remark 1.1. *The result can be generalized to the context of M being non-compact. In order for the same argument to work, we need to assume that the Ricci flow solution $g(t)$ is complete with the curvature and all the covariant derivatives being uniformly bounded (in the space direction).*

Our next result deals with (1.6), which is also a natural evolution equation to consider with, by previous discussion.

Theorem 1.2. *Let $(M, g(t))$, $t \in [0, T)$, be a solution to the Ricci flow on a closed manifold, and suppose that $g(0)$ (and so $g(t)$) has weakly positive curvature operator. Let u be a smooth solution to (1.6), define*

$$(1.8) \quad H = 2\Delta u - |\nabla u|^2 - 3R - \frac{2n}{t}.$$

Then for all time $t \in (0, T)$

$$H \leq 0.$$

Remark 1.2. *If f is a positive function such that $f = e^{-u}$, then f satisfies the following evolution equation*

$$\frac{\partial f}{\partial t} = \Delta f + Rf - \frac{f \ln f}{1 + \frac{t}{2}}.$$

In [2], the first author studied the conjugate heat equation under the Ricci flow. In particular, the following theorem was proved.

Theorem 1.3. [2, Theorem 3.6] *Let $(M, g(t))$, $t \in [0, T]$, be a solution to the Ricci flow, suppose that $g(t)$ has nonnegative scalar curvature. Let f be a positive solution of the conjugate heat equation*

$$\frac{\partial}{\partial t} f = -\Delta f + Rf,$$

let $v = -\ln f - \frac{n}{2} \ln(4\pi\tau)$, $\tau = T - t$ and

$$P = 2\Delta v - |\nabla v|^2 + R - \frac{2n}{\tau}.$$

Then we have

$$(1.9) \quad \frac{\partial}{\partial \tau} P = \Delta P - 2\nabla P \cdot \nabla v - 2|v_{ij} + R_{ij} - \frac{1}{\tau}g_{ij}|^2 - \frac{2}{\tau}P - 2\frac{|\nabla v|^2}{\tau} - 2\frac{R}{\tau}.$$

Moreover, for all time $t \in [0, T)$,

$$P \leq 0.$$

In the last section, we apply a similar trick as in the proof of Theorem 1.1 and obtain a slightly different result, where we no longer need to assume that $g(t)$ has nonnegative scalar curvature.

Acknowledgments Xiaodong Cao wants to thank the organizers of the conference ‘‘Complex and Differential Geometry’’ for their invitation and hospitality. Both

authors want to express their gratitude to East China Normal University, where they started this discussion.

2. Proof of Theorem 1.1 and Application

The evolution equations of u , is very similar to what is considered in [3]. So the computation for the very general setting there can be applied. The only difference is now we have more terms coming from time derivative $\frac{\partial}{\partial t}u$.

Proof of Theorem 1.1. Recall the definition of H from (1.8), comparing with [3, Corollary 2.2], we have

$$(2.1) \quad \begin{aligned} \frac{\partial}{\partial t}H = & \Delta H - 2\nabla H \cdot \nabla u - 2|u_{ij} - R_{ij} - \frac{1}{t}g_{ij}|^2 - \frac{2}{t}H - \frac{2}{t}|\nabla u|^2 \\ & - 2\left(\frac{\partial}{\partial t}R + \frac{R}{t} + 2\nabla R \cdot \nabla u + 2R_{ij}u_iu_j\right) - 2\Delta u + 2|\nabla u|^2, \end{aligned}$$

where the last two terms of the right hand side coming from the extra term $-u$ in (1.3). Plugging in $-2\Delta u + 2|\nabla u|^2 = -H + |\nabla u|^2 - 3R - \frac{2n}{t}$, one arrives at

$$(2.2) \quad \begin{aligned} \frac{\partial}{\partial t}H = & \Delta H - 2\nabla H \cdot \nabla u - 2|u_{ij} - R_{ij} - \frac{1}{t}g_{ij}|^2 - \left(\frac{2}{t} + 1\right)H \\ & + \left(1 - \frac{2}{t}\right)|\nabla u|^2 - 3R - \frac{2n}{t} - \left(\frac{\partial}{\partial t}R + \frac{R}{t} + 2\nabla R \cdot \nabla u + 2R_{ij}u_iu_j\right). \end{aligned}$$

In sight of the definition of H (1.8), for t small enough, we have $H < 0$. Since g_{ij} has weakly positive curvature operator, by the trace Harnack inequality for the Ricci flow proved by R. Hamilton in [8], we have

$$\frac{\partial}{\partial t}R + \frac{R}{t} + 2\nabla R \cdot \nabla u + 2R_{ij}u_iu_j \geq 0.$$

Also we have $R \geq 0$. Notice that the term $(1 - \frac{2}{t})|\nabla u|^2$ prevents us from obtaining an upper bound for H for $t > 2$.

We can deal with this by the following simple manipulation. To begin with, one observes that from the definition of H ,

$$|\nabla u|^2 = 2\left(\Delta u - R - \frac{n}{t}\right) - H - R.$$

We also have the following equality from definition,

$$\text{tr}\left(u_{ij} - R_{ij} - \frac{1}{t}g_{ij}\right) = \Delta u - R - \frac{n}{t}.$$

Now we can continue the computation for the evolution of H as follows,

$$\begin{aligned}
\frac{\partial}{\partial t} H &\leq \Delta H - 2\nabla H \cdot \nabla u - 2|u_{ij} - R_{ij} - \frac{1}{t}g_{ij}|^2 - \left(\frac{2}{t} + 1\right) H - \frac{2}{t}|\nabla u|^2 \\
&\quad - 4R + 2\left(\Delta u - R - \frac{n}{t}\right) - H - \frac{2n}{t} \\
&\leq \Delta H - 2\nabla H \cdot \nabla u - \frac{2}{n}\left(\Delta u - R - \frac{n}{t}\right)^2 - \left(\frac{2}{t} + 1\right) H - \frac{2}{t}|\nabla u|^2 \\
&\quad - 4R + 2\left(\Delta u - R - \frac{n}{t}\right) - H - \frac{2n}{t} \\
&= \Delta H - 2\nabla H \cdot \nabla u - \left(\frac{2}{t} + 2\right) H - \frac{2}{t}|\nabla u|^2 - 4R - \frac{2n}{t} \\
&\quad - \frac{2}{n}\left(\Delta u - R - \frac{n}{t} - \frac{n}{2}\right)^2 + \frac{n}{2} \\
&\leq \Delta H - 2\nabla H \cdot \nabla u - \left(\frac{2}{t} + 2\right) H - \frac{2}{t}|\nabla u|^2 - 4R - \frac{2n}{t} + \frac{n}{2}.
\end{aligned}$$

The essential step is the second inequality where we make use of the elementary inequality

$$|u_{ij} - R_{ij} - \frac{1}{t}g_{ij}|^2 \geq \frac{1}{n}\left(\Delta u - R - \frac{n}{t}\right)^2.$$

Now we can apply maximum principle. The value of H for very small positive t is clearly very negative. So we only need to consider the maximum value point is at $t > 0$ for the desired estimate.

For $\forall T_0 < T$, assume that the maximum in $(0, T_0]$ is taken at $t_0 > 0$. At the maximum value point, using the nonnegativity of $|\nabla u|^2$ and R , one has

$$H \leq \frac{-4n + nt_0}{4 + 4t_0} = \frac{n}{4}\left(1 - \frac{5}{t_0 + 1}\right) \leq \frac{n}{4}\left(1 - \frac{5}{T + 1}\right).$$

So if $T \leq 4$, i.e., for time in $[0, 4)$, $H \leq 0$. In general, we have

$$H \leq \frac{n}{4}.$$

Theorem 1.1 is thus proved. □

As a consequence of Theorem 1.1, we have

Corollary 2.1. *Let $(M, g(t))$, $t \in [0, T)$, be a solution to the Ricci flow on a closed manifold, and suppose that $g(0)$ (and so $g(t)$) has weakly positive curvature operator. Let f be a positive solution to the heat equation*

$$\frac{\partial}{\partial t} f = \Delta f - f \ln f + Rf.$$

Assume that (x_1, t_1) and (x_2, t_2) , $0 < t_1 < t_2$, are two points in $M \times (0, T)$. Let

$$\Gamma = \inf_{\gamma} \int_{t_1}^{t_2} e^t \left(|\dot{\gamma}|^2 + R + \frac{2n}{t} + \frac{n}{4} \right) dt,$$

where γ is any space-time path joining (x_1, t_1) and (x_2, t_2) . Then we have

$$e^{t_1} \ln f(x_1, t_1) \leq e^{t_2} \ln f(x_2, t_2) + \frac{\Gamma}{2}.$$

This inequality is in the type of classical Harnack inequalities. The proof is quite standard by integrating the differential Harnack inequality. We include it here for completeness.

Proof. Pick a space-time curve connecting (x_1, t_1) and (x_2, t_2) , $\gamma(t) = (x(t), t)$ for $t \in [t_1, t_2]$. Recall that $u(x, t) = -\ln f(x, t)$. Using the evolution equation for u , we have

$$\begin{aligned} \frac{d}{dt} u(x(t), t) &= \frac{\partial u}{\partial t} + \nabla u \cdot \dot{\gamma} \\ (2.3) \quad &\leq \Delta u - |\nabla u|^2 - R - u + \nabla u \cdot \dot{\gamma} \\ &\leq \Delta u - \frac{|\nabla u|^2}{2} - R - u + \frac{|\dot{\gamma}|^2}{2}. \end{aligned}$$

Now by Theorem 1.1, we have

$$\Delta u = \frac{1}{2} \left(H + |\nabla u|^2 + 3R + \frac{2n}{t} \right) \leq \frac{1}{2} \left(\frac{n}{4} + |\nabla u|^2 + 3R + \frac{2n}{t} \right).$$

So we have the following estimation,

$$\frac{d}{dt} u(x(t), t) \leq \frac{1}{2} \left(|\dot{\gamma}|^2 + R + \frac{2n}{t} + \frac{n}{4} \right) - u.$$

For any space-time curve γ , we arrives at

$$\frac{d}{dt} (e^t \cdot u) \leq \frac{e^t}{2} \left(|\dot{\gamma}|^2 + R + \frac{2n}{t} + \frac{n}{4} \right).$$

Hence the desired Harnack inequality is proved by integrating t from t_1 to t_2 . \square

3. Proof of Theorem 1.2

In this section we study u satisfying the evolution equation (1.6) originated from gradient expanding Ricci soliton equation. We investigate the same quantity

$$H = 2\Delta u - |\nabla u|^2 - 3R - \frac{2n}{t}$$

as in the last section. The evolution equations of u , is still very similar to what is considered in [3]. We will have slightly different terms coming from time derivative $\frac{\partial}{\partial t} u$ when computing the evolution equation satisfied by H . Comparing with [3, Corollary 2.2], we proceed as follows.

Proof of Theorem 1.2.

$$(3.1) \quad \frac{\partial}{\partial t} H = \Delta H - 2\nabla H \cdot \nabla u - 2|u_{ij} - R_{ij} - \frac{1}{t}g_{ij}|^2 - \frac{2}{t}H - \frac{2}{t}|\nabla u|^2 \\ - 2 \left(\frac{\partial}{\partial t} R + \frac{R}{t} + 2\nabla R \cdot \nabla u + 2R_{ij}u_i u_j \right) + \frac{2}{t+2} (-2\Delta u + 2|\nabla u|^2),$$

where the last two terms of the right hand side come from the extra term $-\frac{u}{1+\frac{t}{2}}$ in (1.6). Plugging in $-2\Delta u + 2|\nabla u|^2 = -H + |\nabla u|^2 - 3R - \frac{2n}{t}$, one arrives at

$$(3.2) \quad \frac{\partial}{\partial t} H = \Delta H - 2\nabla H \cdot \nabla u - 2|u_{ij} - R_{ij} - \frac{1}{t}g_{ij}|^2 - \left(\frac{2}{t} + \frac{2}{t+2} \right) H - \frac{6}{t+2} R \\ + \left(\frac{2}{t+2} - \frac{2}{t} \right) |\nabla u|^2 - \frac{4n}{t^2 + 2t} - \left(\frac{\partial}{\partial t} R + \frac{R}{t} + 2\nabla R \cdot \nabla u + 2R_{ij}u_i u_j \right).$$

In sight of the definition of H , for t small enough, we have $H < 0$. Since $g(t)$ has weakly positive curvature operator, by the trace Harnack inequality for the Ricci flow ([8]), we have

$$\frac{\partial}{\partial t} R + \frac{R}{t} + 2\nabla R \cdot \nabla u + 2R_{ij}u_i u_j \geq 0.$$

Notice that now the coefficient for $|\nabla u|^2$ on the right hand side is $\frac{2}{t+2} - \frac{2}{t} < 0$, and we have $R \geq 0$. So one can conclude directly from Maximum Principle that $H \leq 0$. \square

4. A Remark on the Conjugate Heat Equation

In this section we point out a simple observation for [2, Theorem 3.6]. The assumption on scalar curvature is not needed below. We follow the original set-up in [2] here.

Over a closed manifold M^n , $g(t)$, $t \in [0, T]$, is a solution to the Ricci flow (1.1); $f(\cdot, t)$ is a positive solution of the conjugate heat equation

$$(4.1) \quad \frac{\partial f}{\partial t} = -\Delta f + Rf,$$

where Δ and R are Laplacian and scalar curvature with respect to the evolving metric $g(t)$. Notice that $\int_M f(\cdot, t) d\mu_{g(t)}$ is a constant along the flow.

Set

$$v = -\log f - \frac{n \log(4\pi\tau)}{2},$$

where $\tau = T - t$ and define

$$P := 2\Delta v - |\nabla v|^2 + R - \frac{2n}{\tau}.$$

Now we can prove the following result which is closely related to [2, Theorem 3.6].

Theorem 4.1. *Let $(M, g(t))$, $t \in [0, T]$, be a solution to the Ricci flow on a closed manifold. f is a positive solution to the conjugate heat equation (4.1), and v is defines as above. Then we have*

$$\max_M (2\Delta v - |\nabla v|^2 + R)$$

increases along the Ricci flow.

Proof. Following the computation as in [2, Theorem 3.6], one has

$$\frac{\partial P}{\partial \tau} = \Delta P - 2\nabla P \cdot \nabla v - 2|\nabla^2 v + Rc - \frac{1}{\tau}g|^2 - \frac{2}{\tau}P - \frac{2}{\tau}|\nabla v|^2 - \frac{2}{\tau}R.$$

Applying the elementary inequality

$$|\nabla^2 v + Rc - \frac{1}{\tau}g|^2 \geq \frac{1}{n} \left(\Delta v + R - \frac{n}{\tau} \right)^2,$$

and noticing that

$$P + |\nabla v|^2 + R = 2 \left(\Delta v + R - \frac{n}{\tau} \right),$$

we arrive at

$$\begin{aligned} \frac{\partial P}{\partial \tau} &\leq \Delta P - 2\nabla P \cdot \nabla v - \frac{1}{2n}(P + |\nabla v|^2 + R)^2 - \frac{2}{\tau}(P + |\nabla v|^2 + R) \\ &= \Delta P - 2\nabla P \cdot \nabla v - \frac{1}{2n} \left(P + |\nabla v|^2 + R + \frac{2n}{\tau} \right)^2 + \frac{2n}{\tau^2}. \end{aligned}$$

Thus if we define

$$\tilde{P} := P + \frac{2n}{\tau} = 2\Delta v - |\nabla v|^2 + R,$$

we have

$$\frac{\partial \tilde{P}}{\partial \tau} \leq \Delta \tilde{P} - 2\nabla \tilde{P} \cdot \nabla v.$$

Hence $\max_M (2\Delta v - |\nabla v|^2 + R)$ decreases as τ increases, which means that it increases as t increases. This concludes the proof. \square

Remark 4.1. *Notice that we do not need to introduce τ in Theorem 4.1, but we keep the notation here so it is easy to compare with [2, Theorem 3.6].*

Remark 4.2. *Theorem 4.1 and [2, Theorem 3.6] estimate quantities differ by $\frac{2n}{\tau}$. Here we do not need to assume nonnegative scalar curvature as in [2, Theorem 3.6]. Moreover, one can also prove this result for complete non-compact manifolds with proper boundness assumption.*

REFERENCES

- [1] Mihai Bailesteanua, Xiaodong Cao, and Artem Pulemotov. Gradient estimates for the heat equation under the Ricci flow. *J. Funct. Anal.*, doi:10.1016/j.jfa.2009.12.003, 2010.
- [2] Xiaodong Cao. Differential Harnack estimates for backward heat equations with potentials under the Ricci flow. *J. Funct. Anal.*, 255(4):1024–1038, 2008.
- [3] Xiaodong Cao and Richard S. Hamilton. Differential Harnack estimates for time-dependent heat equations with potentials. *Geom. Funct. Anal.*, 19(4):989–1000, 2009.

- [4] Albert Chau, Luen-Fai Tam, and Chengjie Yu. Pseudolocality for the Ricci flow and applications, 2007. [arXiv.org:math/0701153](https://arxiv.org/math/0701153).
- [5] Bennett Chow, Sun-Chin Chu, David Glickenstein, Christine Guenther, James Isenberg, Tom Ivey, Dan Knopf, Peng Lu, Feng Luo, and Lei Ni. *The Ricci flow: techniques and applications. Part I*, volume 135 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2007. Geometric aspects.
- [6] Christine M. Guenther. The fundamental solution on manifolds with time-dependent metrics. *J. Geom. Anal.*, 12(3):425–436, 2002.
- [7] Richard S. Hamilton. Four-manifolds with positive curvature operator. *J. Differential Geom.*, 24(2):153–179, 1986.
- [8] Richard S. Hamilton. The Harnack estimate for the Ricci flow. *J. Differential Geom.*, 37(1):225–243, 1993.
- [9] Richard S. Hamilton. The formation of singularities in the Ricci flow. In *Surveys in differential geometry, Vol. II (Cambridge, MA, 1993)*, pages 7–136. Internat. Press, Cambridge, MA, 1995.
- [10] Shu-Yu Hsu. Gradient estimates for a nonlinear parabolic equation under Ricci flow, 2008. [arXiv.org:0806.4004](https://arxiv.org/0806.4004).
- [11] Shilong Kuang and Qi S. Zhang. A gradient estimate for all positive solutions of the conjugate heat equation under Ricci flow. *J. Funct. Anal.*, 255(4):1008–1023, 2008.
- [12] Peter Li and Shing-Tung Yau. On the parabolic kernel of the Schrödinger operator. *Acta Math.*, 156(3-4):153–201, 1986.
- [13] Shiping Liu. Gradient estimate for solutions of the heat equation under Ricci flow. *Pacific J. Math.*, 243(1):165–180, 2009.
- [14] Li Ma. Gradient estimates for a simple elliptic equation on complete non-compact Riemannian manifolds. *J. Funct. Anal.*, 241(1):374–382, 2006.
- [15] Lei Ni. Ricci flow and nonnegativity of sectional curvature. *Math. Res. Lett.*, 11(5-6):883–904, 2004.
- [16] Lei Ni. Monotonicity and Li-Yau-Hamilton inequalities. In *Surveys in differential geometry. Vol. XII. Geometric flows*, volume 12 of *Surv. Differ. Geom.*, pages 251–301. Int. Press, Somerville, MA, 2008.
- [17] Grisha Perelman. The entropy formula for the Ricci flow and its geometric applications, 2002. [arXiv:math.DG/0211159](https://arxiv.org/math.DG/0211159).
- [18] Jun Sun. Gradient estimates for positive solutions of the heat equation under geometric flow, 2009. **Preprint**.
- [19] Yunyan Yang. Gradient estimates for a nonlinear parabolic equation on Riemannian manifolds. *Proc. Amer. Math. Soc.*, 136(11):4095–4102, 2008.
- [20] Qi S. Zhang. Some gradient estimates for the heat equation on domains and for an equation by Perelman. *Int. Math. Res. Not.*, pages Art. ID 92314, 39, 2006.

DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, ITHACA, NY 14853-4201, USA.

E-mail address: cao@math.cornell.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, AT ANN ARBOR, MI 48109, USA.

E-mail address: zhangou@umich.edu