

# Chocolate consumption and risk of diabetes mellitus in the Physicians' Health Study<sup>1–4</sup>

Chisa Matsumoto, Andrew B Petrone, Howard D Sesso, J Michael Gaziano, and Luc Djoussé

## ABSTRACT

**Background:** Previous studies reported beneficial effects of cocoa or chocolate on insulin resistance, oxidative stress, and inflammation, which are important risk factors of type 2 diabetes mellitus (DM). However, it is unclear whether chocolate consumption is associated with risk of DM.

**Objective:** We tested the hypothesis that chocolate consumption is inversely associated with incident DM in the Physicians' Health Study (PHS).

**Design:** We prospectively analyzed data on 18,235 PHS participants who were free of DM at baseline (1997–2001). Chocolate consumption was obtained from a baseline food-frequency questionnaire. Incident DM was ascertained via annual follow-up questionnaires and validated in a subsample by a review of medical records. We used Cox proportional hazards models to estimate HRs and 95% CIs of DM.

**Results:** The mean ( $\pm$ SD) age at baseline was  $66.3 \pm 9.2$  y. During a mean follow up of 9.2 y, 1123 men (6.2%) developed DM. For self-reported chocolate consumption of none, 1–3 servings/mo, 1 serving/wk, and  $\geq 2$  servings/wk, multivariable-adjusted HRs (95% CIs) of DM adjusted for lifestyle, clinical, and dietary risk factors including total energy intake were 1.00 (referent), 0.93 (0.79, 1.09), 0.86 (0.72, 1.04), and 0.83 (0.69, 0.99), respectively ( $P$ -trend = 0.047). In secondary analyses, the inverse association of chocolate consumption and risk of DM was slightly stronger in subjects without a history of cardiovascular disease or heart failure ( $P$ -trend = 0.023). In addition, both age and BMI modified the chocolate-DM relation ( $P < 0.05$  each).

**Conclusion:** Our data support an inverse relation of chocolate intake with incident DM, which appears only to apply in younger and normal-body weight men after controlling for comprehensive life styles including total energy consumption. *Am J Clin Nutr* 2015;101:362–7.

**Keywords** chocolate, diabetes mellitus, nutrition, epidemiology, risk factors

## INTRODUCTION

Type 2 diabetes mellitus (DM)<sup>5</sup> has tremendous worldwide social and economic burden partly because of its complications, including cardiovascular disease (CVD), renal dysfunction, retinopathy, or diabetic cardiomyopathy (1). The prevalence of DM is increasing rapidly and estimated to rise from 171 million cases in 2000 to 366 million cases in 2030 (2). Therefore, it is important to identify modifiable risk factors of DM such as diet

and determine beneficial foods that could help reduce risk of DM.

Chocolate contains caffeine, flavonoids (e.g., flavanols and procyanidins), and minerals such as magnesium (3) that could influence risk of DM. An inverse association between the consumption of caffeine, flavonoids, and magnesium with risk of DM has been reported with inconsistent results (4–9). In contrast, cocoa and chocolate have beneficial effects on insulin resistance (10–12), oxidative stress (12–14), and inflammation (15, 16), which play important roles in the pathogenesis of DM, and some observational studies suggested an inverse association of chocolate intake and risk of DM (8, 17). However, it is still unclear whether chocolate consumption is associated with risk of DM and whether this association differs according to characteristics of the study population. Hence, we sought to examine whether chocolate consumption is inversely associated with incident DM in the Physicians' Health Study (PHS).

## SUBJECTS AND METHODS

### Study population

Study subjects were selected from 29,071 participants in the PHS I (clinicaltrials.gov; NCT00000500) and PHS II (clinicaltrials.gov; NCT00270647). The PHS I is a completed randomized trial of the efficacy of low-dose aspirin and  $\beta$ -carotene

<sup>1</sup>From the Division of Cardiology, Tokyo Medical University, Tokyo, Japan (CM); the Divisions of Aging (ABP, HDS, JMG, and LD) and Preventive Medicine (HDS and JMG), Brigham and Women's Hospital, Boston, MA; the Harvard Medical School, Boston, MA (HDS, JMG, and LD); and the Boston Veterans Affairs Healthcare System, Boston, MA (JMG and LD).

<sup>2</sup>Supported by grant R21 HL088081 from the National Heart, Lung, and Blood Institute, Bethesda, MD. The Physicians' Health Study was supported by grants CA-34944, CA-40360, and CA-097193 from the National Cancer Institute and grants HL-26490 and HL-34595 from the National Heart, Lung, and Blood Institute, Bethesda, MD.

<sup>3</sup>Study sponsors had no role in the design and conduct of the study, collection, management, analysis, and interpretation of data, and preparation, review, or approval of the manuscript.

<sup>4</sup>Address correspondence to C Matsumoto, Division of Cardiology, Tokyo Medical University, 6-7-1 Nishishinjyuku Shinjyuku-ku, Tokyo, Japan, 160-0023. E-mail: chisamatsumoto@post.harvard.edu.

<sup>5</sup>Abbreviations used: CVD, cardiovascular disease; DM, type 2 diabetes mellitus; FFQ, food-frequency questionnaire; PHS, Physicians' Health Study.

Received June 10, 2014. Accepted for publication November 10, 2014.

First published online November 26, 2014; doi: 10.3945/ajcn.114.092221.

on CVD and cancer in 22,071 US male physicians aged 40–84 y at baseline (1982). The PHS II was a randomized trial designed to assess the effects of  $\beta$ -carotene, vitamin E, vitamin C, and a multivitamin on risk of cancer, CVD, and other chronic diseases in 14,641 US male physicians aged  $\geq 50$  y at baseline (1997). Detailed descriptions of both studies have been published previously (18, 19). For the current analysis, we started follow-up when the food-frequency questionnaire (FFQ) was assessed (1997–2001).

In PHS participants free of baseline DM ( $n = 19,465$ ), we excluded 342 participants with missing data on chocolate consumption and 1105 participants whose reported total energy intakes were implausible ( $< 500$  or  $> 3500$  kcal/d or missing). Thus, a total of 18,235 men were used for the current analyses.

### Ethics

Each participant gave written informed consent, and the Brigham and Women's Hospital Institutional Review Board approved the study protocol.

### Ascertainment of DM, chocolate consumption, and other covariates

DM was ascertained by self-reports on annual follow-up questionnaires and validated in a subsample of participants through a review of medical records with a positive predictive value of self-reported DM of 98.3% (20). Chocolate consumption was obtained from a baseline semiquantitative FFQ (1997–2001). In a single question, participants were asked to answer "Please fill in your average use, during the year, of each specified food; chocolate (1 oz)." Possible answers included never or  $< 1$  serving/mo, 1–3 servings/mo, 1, 2–4, and 5–6 servings/wk, and 1, 2–3, 4–5, and  $\geq 6$  servings/d. Other dietary factors (whole grain, red meat, eggs, nuts, and candies without chocolate) were assessed via the FFQ. Glycemic index, fiber, magnesium, caffeine, and flavonoids were computed for each food item on the FFQ by using the food-composition database from the Harvard School of Public Health and manufacturer information. The validity and reproducibility of the FFQ in health professionals have been reported previously (21, 22). We used the residual method to adjust nutrients for energy intake (23).

Other baseline covariates included cohort status (recruited in 1982 and participated in the PHS II, recruited in 1982 and did not participate in the PHS II, and recruited in 1997), age, BMI, smoking (never, past, and current), vigorous exercise ( $< 1$  and  $\geq 1$  time/wk), alcohol consumption (rarely or never, 1 drink/mo to 1 drink/wk, 2–6 drinks/wk, and  $\geq 1$  drink/d), history of hypercholesterolemia (no or yes), and history of hypertension (no or yes).

### Statistics

Chocolate consumption was classified into the following 4 categories: never or rarely (subsequently referred to as "none"), 1–3 servings/mo, and 1 and 2 servings/wk (we merged the last 6 extreme categories because of small sample sizes in these groups). There were missing data on exercise ( $n = 291$ ), alcohol intake ( $n = 100$ ), and intakes of whole grain ( $n = 26$ ), red meat ( $n = 17$ ), and nuts ( $n = 172$ ). We created indicator variables for missing variables and included them in the multivariable model.

We presented baseline characteristics across categories of chocolate consumption by using generalized linear models for continuous variables and chi-square tests for categorical variables. The person-time of follow-up was computed from baseline until the first occurrence of DM, death, or the date of last known contact. We used Cox proportional hazards models to estimate HRs with corresponding 95% CIs. An initial model adjusted for age; a second model further controlled for cohort status, smoking status (never, current, and past), vigorous exercise ( $< 1$  and  $\geq 1$  time/wk), alcohol intake (rarely or never, 1 drink/mo to 1 drink/wk, 2–6 drinks/wk, and  $\geq 1$  drink/d), and BMI (continuous) (model 2). The final model adjusted for covariates in model 2 plus total caloric intake (in quartiles), whole grain (in quartiles), red meat (in quartiles), and nut (in quartiles) intakes (model 3). Covariates were selected on the basis of knowledge from previous studies (18, 24, 25) or if the regression coefficient for chocolate consumption ( $\geq 2$  servings/wk) changed by 10% when adding potential confounders to a model singly or jointly (26). We also considered additional adjustment beyond model 3 for other dietary factors, including nonchocolate candy, eggs, glycemic index, fiber, magnesium, caffeine, and flavonoids. We also considered a multivariable model adjusted for potential mediators such as a history of hypertension and hypercholesterolemia. We tested for a linear trend by using median servings per day in each category of chocolate as a continuous variable.

In secondary analyses, we repeated analyses in 15,999 men free of prevalent CVD (myocardial infarction, percutaneous transluminal coronary angioplasty, coronary artery bypass graft, and stroke) or heart failure at baseline because these comorbidities may have led to a change in dietary habits. We evaluated the effect modification by age ( $< 65$  and  $\geq 65$  y), BMI (in  $\text{kg}/\text{m}^2$ ) ( $< 25$  and  $\geq 25$ ), and history of hypercholesterolemia. We performed several sensitivity analyses to assess the robustness of our findings. We excluded men with a history of cancer (except skin cancer) because these men may have changed their dietary habits, and some epidemiologic studies suggested significant association of DM and risk of cancer (27, 28). Furthermore, we excluded men with missing values on the major covariates vigorous exercise, alcohol intake, smoking, and intakes of whole grain, red meat, and nuts. For each of these sensitivity analyses, results were generally similar to main findings and are not presented. All analyses were completed with SAS software (version 9.3; SAS Institute). All statistical tests were 2-sided, and  $P < 0.05$  was considered significant.

### RESULTS

The mean ( $\pm$ SD) age at baseline was  $66.3 \pm 9.2$  y. Baseline characteristics are summarized in **Table 1** by amount of chocolate consumption. Higher intake of chocolate was not associated with a significant trend for higher intake of energy-adjusted micronutrients such as caffeine and flavonoids. During a mean follow up of 9.2 y, 1123 men (6.2%) developed DM. For self-reported chocolate consumption of none, 1–3 servings/mo, and 1 and  $\geq 2$  servings/wk, multivariable-adjusted HRs (95% CI) of DM were 1.00 (reference), 0.93 (0.79, 1.09), 0.86 (0.72, 1.04), and 0.83 (0.69, 0.99), respectively ( $P$ -trend = 0.047) (**Table 2**, model 3). Adjustment for other dietary factors did not alter

**TABLE 1**Distribution of baseline characteristics of physicians according to chocolate intake assessed by FFQ<sup>1</sup>

| Variables                             | Average frequency of chocolate intake |                 |              |                          |
|---------------------------------------|---------------------------------------|-----------------|--------------|--------------------------|
|                                       | None                                  | 1–3 servings/mo | 1 serving/wk | ≥2 servings/wk           |
| Participants, <i>n</i>                | 4432                                  | 5349            | 3649         | 4805                     |
| Age, y                                | 67.0 ± 8.8                            | 66.0 ± 9.0      | 66.1 ± 9.2   | 66.3 ± 9.9 <sup>2</sup>  |
| BMI, kg/m <sup>2</sup>                | 25.5 ± 3.2                            | 25.8 ± 3.3      | 25.7 ± 3.1   | 25.6 ± 3.2 <sup>2</sup>  |
| Smoking status, <sup>2</sup> %        |                                       |                 |              |                          |
| Past                                  | 46.0                                  | 42.8            | 39.3         | 39.0                     |
| Never                                 | 50.3                                  | 54.2            | 57.5         | 57.6                     |
| Current                               | 3.5                                   | 3.0             | 3.0          | 3.3                      |
| Vigorous exercise, %                  |                                       |                 |              |                          |
| <1 time/wk                            | 36.8                                  | 36.5            | 35.1         | 35.9                     |
| ≥1 time/wk                            | 61.7                                  | 62.1            | 63.3         | 62.2                     |
| Alcohol intake, <sup>2</sup> %        |                                       |                 |              |                          |
| Rarely, never                         | 15.3                                  | 14.6            | 16.5         | 18.9                     |
| Monthly                               | 6.1                                   | 6.9             | 7.8          | 8.6                      |
| Weekly                                | 35.2                                  | 38.5            | 39.5         | 39.3                     |
| Daily                                 | 43.0                                  | 39.5            | 35.6         | 32.6                     |
| Total calories, kcal/d                | 1528 ± 462                            | 1617 ± 471      | 1695 ± 481   | 1863 ± 518 <sup>2</sup>  |
| Whole grain, servings/d               | 1.7 ± 1.7                             | 1.8 ± 1.7       | 1.8 ± 1.6    | 1.7 ± 1.6 <sup>2</sup>   |
| Red meat, servings/d                  | 0.6 ± 0.6                             | 0.7 ± 0.6       | 0.7 ± 0.5    | 0.8 ± 0.6 <sup>2</sup>   |
| Eggs, servings/d                      | 1.7 ± 1.4                             | 1.8 ± 1.3       | 1.9 ± 1.3    | 2.0 ± 1.3 <sup>2</sup>   |
| Nuts, servings/d                      | 1.2 ± 1.3                             | 1.4 ± 1.3       | 1.6 ± 1.3    | 1.8 ± 1.4 <sup>2</sup>   |
| Candies without chocolate, servings/d | 0.4 ± 1.0                             | 1.0 ± 1.1       | 1.5 ± 1.2    | 1.7 ± 1.5 <sup>2</sup>   |
| Carbohydrate, g/d                     | 216 ± 51                              | 215 ± 43        | 215 ± 41     | 211 ± 37 <sup>2</sup>    |
| Glycemic index                        | 107 ± 15                              | 107 ± 13        | 107 ± 14     | 106 ± 14 <sup>2</sup>    |
| Total fat, g/d                        | 47.8 ± 14.4                           | 50.5 ± 12.5     | 52.3 ± 11.9  | 55.7 ± 11.5 <sup>2</sup> |
| Fiber, g/d                            | 19.8 ± 7.4                            | 19.3 ± 6.4      | 18.9 ± 5.9   | 18.0 ± 5.5 <sup>2</sup>  |
| Caffeine, mg/d                        | 195 ± 150                             | 199 ± 151       | 192 ± 148    | 185 ± 144 <sup>2</sup>   |
| Total flavonoids, mg/d                | 345 ± 345                             | 312 ± 288       | 298 ± 277    | 294 ± 249 <sup>2</sup>   |
| Magnesium, mg/d                       | 336 ± 95                              | 325 ± 81        | 318 ± 73     | 307 ± 70 <sup>2</sup>    |
| History of hypercholesterolemia, %    | 42.8                                  | 39.8            | 38.4         | 37.8 <sup>2</sup>        |
| History of hypertension, %            | 44.3                                  | 42.3            | 41.0         | 39.7 <sup>2</sup>        |
| History of cardiovascular disease, %  | 13.4                                  | 11.9            | 9.9          | 10.2 <sup>2</sup>        |
| History of heart failure, %           | 1.6                                   | 1.7             | 1.7          | 2.0                      |
| History of cancer, %                  | 11.7                                  | 10.7            | 11.5         | 11.8                     |

<sup>1</sup>All values for continuous variables are means ± SDs, and all values for categorical variables are proportions. All nutrients are presented as energy adjusted values. Data were missing for BMI (*n* = 3), vigorous exercise (*n* = 291), alcohol intake (*n* = 100), history of hypercholesterolemia (*n* = 439), history of hypertension (*n* = 89), and intake of whole grain (*n* = 26), red meat (*n* = 17), eggs (*n* = 95), nuts (*n* = 172), and candies without chocolate (*n* = 280). FFQ, food-frequency questionnaire.

<sup>2</sup>*P* < 0.05, across categories of chocolate. A generalized linear model was used for continuous variables, and the chi-square test was used for categorical variables.

the main results (data not shown). As expected, after additional adjustment for potential intermediate factors (history of hypertension and hypercholesterolemia), there was a slight attenuation of the HR (95% CI) from 0.83 (0.69, 0.99) to 0.86

(0.72, 1.03) for ≥2 chocolate servings/wk compared with non-consumers (*P*-trend = 0.11).

In secondary analyses, the exclusion of men without history of CVD or heart failure led to a slightly stronger inverse association

**TABLE 2**HRs (95% CIs) for DM by category of chocolate intake in 18,235 male physicians<sup>1</sup>

| Frequency of chocolate intake | <i>n</i> | Cases, <i>n</i> (%) | Incidence rate/1000 PY | HRs (95% CIs) for DM |                      |                      |
|-------------------------------|----------|---------------------|------------------------|----------------------|----------------------|----------------------|
|                               |          |                     |                        | Age-adjusted model 1 | Multivariate model 2 | Multivariate model 3 |
| None                          | 4432     | 275 (6.2)           | 6.9                    | Referent             | Referent             | Referent             |
| 1–3 servings/mo               | 5349     | 345 (6.5)           | 7.0                    | 1.02 (0.87, 1.19)    | 0.97 (0.83, 1.14)    | 0.93 (0.79, 1.09)    |
| 1 serving/wk                  | 3649     | 217 (6.0)           | 6.4                    | 0.93 (0.78, 1.11)    | 0.91 (0.76, 1.08)    | 0.86 (0.72, 1.04)    |
| ≥2 servings/wk                | 4805     | 286 (6.0)           | 6.5                    | 0.94 (0.80, 1.11)    | 0.89 (0.75, 1.05)    | 0.83 (0.69, 0.99)    |
| <i>P</i> -linear trend        | —        | —                   | —                      | 0.39                 | 0.17                 | 0.047                |

<sup>1</sup>Cox proportional hazards models were used to estimate HRs (95% CIs). Model 1 was adjusted for age. Model 2 was adjusted as for model 1 and for cohort status, BMI, smoking status, exercise, and alcohol consumption. Model 3 was adjusted as for model 2 and for total caloric intake and intakes of whole grains, nuts, and red meat. DM, type 2 diabetes mellitus; PY, person-years.

**TABLE 3**

HRs (95% CIs) for DM by category of chocolate intake in 15,999 physicians without cardiovascular disease or heart failure at baseline<sup>1</sup>

| Frequency of chocolate intake | n    | Cases, n (%) | Incidence rate/1000 PY | HRs (95% CIs) for DM |                      |                      |
|-------------------------------|------|--------------|------------------------|----------------------|----------------------|----------------------|
|                               |      |              |                        | Age-adjusted model 1 | Multivariate model 2 | Multivariate model 3 |
| None                          | 3806 | 240 (6.3)    | 6.8                    | Referent             | Referent             | Referent             |
| 1–3 servings/mo               | 4672 | 280 (6.0)    | 6.3                    | 0.93 (0.79, 1.11)    | 0.88 (0.74, 1.05)    | 0.85 (0.72, 1.02)    |
| 1 serving/wk                  | 3250 | 182 (5.6)    | 5.9                    | 0.86 (0.71, 1.05)    | 0.84 (0.69, 1.02)    | 0.81 (0.66, 0.98)    |
| ≥2 servings/wk                | 4271 | 241 (5.6)    | 6.0                    | 0.88 (0.73, 1.05)    | 0.82 (0.68, 0.98)    | 0.77 (0.63, 0.93)    |
| <i>P</i> -linear trend        | —    | —            | —                      | 0.21                 | 0.07                 | 0.02                 |

<sup>1</sup>Cox proportional hazards models were used to estimate HRs (95% CIs). Model 1 was adjusted for age. Model 2 was adjusted as for model 1 and for cohort status, BMI, smoking status, exercise, and alcohol consumption. Model 3 was adjusted as for model 2 and for total caloric intake and intakes of whole grains, nuts, and red meat. DM, type 2 diabetes mellitus; PY, person-years.

between chocolate consumption and risk of DM [HR (95% CI) for ≥2 chocolate servings/wk: 0.77 (0.63, 0.93)] (Table 3). No significant association between chocolate consumption and DM was observed in 2236 men with CVD or heart failure (*P*-trend = 0.43).

There was a significant effect modification by BMI (*P*-interaction = 0.04) (Table 4), age, and chocolate intake on DM risk (*P*-interaction = 0.03) (Table 4). In subgroup analysis, only men with normal BMI (<25) or aged <65 y showed an inverse association of chocolate consumption and risk of DM. A history of hypercholesterolemia did not modify the association between chocolate and DM (*P*-interaction = 0.65).

**DISCUSSION**

In this prospective study, self-reported chocolate intake was significantly and inversely associated with incident DM and such relation appeared to be limited to normal-body weight and younger men after comprehensive adjustment for lifestyle, clinical, and dietary risk factors.

Other studies supported our finding that chocolate consumption may lower risk of DM (8, 17). In a Japanese observational study, the consumption of ≥1 serving chocolate/wk compared with none was inversely associated with risk of DM in men [HR (95% CI): 0.65 (0.43, 0.97)] (8). The Japanese population was

**TABLE 4**

HRs (95% CIs) for DM by category of chocolate intake stratified by BMI and age<sup>1</sup>

| Frequency of chocolate intake                             | n    | Cases, n (%) | Incidence rate/1000 PY | HRs (95% CIs) for DM |                      |                      |
|---|------|--------------|------------------------|----------------------|----------------------|----------------------|
|   |      |              |                        | Age-adjusted model 1 | Multivariate model 2 | Multivariate model 3 |
| <b>BMI &lt;25 kg/m<sup>2</sup> (n = 8218)<sup>2</sup></b> |      |              |                        |                      |                      |                      |
| None  | 2059 | 78 (3.8)     | 4.2                    | Referent             | Referent             | Referent             |
| 1–3 servings/mo   | 2332 | 80 (3.4)     | 3.7                    | 0.88 (0.65, 1.21)    | 0.86 (0.63, 1.17)    | 0.84 (0.61, 1.15)    |
| 1 serving/wk  | 1620 | 55 (3.4)     | 3.6                    | 0.87 (0.62, 1.23)    | 0.82 (0.58, 1.16)    | 0.81 (0.57, 1.15)    |
| ≥2 servings/wk  | 2207 | 57 (2.6)     | 2.8                    | 0.67 (0.47, 0.94)    | 0.62 (0.44, 0.87)    | 0.59 (0.42, 0.85)    |
| <i>P</i> -linear trend                                    | —    | —            | —                      | 0.02                 | 0.007                | 0.005                |
| <b>BMI ≥25 kg/m<sup>2</sup> (n = 10,014)<sup>2</sup></b>  |      |              |                        |                      |                      |                      |
| None  | 2373 | 197 (8.3)    | 9.1                    | Referent             | Referent             | Referent             |
| 1–3 servings/mo   | 3016 | 265 (8.8)    | 9.5                    | 1.05 (0.87, 1.26)    | 1.02 (0.85, 1.23)    | 0.97 (0.81, 1.18)    |
| 1 serving/wk  | 2028 | 162 (8.0)    | 8.6                    | 0.94 (0.77, 1.16)    | 0.94 (0.76, 1.16)    | 0.90 (0.72, 1.11)    |
| ≥2 servings/wk  | 2597 | 229 (8.8)    | 9.6                    | 1.06 (0.87, 1.28)    | 1.01 (0.83, 1.22)    | 0.93 (0.76, 1.15)    |
| <i>P</i> -linear trend                                    | —    | —            | —                      | 0.64                 | 0.97                 | 0.57                 |
| <b>Age &lt;65 y (n = 8736)<sup>3</sup></b>                |      |              |                        |                      |                      |                      |
| None  | 2019 | 139 (6.9)    | 7.0                    | Referent             | Referent             | Referent             |
| 1–3 servings/mo   | 2601 | 179 (6.9)    | 6.9                    | 1.01 (0.81, 1.26)    | 0.98 (0.79, 1.23)    | 0.93 (0.74, 1.17)    |
| 1 serving/wk  | 1808 | 114 (6.3)    | 6.3                    | 0.91 (0.71, 1.17)    | 0.91 (0.71, 1.17)    | 0.87 (0.68, 1.13)    |
| ≥2 servings/wk  | 2308 | 134 (5.8)    | 5.7                    | 0.84 (0.66, 1.06)    | 0.78 (0.61, 0.99)    | 0.73 (0.57, 0.94)    |
| <i>P</i> -linear trend                                    | —    | —            | —                      | 0.08                 | 0.02                 | 0.01                 |
| <b>Age ≥65 y (n = 9499)<sup>3</sup></b>                   |      |              |                        |                      |                      |                      |
| None  | 2413 | 136 (5.6)    | 6.7                    | Referent             | Referent             | Referent             |
| 1–3 servings/mo   | 2748 | 166 (6.0)    | 7.0                    | 1.04 (0.83, 1.30)    | 0.99 (0.79, 1.24)    | 0.96 (0.76, 1.21)    |
| 1 serving/wk  | 1841 | 103 (5.6)    | 6.5                    | 0.96 (0.74, 1.24)    | 0.93 (0.72, 1.21)    | 0.90 (0.69, 1.17)    |
| ≥2 servings/wk  | 2497 | 152 (6.1)    | 7.3                    | 1.09 (0.87, 1.38)    | 1.06 (0.84, 1.33)    | 0.98 (0.77, 1.25)    |
| <i>P</i> -linear trend                                    | —    | —            | —                      | 0.44                 | 0.54                 | 0.96                 |

<sup>1</sup>Cox proportional hazards models were used to estimate HRs (95% CIs). Model 1 was adjusted for age. Model 2 was adjusted as for model 1 and for cohort status, BMI, smoking status, exercise, and alcohol consumption. Model 3 was adjusted as for model 2 and for total caloric intake and intakes of whole grains, nuts, and red meat. DM, type 2 diabetes mellitus; PY, person-years.

<sup>2</sup>*P*-interaction = 0.04.

<sup>3</sup>*P*-interaction = 0.03.

younger (mean age: 57 y) than our population (mean age: 66 y); furthermore, BMI was lower in Japanese subjects (mean BMI: 22.3) than the PHS cohort (mean BMI: 25.7). A stronger association observed in PHS subjects who were <65 y old and those with normal BMI was consistent with Japanese results.

A recent prospective analysis of the Atherosclerosis Risk in Communities cohort showed that, in men, consumption of chocolate was associated with 29% (95% CI: 10%, 64%) lower risk of DM for <1 serving/mo, 44% (24%, 59%) lower risk of DM for 1–4 servings/wk, and 32% (95% CI: –6%, 56%) lower risk of DM for  $\geq 2$  servings/wk (17). As with the Japanese study, this relation was stronger than in our study, possibly explained by a younger population (mean age: 53–54 y across categories of chocolate consumption). However, BMI was higher in the Atherosclerosis Risk in Communities study (mean BMI: 27.2–27.8), and results did not change after adjustment for BMI (17).

What might explain a slightly stronger inverse association between chocolate consumption and risk of DM after the exclusion of men with CVD and heart failure? CVD or heart failure may lead to dietary changes that confound the association of chocolate consumption with risk of DM. It is also possible that competing causes of death or different risk factors in overweight or obese men might have increased the background rate of DM, thereby making it more difficult to detect a small increase in risk. In fact, the incidence rate of DM in men with BMI  $\geq 25$  was almost 3 times higher than that of men with normal BMI, especially in the higher chocolate-consumption group (Table 4). In our study, potential benefits of chocolate consumption were limited to individuals with BMI <25 and age <65 y, which suggested that the benefits of chocolate could be limited to this specific subgroup. Lastly, the inverse association of chocolate consumption and risk of DM was only significant when it was controlled for comprehensive life style factors including total energy consumption.

#### Possible mechanisms by which chocolate consumption may influence DM risk

Beneficial effects of cocoa and chocolate on risk factors related to DM were consistently reported in short-term randomized clinical trials as well as long-term observational and experimental studies. For example, cocoa or chocolate improves insulin resistance (12–14), oxidative stress (14–16), and inflammation (17, 18). Whether these results and possible mechanisms extend to an inverse association between chocolate consumption and risk of DM remains unclear.

Also uncertain is which substances in chocolate are responsible for any potential inverse association. Results from previous observational studies that evaluated the association of flavonoids or caffeine with risk of DM have been inconsistent (5–9). Adjustment for these and other key dietary components of chocolate did not greatly affect HRs of DM (data not shown).

#### Strengths and limitations

Our study has several strengths, including a large sample size, long follow-up, and valid ascertainment of DM. Our cohort study has some limitations. First, we were not able to differentiate types of chocolates consumed. Nevertheless, dark chocolate contains greater amounts of cocoa and higher flavonoid concentrations

than does milk or white chocolate (29). Also, cocoa concentrations of chocolate in the market has a wide range [i.e., the required cocoa concentration of milk chocolate produced in the United States is lower (>10%) than for milk chocolate produced in European countries (>25%)] (30, 31). And, this could have led to an underestimation or overestimation of the true association between chocolate consumption and risk of DM. Second, our sample consisted of highly educated male physicians, possibly limiting the generalizability of our findings to other socioeconomic or ethnic groups and women. Additional studies are needed in other ethnic groups and people with lower socioeconomic status to confirm the current findings. Third, unmeasured and residual confounding could have partially explained our findings. Fourth, chocolate was self-reported in this cohort, and we could not exclude a reporting bias in our data.

In conclusion, our data are consistent with an inverse association between chocolate consumption and incident DM, which appeared only to apply in younger and normal-body weight men after controlling for comprehensive lifestyle factors.

The authors' responsibilities were as follows—CM and LD: study design and conception and statistical analyses; CM, ABP, and LD: drafting of the manuscript; LD: obtained funding, collected data, and supervised the work; HDS: collected data, obtained funding for PHS, and critically reviewed manuscript for content; and JMG: obtained funding, collected data for PHS, and critically reviewed manuscript for content. LD received investigator-initiated grants from GlaxoSmithKline and the California Walnut Commission. CM, ABP, HDS, and JMG had no conflicts of interest.

#### REFERENCES

- Bruno G, Landi A. Epidemiology and costs of diabetes. *Transplant Proc* 2011;43:327–9.
- Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care* 2004;27:1047–53.
- The National Agricultural Library. National Nutrient Database for Standard Reference [cited 26 Sep 2014]. Available from: <http://ndb.nal.usda.gov/ndb/search/list>.
- Dong JY, Xun P, He K, Qin LQ. Magnesium intake and risk of type 2 diabetes: meta-analysis of prospective cohort studies. *Diabetes Care* 2011;34:2116–22.
- Iso H, Date C, Wakai K, Fukui M, Tamakoshi A. The relationship between green tea and total caffeine intake and risk for self-reported type 2 diabetes among Japanese adults. *Ann Intern Med* 2006;144:554–62.
- Wedick NM, Pan A, Cassidy A, Rimm EB, Sampson L, Rosner B, Willett W, Hu FB, Sun Q, van Dam RM. Dietary flavonoid intakes and risk of type 2 diabetes in US men and women. *Am J Clin Nutr* 2012;95:925–33.
- Knekt P, Kumpulainen J, Jarvinen R, Rissanen H, Heliovaara M, Reunanen A, Hakulinen T, Aromaa A. Flavonoid intake and risk of chronic diseases. *Am J Clin Nutr* 2002;76:560–8.
- Oba S, Nagata C, Nakamura K, Fujii K, Kawachi T, Takatsuka N, Shimizu H. Consumption of coffee, green tea, oolong tea, black tea, chocolate snacks and the caffeine content in relation to risk of diabetes in Japanese men and women. *Br J Nutr* 2010;103:453–9.
- van Dam RM, Willett WC, Manson JE, Hu FB. Coffee, caffeine, and risk of type 2 diabetes: a prospective cohort study in younger and middle-aged U.S. women. *Diabetes Care* 2006;29:398–403.
- Curtis PJ, Sampson M, Potter J, Dhatariya K, Kroon PA, Cassidy A. Chronic ingestion of flavan-3-ols and isoflavones improves insulin sensitivity and lipoprotein status and attenuates estimated 10-year CVD risk in medicated postmenopausal women with type 2 diabetes: a 1-year, double-blind, randomized, controlled trial. *Diabetes Care* 2012;35:226–32.
- Grassi D, Lippi C, Necozione S, Desideri G, Ferri C. Short-term administration of dark chocolate is followed by a significant increase in insulin sensitivity and a decrease in blood pressure in healthy persons. *Am J Clin Nutr* 2005;81:611–4.

12. Jalil AMM, Ismail A, Pei CP, Hamid M, Kamaruddin SHS. Effects of cocoa extract on glucometabolism, oxidative stress, and antioxidant enzymes in obese-diabetic (Ob-db) rats. *J Agric Food Chem* 2008;56:7877–84.
13. Davison G, Callister R, Williamson G, Cooper KA, Gleeson M. The effect of acute pre-exercise dark chocolate consumption on plasma antioxidant status, oxidative stress and immunoendocrine responses to prolonged exercise. *Eur J Nutr* 2012;51:69–79.
14. Dreosti IE. Antioxidant polyphenols in tea, cocoa, and wine. *Nutrition* 2000;16:692–4.
15. Selmi C, Cocchi CA, Lanfredini M, Keen CL, Gershwin ME. Chocolate at heart: the anti-inflammatory impact of cocoa flavanols. *Mol Nutr Food Res* 2008;52:1340–8.
16. Sies H, Schewe T, Heiss C, Kelm M. Cocoa polyphenols and inflammatory mediators. *Am J Clin Nutr* 2005;81:304S–12S.
17. Greenberg JA. Chocolate intake and diabetes risk. *Clin Nutr* 2014 Feb 17 (Epub ahead of print; DOI: 10.1016/j.clnu.2014.02.005).
18. Steering Committee of the Physicians' Health Study Research Group. Final report on the aspirin component of the ongoing Physicians' Health Study. *N Engl J Med* 1989;321:129–35.
19. Christen WG, Gaziano JM, Hennekens CH. Design of Physicians' Health Study II—a randomized trial of beta-carotene, vitamins E and C, and multivitamins, in prevention of cancer, cardiovascular disease, and eye disease, and review of results of completed trials. *Ann Epidemiol* 2000;10:125–34.
20. Djoussé L, Driver J, Gaziano J, Buring J, Lee I. Association between modifiable lifestyle factors and residual lifetime risk of diabetes. *Nutr Metab Cardiovasc Dis* 2013;23:17–221.
21. Rimm EB, Giovannucci EL, Stampfer MJ, Colditz GA, Litin LB, Willett WC. Reproducibility and validity of an expanded self-administered semiquantitative food frequency questionnaire among male health professionals. *Am J Epidemiol* 1992;135:1114–26.
22. Willett WC, Sampson L, Stampfer MJ, Rosner B, Bain C, Witschi J, Hennekens CH, Speizer FE. Reproducibility and validity of a semi-quantitative food frequency questionnaire. *Am J Epidemiol* 1985;122:51–65.
23. Hu FB, Stampfer MJ, Rimm E, Ascherio A, Rosner BA, Spiegelman D, Willett WC. Dietary fat and coronary heart disease: a comparison of approaches for adjusting for total energy intake and modeling repeated dietary measurements. *Am J Epidemiol* 1999;149:531–40.
24. Kochar J, Gaziano JM, Djoussé L. Nut consumption and risk of type II diabetes in the Physicians' Health Study. *Eur J Clin Nutr* 2010;64:75–9.
25. Kochar J, Djoussé L, Gaziano JM. Breakfast cereals and risk of type 2 diabetes in the Physicians' Health Study I. *Obesity (Silver Spring)* 2007;15:3039–44.
26. Greenland S, Rothman KJ. Introduction to stratified analysis. In: Anonymous modern epidemiology. Philadelphia: Lippincott Williams & Wilkins; 1998. p. 258.
27. Inoue M, Tsugane S. Insulin resistance and cancer: epidemiological evidence. *Endocr Relat Cancer* 2012;19:F1–8.
28. Hillon P, Guiu B, Vincent J, Petit J. Obesity, type 2 diabetes and risk of digestive cancer. *Gastroenterol Clin Biol* 2010;34:529–33.
29. Steinberg FM, Bearden MM, Keen CL. Cocoa and chocolate flavonoids: implications for cardiovascular health. *J Am Diet Assoc* 2003;103:215–23.
30. Food and Drug Administration Department of Health and Human Services. "Title 21—food and drugs, chapter I, subchapter B—food for human consumption, part 163—cocoa products" (cited 1 Oct 2014). Available from: <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/CFRSearch.cfm?fr=163.130>.
31. Food Standard Agency. Guidance on the cocoa and chocolate products regulations 2003 (cited 1 Oct 2014). Available from: <http://www.food.gov.uk/sites/default/files/multimedia/pdfs/chocguidancejun2009.pdf>.