

多种RNA分子在肿瘤热疗研究中的进展

《现代肿瘤医学》[ISSN:1672-4992/CN:61-1415/R] 期数: 2019年12期 页码: 2185-2188 栏目: 综述 出版日期: 2019-05-08

Title: The progress of various RNA molecular in tumor hyperthermia research

作者: 刘佳佳¹; 蒙洁¹; 陈彻¹; 李莉¹; 王晶¹; 2

1.甘肃中医药大学临床医学院,甘肃 兰州 730000;2.甘肃省中医方药挖掘与创新转化重点实验室,甘肃 兰州 730000

Author(s): Liu Jiajia¹; Meng Jie¹; Chen Che¹; Li Li¹; Wang Jing¹; 2

1.School of Clinical Medicine,Gansu University of Chinese Medicine,Gansu Lanzhou 730000,China;2.The Key Laboratory of Traditional Chinese Herbs and Prescription Innovation and Transformation of Gansu Province,Gansu University of Chinese Medicine,Gansu Lanzhou 730000,China.

关键词: 肿瘤; 热疗; mRNA; 微小RNA; 长链RNA

Keywords: tumor; hyperthermia; mRNA; miRNA; lncRNA

分类号: R730.5

DOI: 10.3969/j.issn.1672-4992.2019.12.038

文献标识码: A

摘要: 随着热疗技术的发展和进步,热疗在肿瘤治疗领域发挥重大作用。RNA分子运载着基因和调控信息,并能反映细胞的实时状态,因此敏感特异性的RNA分子在临床精准医疗中显得尤为重要。关键RNA分子在肿瘤细胞热疗中的特异性表达,一方面可以揭示肿瘤热疗的分子机制,另一方面也作为肿瘤预后和诊断的重要标记。

Abstract: With the development of heat treatment technology, the problem such as temperature control, distribution and measurement has been resolved gradually, tumor hyperthermia will also play a major role. RNA for carrying information and can reflect the cell gene regulation and real-time state, using the sensitive specific RNA molecule markers in clinical precision medical is great important. The key RNAs show a specific expression in tumor cells treated by hyperthermia, which can reveal the molecular mechanism of tumor therapy, can also be used as biomarkers in tumor prognosis and diagnosis.

参考文献/REFERENCES

- [1] Wang XL,Sun RM.Mechanism and clinical study of tumor hyperthermia [J].Technology Information,2016,25(10):181-182. [王晓莉,孙瑞梅.肿瘤热疗的机理及临床研究 [J].科技资讯,2016,25(10):181-182.]
- [2] Ando K,Suzuki Y,Kaminuma T,et al.Tumor-specific CD8-positive T cell-mediated anti-tumor immunity is implicated in the antitumor effect of local hyperthermia [J].Int J Hyperthermia,2018,9(17):1-6.
- [3] Camacho CP,Lindsey SC,Melo MCC,et al.Measurement of calcitonin and calcitonin gene-related peptide mRNA refines the management of patients with medullary thyroid cancer and may replace calcitonin-stimulation tests [J].Thyroid,2013,23(3):308-314.
- [4] Hamam R,Hamam D,Alsaleh KA,et al.Circulating microRNAs in breast cancer:novel diagnostic and prognostic biomarkers [J].Cell Death Dis,2017,8(9):1-10.
- [5] Yang J,Ma DL,Fesler A,et al.Expression analysis of microRNA as prognostic biomarkers in colorectal cancer [J].Oncotarget,2016,8(32):53403-52411.
- [6] Wang HN,Crawford BM,Fales AM,et al.Multiplexed detection of microRNA biomarkers using SERS-based inverse molecular sentinel (iMS) nanoprobes [J].J Phys Chem C,2016,120(37):21047-21050.
- [7] Nandagopal L,Sonpavde G.Circulating biomarkers in bladder cancer [J].Bladder Cancer,2016,2(4):369-379.
- [8] Jiang N,Tian HH,Pan JC,et al.Application of circulating long-chain non-coding RNA as biomarker in tumor molecular diagnosis [J].J Bioengineering,2017,33(6):910-922. [江楠,田海华,潘金昌,等.循环长链非编码RNA作为生物标志物在肿瘤分子诊断中的应用 [J].生物工程学报,2017,33(6):910-922.]
- [9] Erbes T,Hirschfeld M,Waldeck S,et al.Hyperthermia-driven aberrations of secreted microRNAs in breast cancer in vitro [J].Int J Hyperthermia,2016,31(6):630-642.
- [10] Zhang P,Hong H,Sun X,et al.MicroRNA-10b regulates epithelial-mesenchymal transition by modulating KLF4/Notch1/E-cadherin in cisplatin-resistant nasopharyngeal carcinoma cells [J].Am J Cancer Res,2016,6(2):141-156.

- [11] Wendt MK,Tian M,Schiemann WP.Deconstructing the mechanisms and consequences of TGF- β -induced EMT during cancer progression [J]. *Cell and Tissue Research*,2012,347(1):85-101.
- [12] Lu YC,Chen YJ,Wang HM,et al.Oncogenic function and early detection potential of miRNA-10b in oral cancer as identified by microRNA profiling [J]. *Cancer Prev Res (Phila)*,2012,5(4):6207.
- [13] Ng L,Wan TM,Man JH,et al.Identification of serum miR-139-3p as a non-invasive biomarker for colorectal cancer [J]. *Oncotarget*,2017,8(16):27393-27400.
- [14] Xie D,Yin RT,Li KM,et al.Effect of hyperthermia combined with three-generation platinum drugs on ovarian cancer cell line SKOV3 [J]. *Journal of Sichuan University*,2014,45(4):606-611. [谢丹,尹如铁,李克敏,等.热疗联合三代铂类药物对卵巢癌细胞株SKOV3的影响 [J].*四川大学学报*,2014,45(4):606-611.]
- [15] Ryu H,Song IC,Choi YS,et al.ERCC1 expression status predicts the response and survival of patients with metastatic or recurrent cervical cancer treated via platinum-based chemotherapy [J]. *Medicine (Baltimore)*, 2017,96(51):e9402.
- [16] Wu LJ,Wang DL,Li L,et al.Sequential release of autophagy inhibitor and chemotherapeutic drug with polymeric delivery system for oral squamous cell carcinoma therapy [J]. *Mol Pharm*,2014,11(5):1662-1675.
- [17] Ju H,Mao L,Zhang L,et al.Ultrasound hyperthermia enhances chemo-sensitivity in oral squamous cell carcinoma by TRIF-mediated pathway [J]. *J Oral Pathol Med*,2018,10:12774.
- [18] Zhang XL,Shi HJ,Wang JP,et al.MicroRNA-218 is upregulated in gastric cancer after cytoreductive surgery and hyperthermic intraperitoneal chemotherapy and increases chemosensitivity to cisplatin [J]. *World J Gastroenterol*,2014,20(32):11347-11355.
- [19] Ruan Q,Fang ZY,Cui SZ,et al.Thermo-chemotherapy induced miR-218 upregulation inhibits the invasion of gastric cancer via targeting Gli2 and E-cadherin [J]. *Tumor Biol*,2015,36(8):5807-5814.
- [20] Pennacchi PC,Sandri S,Vicente AL,et al.Targeting the hedgehog transcription factors GLI1 and GLI2 restores sensitivity to vemurafenib-resistant human melanoma cells [J]. *Oncogene*,2017,36(13):1849-1861.
- [21] Li ZJ,Mack SC,Mak TH,et al.Evasion of p53 and G2/M checkpoints are characteristic of Hh-driven basal cell carcinoma [J]. *Oncogene*,2014,33(20):2674-2680.
- [22] Sadam H,Liivas U,Kazantseva A,et al.GLI2 cell-specific activity is controlled at the level of transcription and RNA processing:Consequences to cancer metastasis [J]. *Biochimica Et Biophysica Acta*,2016,1862(1):46-55.
- [23] Yu BQ,Gu DS,Zhang XL,et al.The role of GLI2-ABCG2 signaling axis for 5Fu resistance in gastric cancer [J]. *J Genet Genomics*,2017,8:375-383.
- [24] Xiao F,Liu B,Zhu QX.c-Jun N-terminal kinase is required for thermotherapy induced apoptosis in human gastric cancer cells [J]. *World J Gastroenterol*,2012,18(48):7348-7356.
- [25] Zeng JZ,Cai XR,Hao XB,et al.LncRNA FUNDC2P4 down-regulation promotes epithelial-mesenchymal transition by reducing E-cadherin expression in residual hepatocellular carcinoma after insufficient radiofrequency ablation [J]. *Int J Hyperthermia*,2018,34(6):802-811.
- [26] Bhatia S,Monkman J,Toh AKL,et al.Targeting epithelial-mesenchymal plasticity in cancer:clinical and preclinical advances in therapy and monitoring [J]. *Biochem J*,2017,474(19):3269-3306.
- [27] Huang CY,Hsieh MJ,Liu TC,et al.Correlation of E-cadherin gene polymorphisms and epidermal growth factor receptor mutation in lung adenocarcinoma [J]. *Cancer*,2018,15(8):765-770.
- [28] Kang K,Nam S,Kim B,et al.Inhibition of osteoclast differentiation by overexpression of NDRG2 in monocytes [J]. *Biochem Biophys Res Commun*,2015,468(4):611-616.
- [29] Yang CL,Zheng XL,Ye K,et al.NDRG2 suppresses proliferation,migration,invasion and epithelial-mesenchymal transition of esophageal cancer cells through regulating the AKT/XIAP signaling pathway [J]. *Int J Biochem Cell Biol*,2018,99:43-51.
- [30] Guo Y,Ma J,Wu L,et al.Hyperthermia-induced NDRG2 upregulation inhibits the invasion of human hepatocellular carcinoma via suppressing ERK1/2 signaling pathway [J]. *Plos One*,2013,8(4):e61079.
- [31] Min KJ,Um HJ,Kim JL,et al.The coffee diterpene kahweol enhances sensitivity to sorafenib in human renal carcinoma Caki cells through down-regulation of Mcl-1 and c-FLIP expression [J]. *Oncotarget*,2017,8(47):83195-83206.
- [32] Song X,Kim SY,Zhou Z,et al.Hyperthermia enhances mapatumumab-induced apoptotic death through ubiquitin-mediated degradation of cellular FLIP(long) in human colon cancer cells [J]. *Cell Death Dis*,2013,4:e577.
- [33] Resende MF,Vieira S,Chinen LT,et al.Prognostication of prostate cancer based on TOP2A protein and gene assessment:TOP2A in prostate cancer [J]. *J Transl Med*,2013,11:36.
- [34] Ludwig R,Teran FJ,Teichgraeber U,et al.Nanoparticle-based hyperthermia distinctly impacts production of ROS,expression of Ki-67, TOP2a, and TPX2, and induction of apoptosis in pancreatic cancer [J]. *Int J Nanomed*,2017,12:1009-1018.
- [35] Rampersand EN,Vujaskovic Z,Inman BA.Hyperthermia as a treatment for bladder cancer [J]. *Oncology*,2010,24(12):1149-1155.
- [36] Xie WY,Zhou XD,Yang J,et al.Inhibition of autophagy enhances heat-induced apoptosis in human non-small cell lung cancer cells through ER stress pathways [J]. *Arch Biochem Biophys*,2016,607:55-66.
- [37] Rezaie P,Khoei S,Khoei S,et al.Evaluation of combined effect of hyperthermia and ionizing radiation on cytotoxic damages induced by IUDR-loaded PCL-PEG-coated magnetic nanoparticles in spheroid culture of U87MG glioblastoma cell line [J]. *Int J Radiat Biol*,2018,94(11):1027-1037.
- [38] Yin PT,Shah BP,Lee KB.Combined magnetic nanoparticle-based microRNA and hyperthermia therapy to enhance apoptosis in brain cancer cells [J]. *Small*,2014,10(20):4106-4112.

[39] Wu JM,Liu T,Rios Z,et al.Heat shock proteins and cancer [J].Trends in Pharmacological Sciences,2017,38(3):226-256.

备注/Memo: National Natural Science Foundation of China(No.81760835);国家自然科学基金计划项目 (编号: 81760835) ;甘肃省自然科学基金 (编号: 1606RJZA192) ; 甘肃省卫生行业科研计划项目 (编号: GSWSKY2017-27) ;甘肃省高等学校基本科研业务费专项资金 (编号: 2013-005)

更新日期/Last Update: 1900-01-01